

Cell-autonomous axon growth of young motoneurons is triggered by a voltage-gated sodium channel

Andrea Wetzel, Sibylle Jablonka and Robert Blum*

Institute for Clinical Neurobiology; University of Würzburg; Würzburg, Germany

Spontaneous electrical activity preceding synapse formation contributes to the precise regulation of neuronal development. Examining the origins of spontaneous activity revealed roles for neurotransmitters that depolarize neurons and activate ion channels. Recently, we identified a new molecular mechanism underlying fluctuations in spontaneous neuronal excitability. We found that embryonic motoneurons with a genetic loss of the low-threshold sodium channel $Na_v1.9$ show fewer fluctuations in intracellular calcium in axonal compartments and growth cones than wild-type littermates. As a consequence, axon growth of $Na_v1.9$ -deficient motoneurons in cell culture is drastically reduced while dendritic growth and cell survival are not affected. Interestingly, $Na_v1.9$ function is observed under conditions that would hardly allow a ligand- or neurotransmitter-dependent depolarization. Thus, $Na_v1.9$ may serve as a cell-autonomous trigger for neuronal excitation. In this addendum, we discuss a model for the interplay between cell-autonomous local neuronal activity and local cytoskeleton dynamics in growth cone function.

In the developing nervous system, electrical activity is observed long before neurons are embedded in a synaptic network.^{1,2} Early electrical activity is an evolutionarily conserved phenomenon and regulates a broad spectrum of developmental processes, such as cell proliferation, neuronal differentiation, cell migration, neuronal survival and neurite growth, as well as the refinement of synaptic connections.² Excitation of

neurons before synapse formation is often termed as “spontaneous excitation” and in many cases the neuronal depolarization is accompanied by spontaneous influx of calcium ions to the neuronal cytosol. Investigating the molecular mechanisms underlying spontaneous calcium influx revealed two principle mechanisms of how spontaneous excitation is initiated. Either spontaneous excitation is ligand-dependent and caused by the non-synaptic release of transmitters such as glutamate or GABA; both act excitatory on immature neurons. Alternatively, excitability is part of a developmental program and thus a cell-autonomous feature of a young neuron. While there is an enormous progress in understanding the molecular mechanisms of ligand-dependent regulation of early electrical activity, the molecules regulating cell-autonomous excitability are barely defined.

In vertebrates, motoneurons exhibit spontaneous calcium transients while they are growing over long distances to the skeletal muscle.²⁻⁴ When embryonic motoneurons from rodents are cultured at low density the phenomenon of spontaneous excitation is preserved and spontaneous calcium transients are preferentially observed in axons and axonal growth cones.^{5,6} Spontaneous calcium transients in motoneurons contribute to axon elongation and presynaptic differentiation.⁵ This is of pathophysiological relevance. Motoneurons isolated from a mouse model for the motoneuron disease spinal muscular atrophy generate less spontaneous calcium transients and this causes deficits in axonal growth and growth cone differentiation.⁵

Keywords: motoneurons, sodium channel, axon growth, spontaneous excitation, $Na_v1.9$, local protein synthesis, spinal muscular atrophy

Submitted: 11/30/12

Revised: 12/05/12

Accepted: 12/06/12

<http://dx.doi.org/10.4161/chan.23153>

*Correspondence to: Robert Blum;
Email: blum_r@klinik.uni-wuerzburg.de

Addendum to: Subramanian N, Wetzel A, Dombert B, Yadav P, Havlicek S, Jablonka S, et al. Role of $Na_v1.9$ in activity-dependent axon growth in motoneurons. *Hum Mol Genet* 2012; 21:3655–67; PMID:22641814; <http://dx.doi.org/10.1093/hmg/dds195>

Cell-Autonomous Excitability in Embryonic Motoneurons

Research aiming to decipher the pathophysiology of the motoneuron disease spinal muscular atrophy gave deeper insights in the molecular mechanisms and function underlying growth-mediating spontaneous excitability.^{5,7,8}

Spinal muscular atrophy, commonly referred to as SMA, affects motoneurons in the anterior horn of the spinal cord and causes axonal defects and disturbed neuromuscular transmission, hence causing muscle weakness.⁹⁻¹¹ SMA is the most common genetic cause of infant mortality and is initiated by mutations in the Survival of Motor Neuron 1 (*SMN1*) gene.¹⁰

When embryonic motoneurons from a mouse model for a severe form of SMA (SMA type II) are analyzed in vitro, they show drastic changes in cell-autonomous functions. When these neurons are cultured on a laminin-isoform that supports axonal elongation, axon growth and growth cone size are reduced.^{5,7} This coincides with lower levels of β -actin mRNA and β -actin protein in growth cones.^{5,7} Recently, we confirmed that β -actin is locally synthesized in growth cones of motoneurons and this process is deregulated in *SMN*-deficient motoneurons.¹² Surprisingly, cytoskeletal defects in motoneuron growth cones cause defects in cell-surface clustering of the N-type calcium channel $Ca_v2.2$ and less local calcium influx events are observed.⁵ Defects in the typical cell-autonomous pathology of *SMN*-deficient motoneurons in vitro can partly be rescued. When motoneurons are treated with a cell membrane permeable cAMP analog, distal actin mRNA and actin protein levels are increased, $Ca_v2.2$ levels are augmented in growth cones, and subsequently spontaneous calcium influx frequencies are normalized.⁵ Thus, local translation of β -actin is most likely upstream of the mechanism for proper N-type channel clustering in distal axons.

Spontaneous calcium transients in motoneurons are observed in cells without synaptic contact, even under rapid perfusion with defined artificial cerebrospinal fluid solutions (Fig. 1A shows some examples, visualized at a slow time-scale).⁶ However, the N-type calcium

channel $Ca_v2.2$ and the P/Q-type channel $Ca_v2.1$, both reside at presynaptic sites in motoneurons, normally require a strong depolarization for activation.¹³ This predicts a trigger molecule with the following features: (1) expressed in motoneurons, (2) present in axons and growth cones, (3) low activation threshold, (4) spontaneous opening close to resting membrane potentials, (5) mediator of excitatory ion influx.

To characterize this trigger molecule was the aim of our study by Subramanian and Wetzel et al. (2012).⁶ We investigated whether low threshold voltage-gated sodium channels mediate cell-autonomous excitability in young motoneurons.

As a first test, wild-typic motoneurons were cultured in the presence of low concentrations of pore blockers of voltage-gated sodium channels. Indeed, after several days in culture axonal elongation of motoneurons was reduced when motoneurons were treated with low amounts of saxitoxin (STX) and tetrodotoxin (TTX). Dendritic growth and motoneuron survival remained unaffected.

Voltage-gated sodium channels are key proteins for neuronal excitation and action potential initiation. However, as discussed above, spontaneous electrical activity in motoneurons is also a local event, and is therefore unlikely to be mediated by ligand-dependent initiation of action potentials. This raised the question which voltage-gated sodium channel is responsible for fluctuations in local excitability close to the resting membrane potential.

A Low-Threshold Voltage-Gated Sodium Channel Triggers Cell-Autonomous Excitation

The family of VGSCs comprises nine members, classified as $Na_v1.1$ to $Na_v1.9$.¹⁴ In comparison to the other Na_v -channels, the TTX-insensitive channel $Na_v1.9$ has unique electrophysiological properties. $Na_v1.9$ currents were classified in nociceptive C-type sensory neurons in dorsal root ganglia, and in sensory neurons of the myenteric plexus.¹⁵⁻¹⁸ In these cells, the channel opens at very low membrane potentials, mediates spontaneous excitation and shows slow activation and

inactivation kinetics.^{15,16,18,19} Thus, $Na_v1.9$ is thought to act as a regulator of sub-threshold electrogenesis.^{15,16,18-20}

These specific features of $Na_v1.9$ prompted us to investigate whether $Na_v1.9$ is involved in cell-autonomous, activity-dependent axon growth in motoneurons. When motoneurons are cultured from $Na_v1.9$ knockout mice, axons are much shorter than axons from wild-type mice (Fig. 1B).⁶ This reduced axonal elongation in $Na_v1.9$ -deficient motoneurons correlates with reduced numbers of spontaneous Ca^{2+} transients. Highest numbers of spontaneous calcium influx events are typically observed in growth cones and distal axons, but “global” as well as “local” calcium transients are strongly reduced in $Na_v1.9$ -deficient motoneurons.⁶

Expression of $Na_v1.9$ in Young Motoneurons

In DRG neurons, $Na_v1.9$ expression levels are high²¹ and $Na_v1.9$ specific currents are confirmed with the help of loss-of-function and gain-of-function experiments.¹⁷ PCR data indicate that $Na_v1.9$ is widely expressed in excitable cells and neuroblastoma cell lines,^{22,23} but, in contrast to data from DRG neurons or enteric sensory neurons, these expression data are not supported by a verification of voltage-dependent $Na_v1.9$ currents.

Earlier data showed no²¹ or weak expression levels^{22,23} for $Na_v1.9$ in the spinal cord. Therefore, we tested whether there are substantial differences in the expression levels of $Na_v1.9$ by comparing DRG neurons with the embryonic spinal cord, where young motoneurons reside, with motoneurons in culture and with in vivo motoneurons.

Quantitative PCR confirmed an exceptionally high expression rate for $Na_v1.9$ in DRG neurons⁶ and revealed $Na_v1.9$ expression during early neuronal development. $Na_v1.9$ expression in DRG tissue starts already at embryonic day 12, increases to 1.150 $Na_v1.9$ transcripts per ng RNA at E18 and becomes very high in older mice (e.g., P19, 2.700 copies/ng RNA).⁶ In spinal cord tissue, $Na_v1.9$ transcription gradually increases from E12 and peaks at late embryonic stages. At E18,




Figure 1. Na_v1.9 triggers spontaneous calcium influx in motoneurons and contributes to activity-dependent axon growth. **(A)** Characteristics of spontaneous Ca²⁺ transients in young cultured motoneurons from wild type mice. Motoneurons were cultivated for 5 d in vitro and afterwards loaded with the calcium indicator Oregon green BAPTA1. In live cell imaging experiments changes in fluorescence intensity ($\Delta f/f_0$) are illustrated from different cells. Calcium transients show variability in frequency and amplitude (mn1–3) and they change between on and off conformation within single cells (mn4 and 5). **(B)** Axon elongation is reduced in motoneurons from Na_v1.9 knockout mice. Representative motoneurons from wild-type and Na_v1.9 knockout mice are shown. Cells were cultured for 7 d in vitro. After anti- α -tubulin staining cells were analyzed by standard laser scanning microscopy. Bar: 40 μ m. **(C)** Representative motoneuron at DIV5 is loaded with the calcium indicator Oregon green BAPTA1. **(D)** Calcium transients of roi1–roi4 as indicated in **(C)**. Note here that local spontaneous calcium influx activity is observed in the growth cone only and does not spread to neighboring axonal regions. **(E)** Model for the interplay between local regulation of actin dynamics and ion channel gating in the plasma membrane in the growth cone of embryonic motoneurons. Actin mRNA is anterogradely transported from the soma into the axon and local synthesis of β -actin takes place in the growth cone. Axonal elongation is an activity-dependent process. The voltage-gated sodium channel Na_v1.9 opens spontaneously near the resting membrane potential and the influx of sodium leads to a depolarization cascade to gate clusters of the N-type voltage-gated calcium channels Ca_v2.2. The influx of free calcium supports axon growth by an unknown mechanism. The receptor tyrosine kinases TrkB might modulate the open probability of Na_v1.9 and/or leads to an opening of non-selective TrpC ion channels. Data were acquired during our previous study by Subramanian et al. (2012) with materials and methods described there in detail.⁶

Na_v1.9 transcription in the spinal cord is \sim 30-fold lower (37,3 copies/ng RNA) compared with DRG neurons.⁶ Na_v1.9 transcripts are also found in young motoneurons in vivo. Laser-assisted microdissection of young motoneurons from the anterior horn of the spinal cord revealed a low abundance of Na_v1.9 transcripts, approximately 80x lower than in DRG neurons.⁶

This indicates that even low expression levels of Na_v1.9 are sufficient to augment neuronal excitability, to trigger spontaneous voltage-dependent calcium influx,

and to support activity-dependent axon growth, at least in cultured embryonic motoneurons.

Early Sorting of Na_v1.9 Protein to Axons and Growth Cones

In young motoneurons, local Na_v1.9-dependent calcium signals are preferentially measured in axons and axonal growth cones. Therefore, Na_v1.9 protein should also be present in local regions within axons and growth cones. In order to identify the Na_v1.9 protein, an antibody

was generated and tested on DRG tissue samples of the mouse. Western blots labeled endogenous mouse Na_v1.9 protein from DRG preparations and recombinant Na_v1.9 from different species at two bands of 180 kDa and approximately 280 kDa.⁶ This typical western blot pattern is lost in preparations from a Na_v1.9 knock out mouse model. In this specific Na_v1.9 knock out mouse, introduced by Östman et al.,¹⁷ the S4 voltage sensor segment of domain I in exon 4 and 5 of the Na_v1.9 gene was replaced by a neomycin resistance cassette, thus causing nonsense

mutations in exon 6. Immunofluorescence labeling of Na_v1.9 combined with super resolution imaging using stimulated emission depletion microscopy (STED) revealed that Na_v1.9 resides preferentially in axons and growth cones. Anti-Na_v1.9 immunolabels are distributed non-uniformly along the axon and are locally enriched at distinct axonal regions.⁶ The spatial restriction of local calcium influx and the typical uneven distribution of Na_v1.9 protein predict mechanisms that help to localize Na_v1.9. Perhaps, the channel maintains this localization by an attachment to the cytoskeleton or by a diffusion barrier in axonal microdomains.

Na_v1.9 Might Act within Axonal Subcompartments

As discussed above, local translation of β-actin mRNA in growth cones is essential for proper growth cone function and is linked with activity-dependent axon growth.^{5,7,12} The actin cytoskeleton in growth cones is essential for cell surface clustering of N-type calcium channels in distal axons, and activation of these channels is then triggered by Na_v1.9 activity (Fig. 1E).^{5,6}

Local calcium transients in axonal regions are often observed within small, distinct axonal regions. **Figure 1C and D** shows an example of a “local” calcium transient, here imaged with the help of a high-affinity Ca²⁺ indicator at 10 Hz, in a single confocal plane. In this example, the calcium transient is only observed in a small torpedo-like growth cone of a cultured motoneuron and does not spread to adjacent axonal regions; hence it forms a microdomain or even a restricted axonal subcompartment. Cytoskeleton remodeling is upstream of increased local excitability in motoneurons. Rescue of β-actin dynamics in SMN-deficient motoneurons also rescues local calcium influx and activity-dependent axonal elongation. High motility at growth cones might be accompanied by a high elasticity of the lipid bilayer. The high motility of the lipid bilayer and the cytoskeleton in the “healthy” growth cone of motoneurons might increase the activation sensitivity of Na_vs. For instance, Na_v1.5 is sensitive to mechanical stimuli and this causes shifts

in activation and inactivation properties due to mechanical modulation of the voltage sensors.²⁴ Therefore, it is conceivable that proper cytoskeleton dynamics may directly influence the region-specific and local excitability of voltage-gated sodium channels in the growth cone.

Critical Aspects

There is a major critical aspect in our study. Na_v1.9 function was monitored indirectly and the electrophysiological properties of the channel have not been defined in motoneurons yet. Electrophysiological studies may help us to solve a critical issue regarding the use of pore blockers of voltage-gated sodium channels as growth inhibiting compounds. Axonal elongation of motoneurons is reduced in presence of 1–10 nM saxitoxin, more than 50 nM tetrodotoxin is necessary to get a comparable effect.⁶ This indicates that STX has a higher affinity to Na_v1.9 than TTX. In line with this finding, it has already been described that low doses of STX reduce neurotrophin-induced neurite growth of young cultured hippocampal neurons of the rat.²⁵ In addition, reconstitution experiments mimicking neurotrophin-induced excitation in hippocampal neurons indicated that 10 nM STX inhibits neurotrophin-induced activation of Na_v1.9, while 50 nM TTX does not.^{22,26}

Electrophysiological recordings with DRG neurons argue against the idea of a high specificity of saxitoxin to Na_v1.9. In DRGs, STX and TTX are both quite ineffective on persistent sodium currents attributed to the Na_v1.9 channel.²⁰ Dose-response experiments regarding voltage-dependent activation currents of Na_v1.9 revealed an inhibitor constant (K_i) value for TTX of 39 ± 9 μM.¹⁶ Thus, it remains a puzzle why 10 nM STX has such a strong inhibitory effect on activity-dependent motoneuron axonal elongation⁶ and neurite growth of hippocampal neurons.^{22,25} When motoneurons are treated with 10 nM STX by fast perfusion with artificial cerebrospinal fluid (ACSF), local and global spontaneous calcium transients are rapidly blocked.⁶ This effect is reversible. STX can be washed out by continuous perfusion with ACSF and spontaneous calcium influx recovers after

10–20 min.⁶ STX and TTX seem to bind to all members of the Na_v1 family, and the differences in biological responses on neurite growth observed with these channel blockers at very low concentrations are not easy to explain. It may be that the growth-inhibiting effect of STX is observed, because it blocks sodium channels that are downstream of Na_v1.9 and have a high-affinity to STX. According to its properties, Na_v1.2²⁷ is a good candidate for being this sodium channel in young neurons, but for this we have no proof, yet.

In addition, it will be helpful to define the properties and constitution of these commercially available toxin preparations by independent methods. For this, a reconstitution system enabling the investigation of voltage-dependent and pharmacological properties of Na_v1.9 by heterologous expression, e.g., in HEK293 cells or another reconstitution cell line, would be helpful, but such an in vitro system is still a major issue in Na_v1.9 research.

Ion Channel Interplay in Growth Cones of Motoneurons

Na_v1.8 is the concomitant partner of Na_v1.9 in DRG neurons and seem not to be expressed in embryonic motoneurons⁶ or healthy adult motor nerves.²⁸ We found 20-fold higher Na_v1.5 vs. Na_v1.9 expression levels in embryonic motoneurons.⁶ Na_v1.5 has a quite low activation threshold¹³ and would be an ideal physiological partner to potentiate Na_v1.9-mediated excitability. This led us to speculate that Na_v1.9 is first activating Na_v1.5, which then initiates a rapid depolarization and a fast voltage-dependent gating of Ca_v2.2 (Fig. 1E). During live imaging of spontaneous activity in motoneurons (minutes to hours), there are always phases of very low or very high activity and motoneurons switch between stages of no or low activity and stages of high activity (Fig. 1A). This raises the question how phases of high activity are stimulated. We have no evidence that increased levels of excitability in motoneurons are predictable or whether the channel opening is a stochastic event. In DRG neurons, Na_v1.9 action is regulated by intracellular signaling cascades.^{15,17} In motoneurons, the tropomyosin receptor kinase B (TrkB)

might be a good candidate for regulating phases of higher vs. lower excitability. These receptors are expressed on cultured embryonic motoneurons and activation of these receptors by the neurotrophin brain-derived neurotrophic factor (BDNF) supports motoneuron survival.²⁹ In neurons of the hippocampus BDNF and TrkB are modulators of Na_v1.9 activity and are claimed to be involved in instructive effects of BDNF in synaptic plasticity.^{22,26,30,31} To our opinion, it is unlikely that BDNF activates TrkB and induces Na_v1.9 activity and subsequent N-type channel gating in motoneurons. More likely is a model that includes signaling mechanisms such as transactivation of an intracellular domain of TrkB^{32,33} and a subsequent increase of the open probability of Na_v1.9 (Fig. 1E).

It is also important to consider a potential function of transient receptor potential channels (Trp) in motoneuron growth cones. In young hippocampal neurons, TrpC5 channels are localized to neurite growth cones, are calcium permeable and regulate neurite extension and growth cone motility.⁵ TrpC family members have also been identified in growth cones of cultured embryonic motoneurons.⁵ It may be that kinase-active TrkB first acts via PLC γ recruitment to start a subsequent increase in excitability by a non-selective ion influx through TrpC channels. Such a mechanism was first observed in pontine neurons³⁴ and TrpC activity may offer a weak and local increase in excitability, Na_v1.9 activation and subsequent depolarization that gates voltage-dependent calcium influx.

In the last decade, investigating cell-autonomous growth mechanisms of motoneurons has offered important insights in the function of mammalian ion channels in activity-dependent growth. However, even though there is a concept on how ion channels in growth cones may interact, the molecular mechanism of the pivotal physiological role of increased free calcium in growth cones remains elusive.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

This work was supported by grants from the Deutsche Forschungsgemeinschaft. We thank Samira Samtleben, Michael Sendtner and Erich Buchner for helpful comments on this manuscript.

References

- Gu X, Spitzer NC. Distinct aspects of neuronal differentiation encoded by frequency of spontaneous Ca²⁺ transients. *Nature* 1995; 375:784-7; PMID:7596410; <http://dx.doi.org/10.1038/375784a0>
- Spitzer NC. Electrical activity in early neuronal development. *Nature* 2006; 444:707-12; PMID:17151658; <http://dx.doi.org/10.1038/nature05300>
- Hanson MG, Landmesser LT. Characterization of the circuits that generate spontaneous episodes of activity in the early embryonic mouse spinal cord. *J Neurosci* 2003; 23:587-600; PMID:12533619
- O'Donovan MJ, Landmesser L. The development of hindlimb motor activity studied in the isolated spinal cord of the chick embryo. *J Neurosci* 1987; 7:3256-64; PMID:3668626
- Jablonka S, Beck M, Lechner BD, Mayer C, Sendtner M. Defective Ca²⁺ channel clustering in axon terminals disturbs excitability in motoneurons in spinal muscular atrophy. *J Cell Biol* 2007; 179:139-49; PMID:17923533; <http://dx.doi.org/10.1083/jcb.200703187>
- Subramanian N, Wetzell A, Dombert B, Yadav P, Havlicek S, Jablonka S, et al. Role of Na(v)1.9 in activity-dependent axon growth in motoneurons. *Hum Mol Genet* 2012; 21:3655-67; PMID:22641814; <http://dx.doi.org/10.1093/hmg/dds195>
- Rossoll W, Jablonka S, Andreassi C, Kröning AK, Karle K, Monani UR, et al. Snn, the spinal muscular atrophy-determining gene product, modulates axon growth and localization of beta-actin mRNA in growth cones of motoneurons. *J Cell Biol* 2003; 163:801-12; PMID:14623865; <http://dx.doi.org/10.1083/jcb.200304128>
- Rossoll W, Kröning AK, Ohndorf UM, Steegborn C, Jablonka S, Sendtner M. Specific interaction of Snn, the spinal muscular atrophy determining gene product, with hnRNP-R and gry-rbp/hnRNP-Q: a role for Snn in RNA processing in motor axons? *Hum Mol Genet* 2002; 11:93-105; PMID:11773003; <http://dx.doi.org/10.1093/hmg/dd1193>
- McGovern VL, Gavriliina TO, Beattie CE, Burghes AH. Embryonic motor axon development in the severe SMA mouse. *Hum Mol Genet* 2008; 17:2900-9; PMID:18603534; <http://dx.doi.org/10.1093/hmg/ddn189>
- Monani UR. Spinal muscular atrophy: a deficiency in a ubiquitous protein; a motor neuron-specific disease. *Neuron* 2005; 48:885-96; PMID:16364894; <http://dx.doi.org/10.1016/j.neuron.2005.12.001>
- Murray LM, Lee S, Bäumer D, Parson SH, Talbot K, Gillingwater TH. Pre-symptomatic development of lower motor neuron connectivity in a mouse model of severe spinal muscular atrophy. *Hum Mol Genet* 2010; 19:420-33; PMID:19884170; <http://dx.doi.org/10.1093/hmg/ddp506>
- Rathod R, Havlicek S, Frank N, Blum R, Sendtner M. Laminin induced local axonal translation of β -actin mRNA is impaired in SMN-deficient motoneurons. *Histochem Cell Biol* 2012; 138:737-48; PMID:22810847; <http://dx.doi.org/10.1007/s00418-012-0989-1>
- Catterall WA, Perez-Reyes E, Snutch TP, Striessnig J. International Union of Pharmacology. XLVIII. Nomenclature and structure-function relationships of voltage-gated calcium channels. *Pharmacol Rev* 2005; 57:411-25; PMID:16382099; <http://dx.doi.org/10.1124/pr.57.4.5>
- Catterall WA, Goldin AL, Waxman SG. International Union of Pharmacology. XLVII. Nomenclature and structure-function relationships of voltage-gated sodium channels. *Pharmacol Rev* 2005; 57:397-409; PMID:16382098; <http://dx.doi.org/10.1124/pr.57.4.4>
- Baker MD, Chandra SY, Ding Y, Waxman SG, Wood JN. GTP-induced tetrodotoxin-resistant Na⁺ current regulates excitability in mouse and rat small diameter sensory neurones. *J Physiol* 2003; 548:373-82; PMID:12651922; <http://dx.doi.org/10.1113/jphysiol.2003.039131>
- Cummins TR, Dib-Hajj SD, Black JA, Akopian AN, Wood JN, Waxman SG. A novel persistent tetrodotoxin-resistant sodium current in SNS-null and wild-type small primary sensory neurons. *J Neurosci* 1999; 19:RC43; PMID:10594087
- Östman JA, Nassar MA, Wood JN, Baker MD. GTP up-regulated persistent Na⁺ current and enhanced nociceptor excitability require Nav1.9. *J Physiol* 2008; 586:1077-87; PMID:18096591; <http://dx.doi.org/10.1113/jphysiol.2007.147942>
- Rugiero F, Mistry M, Sage D, Black JA, Waxman SG, Crest M, et al. Selective expression of a persistent tetrodotoxin-resistant Na⁺ current and Nav1.9 subunit in myenteric sensory neurons. *J Neurosci* 2003; 23:2715-25; PMID:12684457
- Herzog RI, Cummins TR, Waxman SG. Persistent TTX-resistant Na⁺ current affects resting potential and response to depolarization in simulated spinal sensory neurons. *J Neurophysiol* 2001; 86:1351-64; PMID:11553682
- Priest BT, Murphy BA, Lindia JA, Diaz C, Abbadie C, Ritter AM, et al. Contribution of the tetrodotoxin-resistant voltage-gated sodium channel Nav1.9 to sensory transmission and nociceptive behavior. *Proc Natl Acad Sci U S A* 2005; 102:9382-7; PMID:15964986; <http://dx.doi.org/10.1073/pnas.0501549102>
- Dib-Hajj SD, Tyrrell L, Black JA, Waxman SG. NaN, a novel voltage-gated Na channel, is expressed preferentially in peripheral sensory neurons and down-regulated after axotomy. *Proc Natl Acad Sci U S A* 1998; 95:8963-8; PMID:9671787; <http://dx.doi.org/10.1073/pnas.95.15.8963>
- Blum R, Kafitz KW, Konnerth A. Neurotrophin-evoked depolarization requires the sodium channel Na_v1.9. *Nature* 2002; 419:687-93; PMID:12384689; <http://dx.doi.org/10.1038/nature01085>
- Jeong SY, Goto J, Hashida H, Suzuki T, Ogata K, Masuda N, et al. Identification of a novel human voltage-gated sodium channel alpha subunit gene, SCN12A. *Biochem Biophys Res Commun* 2000; 267:262-70; PMID:10623608; <http://dx.doi.org/10.1006/bbrc.1999.1916>
- Beyder A, Rae JL, Bernard C, Strege PR, Sachs F, Farrugia G. Mechanosensitivity of Nav1.5, a voltage-sensitive sodium channel. *J Physiol* 2010; 588:4969-85; PMID:21041530; <http://dx.doi.org/10.1113/jphysiol.2010.199034>
- Kafitz KW, Lepier A, Thoenen H, Konnerth A. Saxitoxin-sensitivity of neurotrophin-induced rapid excitation and neurite growth. *Pflugers Arch* 2001; 441:R125, O2-2
- Blum R, Konnerth A. Neurotrophin-mediated rapid signaling in the central nervous system: mechanisms and functions. *Physiology (Bethesda)* 2005; 20:70-8; PMID:15653842; <http://dx.doi.org/10.1152/physiol.00042.2004>
- West JW, Scheuer T, Maechler L, Catterall WA. Efficient expression of rat brain type IIA Na⁺ channel alpha subunits in a somatic cell line. *Neuron* 1992; 8:59-70; PMID:1309650; [http://dx.doi.org/10.1016/0896-6273\(92\)90108-P](http://dx.doi.org/10.1016/0896-6273(92)90108-P)

28. Ulzheimer JC, Peles E, Levinson SR, Martini R. Altered expression of ion channel isoforms at the node of Ranvier in P0-deficient myelin mutants. *Mol Cell Neurosci* 2004; 25:83-94; PMID:14962742; <http://dx.doi.org/10.1016/j.mcn.2003.09.015>
29. Sendtner M, Holtmann B, Kolbeck R, Thoenen H, Barde YA. Brain-derived neurotrophic factor prevents the death of motoneurons in newborn rats after nerve section. *Nature* 1992; 360:757-9; PMID:1465147; <http://dx.doi.org/10.1038/360757a0>
30. Kovalchuk Y, Hanse E, Kafitz KW, Konnerth A. Postsynaptic Induction of BDNF-Mediated Long-Term Potentiation. *Science* 2002; 295:1729-34; PMID:11872844; <http://dx.doi.org/10.1126/science.1067766>
31. Lang SB, Stein V, Bonhoeffer T, Lohmann C. Endogenous brain-derived neurotrophic factor triggers fast calcium transients at synapses in developing dendrites. *J Neurosci* 2007; 27:1097-105; PMID:17267564; <http://dx.doi.org/10.1523/JNEUROSCI.3590-06.2007>
32. Lee FS, Chao MV. Activation of Trk neurotrophin receptors in the absence of neurotrophins. *Proc Natl Acad Sci U S A* 2001; 98:3555-60; PMID:11248116; <http://dx.doi.org/10.1073/pnas.061020198>
33. Wiese S, Jablonka S, Holtmann B, Orel N, Rajagopal R, Chao MV, et al. Adenosine receptor A2A-R contributes to motoneuron survival by transactivating the tyrosine kinase receptor TrkB. *Proc Natl Acad Sci U S A* 2007; 104:17210-5; PMID:17940030; <http://dx.doi.org/10.1073/pnas.0705267104>
34. Li HS, Xu XZ, Montell C. Activation of a TRPC3-dependent cation current through the neurotrophin BDNF. *Neuron* 1999; 24:261-73; PMID:10677043; [http://dx.doi.org/10.1016/S0896-6273\(00\)80838-7](http://dx.doi.org/10.1016/S0896-6273(00)80838-7)