

ORIGINAL ARTICLE

Prescription of antibiotics in the medical care of newly arrived refugees and migrants

Evelyn Kleinert¹  | Nele Hillermann¹ | Alexandra Jablonka^{2,3} |
Christine Happle⁴ | Frank Müller¹ | Anne Simmenroth⁵

¹Department of General Practice, University Medical Center Göttingen/Georg-August-University, Göttingen, Germany

²Department of Clinical Immunology and Rheumatology, Hannover Medical School, Hannover, Germany

³German Center for Infection Research, partner site Hanover-Braunschweig, Hannover, Germany

⁴Department of Pediatrics, Neonatology, and Allergology, Hannover Medical School, Hannover, Germany

⁵Department of General Practice, University Medical Center Würzburg, Würzburg, Germany

Correspondence

Evelyn Kleinert, Department of General Practice, University Medical Center Göttingen/Georg-August-University, Humboldtallee 38, 37073 Göttingen, Germany.
Email: evelyn.kleinert@med.uni-goettingen.de

Funding information

European Social Fund; German Centre for Infection Research at Hannover Medical School; Land Niedersachsen; Medizinischen Hochschule Hannover; Robert Bosch Stiftung

Abstract

Purpose: Unnecessary and inappropriate use of antibiotics is a widespread problem in primary care. However, current data on the care of refugees and migrants in initial reception centers is pending. This article provides data on prescription frequencies of various antibiotics and associated diagnoses.

Methods: In this retrospective observational study, patient data of 3255 patients with 6376 medical contacts in two initial reception centers in Germany were analyzed. Patient data, collected by chart review, included sociodemographic characteristics, diagnoses, and prescriptions. Antibiotic prescription behavior and corresponding physician-coded diagnoses were analyzed.

Results: Nineteen percent of all patients in our study received systemic antibiotics during the observation period, with children below the age of 10 years receiving antibiotics most frequently (24%). The most commonly prescribed antibiotics were penicillins (65%), macrolides (12%), and cephalosporins (7%). The most frequent diagnoses associated with antibiotic prescription were acute tonsillitis (26%), bronchitis (21%), infections of the upper respiratory tract (14%), and urinary tract infections (10%). In case of acute bronchitis 74% of the antibiotic prescriptions were probably not indicated. In addition, we found a significant number of inappropriate prescriptions such as amoxicillin for tonsillitis (67%), and ciprofloxacin and cotrimoxazol for urinary tract infections (49%).

Conclusion: Regarding inappropriate prescription of antibiotics in refugee healthcare, this study shows a rate ranging from 8% for upper respiratory tract infections to 75% for acute bronchitis. Unnecessary use of antibiotics is a global problem contributing to gratuitous costs, side effects, and antimicrobial resistance. This research contributes to the development of stringent antibiotic stewardship regimens in the particularly vulnerable population of migrants and refugees.

KEYWORDS

antibiotic prescription, antimicrobial resistance, inappropriate prescription, pharmacoepidemiology, primary healthcare, refugee healthcare, viral infection

Evelyn Kleinert and Nele Hillermann are joint first authors, and Frank Müller and Anne Simmenroth are joint senior authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Pharmacoepidemiology and Drug Safety* published by John Wiley & Sons Ltd.

1 | INTRODUCTION

The annual Global Trends Report of the United Nations High Commissioner for Refugees (UNHCR) shows that at the end of 2018, 70.8 million people worldwide were forcibly displaced.¹ Since the beginning of 2015, more than 1.6 million people applied for asylum in Germany.² This high number of refugees and their integration, particularly into the German healthcare system, poses a great challenge. In a 2016 questionnaire study, 86% of medical practitioners declared that they treated patients who had arrived in Germany as asylum seekers since 2015.³ Common reasons of refugee patients for seeking treatment are respiratory and neuropsychological diseases, gastrointestinal disorders, diseases of the musculoskeletal system, and skin complaints. Studies showed that the reasons for medical encounters in refugee patients differ little from those of local patients.^{4,5}

In Germany, physicians prescribed 368 million defined daily doses (DDD) of systemic antibiotics in 2016,⁶ which is equivalent to 14.14 DDD per 1000 inhabitants per day.⁷ Systemic antibiotics are more often prescribed to children, especially at preschool age, than to all other age groups.⁷⁻⁹ Generally, women receive more antibiotics than men.^{8,9} A key reason for this finding can be seen in the fact that some infectious diseases, for example, urinary tract infections (UTI), are more common in women due to the anatomy of the female urinary tract. Additionally, women seek medical care more often than men and are therefore more likely to receive antibiotic prescriptions.^{10,11}

Antibiotics are frequently prescribed to patients with respiratory diseases, even though antibiotics are not indicated because most respiratory infections are of viral origin. Moreover, antibiotics do not reduce disease duration.¹²⁻¹⁴ Doctors reported that their reasons for prescribing not indicated antibiotics were most commonly due to patient demands and also that the demand arose in urgent care settings.¹⁵ Frequently prescribed antibiotics are penicillins and aminopenicillins, cephalosporins, tetracyclines, and quinolones.^{6,7} Inadequate use of antibiotics causes antimicrobial resistance, which significantly limits treatment options in the treatment of severe infections. Antibiotic resistance is considered one of the largest threats to global health. One of the most important strategies worldwide is responsible and prudent use of systemic antibiotics, realized by thorough guidelines and so-called antibiotic stewardship.^{16,17} Antibiotic stewardship programs subsume different measures for targeted and adequate antibiotic therapy and for avoiding unnecessary prescriptions. These can include training of physicians, patient education and empowerment, and joint patient visiting rounds with microbiologists and pharmacists, as well as tracking and reporting the use and prescription of antibiotics. Typically, interprofessional antibiotic stewardship units are established in hospitals for this purpose.^{18,19}

Several studies have demonstrated that people with low educational status are particularly unaware of the mode of functioning of antibiotics and at risk of antibiotic resistance.²⁰⁻²³ This might be an important aspect in refugee healthcare considering that 30% of asylum seekers in Germany have only attended primary school or have never attended school at all.²⁴ Additionally, language and cultural barriers complicate doctor-patient relationship and general healthcare

Key Points

- We observed a high rate of inappropriate prescription of antibiotics in a large current cohort of refugees newly arriving in Germany.
- Especially in the context of common viral infection such as acute bronchitis, the observed high rate of prescription of antibiotics appears to be unnecessary.
- To avoid unnecessary costs, side effects, and the emergence of antimicrobial resistance, healthcare providers caring for refugees must be aware of this problem and be trained in antibiotic stewardship.
- In our setting of healthcare provision in initial reception centers, cultural and language barriers as well as changing medical staff with limited time resources may have contributed to the high rate of prescription of antibiotics.

provision.²⁵⁻²⁸ Still, there is a gap in research on antibiotic prescribing in healthcare facilities for refugees and asylum seekers.

This study collects data on the prescription of antibiotics in primary healthcare centers in two German initial reception facilities for refugees and asylum seekers. We provide an overview of antibiotic prescription behavior in primary refugee healthcare in Germany in the context of diagnoses and demographic characteristics.

2 | METHODS

2.1 | Study population

In this retrospective observational study, all patient charts from two on-site primary healthcare wards of the initial reception facilities Celle and Friedland in Germany were analyzed. In total, we included data from 3255 patients with 6376 medical contacts in the study. The first cohort we analyzed consists of 1017 refugee patients (with a total of 2282 encounters) who sought medical care between September and December 2015 at the medical ward of the refugee reception facility in Celle (in Northern Germany). This facility was a temporary shelter that was built to cope with the large number of refugees during the 2015 refugee crisis and was closed with the decrease in numbers in mid-2016. Parts of this cohort have been described previously.^{29,30} The second cohort consists of 2238 refugee patients who had 4094 encounters in total and arrived at the reception facility in Friedland (in Central Germany) between August 2017 and August 2018. Other aspects of this cohort and their typical healthcare needs have been published previously.⁵ The reception facilities provided primary medical care by on-site medical wards to all residents of the facilities according to the German asylum seeker benefits law (Asylbewerberleistungsgesetz), which restricts medical services in comparison to those covered by the statutory health insurance. In brief, patients receive primary care to meet basic healthcare needs

and can be referred to a specialist or hospital for treatment if necessary. Each on-site ward was staffed by emergency medical personnel 24 h per day, supplemented by consultation hours by various doctors on each weekday.

2.2 | Data collection and analysis

Patient data were extracted from electronic (Celle) or paper-based (Friedland) health records, including age, sex, refugee status (asylum seeker or resettlement refugee), and country of origin. Whenever available, diagnosis coding according to the International Statistical Classification of Diseases and Related Health Problems (ICD-10) was also recorded for each consultation. Prescribed drugs were coded using the Anatomical Therapeutic Chemical Classification System (ATC). Coding was performed by a late-stage medical student and supervised by experienced physicians. Random audits were performed to ensure data quality. Because patients often consulted their doctors with several health concerns (e.g., tonsillitis, migraine, and depression), the antibiotic-relevant diagnosis (tonsillitis) was identified manually (see File S1).

Prescriptions were coded using different ATC coding levels including anatomical main group (first level), therapeutic/pharmacological subgroup (third level), and chemical substance (fifth level). In this study, we only evaluate the prescription of systemic antibiotics, therefore antibiotics for topical use were not taken into account. The statistical software package IBM SPSS statistics 25 (IBM, Armonk, NY) was used for all analyses. Descriptive statistical methods were used to describe the patient cohort and their diagnoses and prescriptions. Differences in age and sex were tested for significance by using the Mann-Whitney *U* test (for metric non-normally distributed variables) and the Pearson χ^2 test (for categorical variables).

Our findings are discussed in consideration of the national guidelines of the German Society of General Practice and Family Medicine (Deutsche Gesellschaft für Allgemeinmedizin und Familienmedizin, DEGAM) and further literature with focus on the most frequent

infectious diseases leading to prescription of antibiotics in our cohort: tonsillitis, acute bronchitis, upper respiratory tract infections (URTI), urinary tract infections (UTI), and otitis media (Table 1).

Acute bronchitis is typically characterized by cough, sore throat, and fever. Because it is usually caused by a viral infection, antibiotic treatment is not recommended.³¹ Nevertheless, antibiotic treatment may be indicated when patients have certain risk constellations.

Tonsillitis is a frequent infection in patients between five and 15 years and is in this age group often caused by group-A streptococci (GAS).³² Patients report sore throat, headache, and fever, accompanied by gastro-intestinal symptoms in children. The Centor- and McIsaac Score are used to estimate the probability of an infection with GAS based on the occurrence of fever, absence of cough, cervical lymph node swelling, enlarged or occupied tonsils, and age under 15 years. High scores (≥ 3) are used as an indicator for antibiotic treatment. Substance of choice is penicillin V or erythromycin in case of allergy.

Viruses cause the vast majority of acute URTI, often without fever, but with headache, cough, and sore throat, often accompanied by lassitude. In most cases, these infections heal without later complications.^{31,33}

Community-acquired uncomplicated UTI are most commonly caused by *E. coli*. First-line antibiotic therapies are currently fosfomycin, nitrofurantoin, nitroxolin, pivmecillinam, or trimethoprim.

With regard to otitis, it is recommended to abstain from immediate prescription of antibiotics for patients with otitis without risk factors and toddlers over the age of 6 months. In case of aggravation or lack of amelioration within 48 h, the patient should be reevaluated. When an antibiotic therapy is required, amoxicillin or a second-generation cephalosporin are recommended.³⁵

3 | RESULTS

Of all patients in both reception facilities, 45% were female, with significantly more female patients in Friedland than in Celle ($p \leq 0.001$).

TABLE 1 Most frequent diagnoses with antibiotic prescription and recommended treatment according to the DEGAM guidelines

Diagnosis	Antibiotic prescription 1st line	Antibiotic prescription 2nd line
Tonsillitis	<ul style="list-style-type: none"> • Penicillin V if Centor- or McIsaac Score ≥ 3 or delayed prescription (Exacerbation of symptoms/no improvement after 3–5 days) • Score values up to 2: no antibiotics, symptomatic therapy only 	Erythromycin if Centor- or McIsaac Score ≥ 3 and Penicillin allergy
Acute bronchitis	No antibiotics recommended	No antibiotics recommended
URTI	No antibiotics recommended	No antibiotics recommended
UTI	Fosfomycin, nitrofurantoin, nitroxolin, pivmecillinam, trimethoprim	in case of recurrence or new infection <14 days -> change of first-line antibiotic
Otitis media	<ul style="list-style-type: none"> • No antibiotics recommended for patients over the age of 6 months and without fever • Or back up/delayed prescription of amoxicillin 	<ul style="list-style-type: none"> • Second-generation cephalosporins • Makrolid in case of allergies to penicillins/cephalosporins (e.g., erythromycin)

Abbreviations: DEGAM, German Society of General Practice and Family Medicine (Deutsche Gesellschaft für Allgemeinmedizin und Familienmedizin); URTI, upper respiratory tract infections; UTI, urinary tract infections.

On average, patients in Celle were younger (mean: 21.89 years) than those in Friedland (mean: 25.79 years, $p \leq 0.001$), although there were more children below the age of 10 years in Friedland (Table 2). All patients in Celle were asylum seekers, whereas 43% of patients in Friedland were resettlement refugees. Most common countries of origin of patients in both camps were Syria, Iraq, and Afghanistan. On average, patients consulted the doctor 1.9 times during their stay. Of all patients, 50% presented with diagnoses within the ICD-10

category R00-R99 comprising symptoms, signs, abnormal clinical and laboratory findings, and ill-defined conditions for which no diagnosis classifiable elsewhere is recorded. In this category, symptoms such as cough, sore throat, fever and headache, and abdominal pain are usually coded, which are very common in general practice. Other common reasons for consultation in our cohorts were diseases of the respiratory system (35%) and diseases of the musculoskeletal system and connective tissue (15%). Most frequently prescribed drugs were

TABLE 2 Sociodemographic and medical characteristics of the patients in the two reception facilities

	Friedland (n = 2238)	Celle (n = 1017)	Total (N = 3255)	
Sociodemographic characteristics				
Age (years)	Mean = 25.79 (0–90, SD = 18.65)	Mean = 21.89 (0–75, SD = 14.38)	Mean = 24.56 (0–90, SD = 17.49)	
	n (%)	n (%)	n (%)	
Age groups (years)	0–9	610 (27.3)	259 (25.5)	869 (26.7)
	10–19	280 (12.5)	172 (17.2)	452 (13.9)
	20–29	395 (17.7)	304 (29.8)	699 (21.5)
	30–39	450 (20.1)	150 (14.8)	600 (18.4)
	40–49	233 (10.4)	86 (8.5)	319 (9.8)
	50–59	144 (6.5)	34 (3.3)	178 (5.5)
	60–69	92 (4.1)	6 (0.6)	98 (3.0)
	70+	29 (1.3)	1 (0.1)	30 (0.9)
Missing	5 (0.2)	5 (0.5)	10 (0.3)	
Sex (female)	Total female patients	1103 (49.9)	371 (35.5)	1474 (45.3)
	≤19 years ^a	405 (45.5)	183 (42.5)	588 (44.5)
	≥20 years ^a	696 (53.0)	186 (32.0)	882 (45.9)
Residence permit status	Asylum seekers	1275 (57.4)	1017 (100.0)	2292 (70.9)
	Resettlement refugees	948 (42.6)	0 (0)	948 (29.1)
Countries of origin	Syria	906 (40.8)	453 (44.5)	1358 (41.8)
	Iraq	224 (10.1)	191 (18.8)	415 (12.7)
	Afghanistan	130 (5.9)	151 (14.9)	281 (8.6)
	other	978 (43.2)	222 (21.8)	1201 (36.9)
Medical characteristics				
	n	n	n	
Total consultations	4094	2282	6376	
Consultations per patient	1.8	2.2	1.9	
	n (%)	n (%)	n (%)	
Diagnoses ^{b,c} (ICD-10)	R00-R99	949 (42.4)	669 (65.8)	1618 (49.7)
	J00-J99	680 (30.4)	445 (43.7)	1125 (34.6)
	M00-M99	323 (14.4)	153 (15.0)	508 (14.6)
Prescribed drugs ^c (ATC, level 1)	Respiratory system (R)	712 (31.8)	522 (51.3)	1234 (37.9)
	Nervous system (N)	530 (23.7)	258 (25.4)	788 (24.2)
	Anti-infectives for systemic use (J)	491 (21.9)	187 (18.4)	678 (19.7)

Abbreviation: SD, standard deviation.

^aPercentage of all female patients. Due to missing values for age, four persons are missing here.

^bDiagnoses are described based on ICD-10 coding: R00-R99: symptoms, signs, abnormal clinical and laboratory findings, and ill-defined conditions for which no diagnosis classifiable elsewhere is available; J00-J99: diseases of the respiratory system; M00-M99: diseases of the musculoskeletal system and connective tissue.

^cMultiple answers are possible.

those of the ATC groups R = respiratory system (38%), N = nervous system (24%), and J = anti-infectives for systemic use (20%).

3.1 | Prescription of antibiotics

In total, 11% of all consultations resulted in antibiotic prescription. 19% ($N = 624$) of all patients were prescribed at least one antibiotic medication for systemic use. Five hundred and fifty-one patients of these (89%) received only one antibiotic prescription, and 54 patients (9%) received two antibiotic prescriptions. More frequent prescriptions were rare (see Table 3). Patients with at least one antibiotic prescription were significantly younger than those without (mean: 22.8 years, SD: 18.23 vs. 24.8 years, SD: 17.22, $p = 0.003$). Antibiotic

prescriptions were most prevalent in children under the age of 10 years ($N = 209$, 24%) and patients over the age of 70 years ($N = 7$, 24%), (Figure 1). However, the group of people over 70 is very small ($n = 22$). As shown in Table 3, the most commonly prescribed antibiotics were penicillins, macrolides, and cephalosporins. Tonsillitis, acute bronchitis, URTI, UTI, and otitis were the most common diagnoses leading to antibiotic prescriptions. Regarding the total number of antibiotic prescriptions, no differences regarding sex were observed, but cephalosporins were prescribed significantly less to female (31%) than to male patients (69%, $p = 0.048$). Considering only adult patients, women were treated more often (51% out of 45% female patients) with antibiotics than men (49% out of 55% male patients, $p = 0.041$).

Among antibiotic-related diagnoses, female patients were significantly less likely to receive a diagnosis of URTI (30%, $p \leq 0.001$) and

TABLE 3 Characteristics of antibiotic prescriptions

	n (%)	Female sex ^a	
		n (%)	p-Value
Patients with antibiotic prescriptions			
Total patients with antibiotics	621 (19.1) ^b	291 (46.8)	0.345
≤19 years	282 (45.5) ^c	118 (41.8)	0.316
≥20 years	338 (54.5) ^c	172 (50.9)	0.041
Number of antibiotic prescriptions per patient ($n = 621$)			
1 prescription	551 (88.7)	Not calculated due to too small subgroups	
2 prescriptions	54 (8.7)		
3 prescriptions	12 (1.9)		
4 prescriptions	2 (0.3)		
5 prescriptions	1 (0.2)		
6 prescriptions	1 (0.2)		
Consultations with antibiotic prescriptions			
Total consultations with antibiotic prescriptions	691 (10.8) ^b	317 (45.9)	0.799
Most common substance groups (ATC, level 3/4; $n = 691$)			
Penicillins	449 (65.0) ^d	199 (44.4)	0.659
Macrolides	82 (11.9) ^d	41 (50.0)	0.402
Cephalosporins	48 (7.0) ^d	15 (31.3)	0.048
Other	112 (16.2)	62 (53.4)	0.073
Most frequent diagnoses leading to antibiotic prescription ($n = 691$)			
Tonsillitis	179 (25.9) ^d	86 (48.0)	0.449
Acute bronchitis	143 (20.7) ^d	69 (48.3)	0.533
URTI	95 (13.6) ^d	29 (30.3)	0.001
UTI	70 (10.1) ^d	52 (74.3)	0.001
Otitis media	60 (8.7) ^d	28 (46.7)	0.898
Other	109 (15.8) ^d	45 (37.2)	0.035
No diagnosis	35 (5.1) ^d	18 (51.4)	0.499

Note: The whole column p -value represents the corresponding significance values. For a quicker overview, we have marked the values <0.05 in bold. Abbreviations: URTI, upper respiratory tract infections; UTI, urinary tract infections.

^aPercentages must be interpreted in the context of the proportion of 45.2% women in the total patient cohort. p -Values show Chi²-tests regarding the distribution between women and men.

^bPercentage of total sample.

^cPercentage of patients with antibiotics. Due to missing in age one person is missing here.

^dPercentage of consultations with antibiotic prescriptions.




FIGURE 1 Age of patients with at least one antibiotic prescription

significantly more likely to receive a diagnosis of UTI (74%, $p \leq 0.001$; Table 3).

3.2 | Diagnosis-specific antibiotic prescription

As shown in Table 4, 89% of patients diagnosed with tonsillitis received antibiotic treatment. They were treated with amoxicillin in 67% of cases followed by phenoxymethylpenicillin (12%). Phenoxymethylpenicillin complies with DEGAM guidelines if Centor- or McIsaac Score are >3 or if symptoms exacerbate or do not improve after 3–5 days. Relevant McIsaac Score parameters were age less than 15 years in 44%, fever in 11%, and cervical lymph node swelling in 3% of cases of tonsillitis with antibiotic prescription.

Acute bronchitis was treated with antibiotics in 74% of cases, contrary to the DEGAM guidelines in Table 1. Most frequently prescribed antibiotics in this context were amoxicillin (45%) and doxycyclin (13%).

In the third largest group of antibiotic-relevant diagnoses, URTI ($n = 95$), 8% of all patients received antibiotic treatment, of whom half were treated with amoxicillin (4%). The very low prescribing rate for URTI is consistent with the guidelines.

Patients with UTI were treated with antibiotics in 73% of cases, mostly with sulfamethoxazole/trimethoprim and ciprofloxacin, although neither is recommended by the DEGAM guideline.

Forty-two percent of patients with otitis media were treated with antibiotics and mostly with amoxicillin (37%), which is in accordance with the guidelines in case of age under 6 months or high fever and if a re-presentation after 24 h is not possible. Fever was documented in 15% of antibiotic-treated patients with otitis media.

4 | DISCUSSION

To our knowledge, this is the first study that collected systematic data on antibiotic prescriptions in a representative group of asylum seekers in Germany during the current exodus. Given that a large proportion of migrants in our cohort were younger than 30 years, the majority of

refugees were male, and home countries frequently were Syria, Iraq, and Afghanistan, the demographic characteristics match the group of all asylum seekers in Germany in the immigration statistics since 2015.^{2,36}

In most cases, doctors coded patients' symptoms such as cough, sore throat, fever, and headache using the ICD-10 category R00-99. The underlying respiratory ailments are in accordance with primary healthcare demands in the respective German population, where respiratory diseases are a highly prevalent complaint leading to medical attendance.³⁷

Overall, 19% patients in our cohorts received systemic antibiotics during the observation period. Patients with antibiotic prescription were significantly younger than those without. Nearly a quarter of the patients below the age of 10 years received antibiotic treatment, and about 45% of all antibiotics were prescribed to patients up to age of 19 years. This result complies with previous research on age distribution of antibiotic prescriptions.⁷⁻⁹ In general, women in Germany are reported to receive more antibiotics than men.^{9,38} This is seen as a result of an overall higher prevalence of UTI in women than in men, and of sex-specific differences in utilization of health care services.^{39,40} The latter could be confirmed for the Celle cohort, as well for children and adolescents²⁹ as for adults.³⁰ For the Friedland cohort, sex-dependent differences in medical attendance could not be investigated as there were no sociodemographic characteristics available for all Friedland residents.

In our whole patient cohort, we found no sex dependency in overall antibiotic prescribing: The proportion of female individuals was 47% in the group of patients with antibiotic prescription and corresponded to that of female individuals (45%) in the whole cohort. As mentioned above, about 45% of antibiotics ($n = 282$ out of 621) in our study were prescribed to patients up to the age of 19 years, and sex dependencies are presumably less likely in diagnoses most frequently treated with antibiotics in this age group. On the other hand, only 10% ($n = 70$ out of 691) of all antibiotic prescriptions were due to UTI, a diagnosis that women are significantly more often diagnosed with and treated for. These two subgroups' respective proportions of all antibiotic prescriptions might reduce sex-dependent influences in the whole patient cohort. However, considering only adult patients (age ≥ 20 years), our data do confirm the formerly described higher frequency of antibiotic prescription to women.

In our setting, most antibiotics were prescribed for respiratory diseases, which is in line with previous findings in German^{6,21,41} and international primary care.^{9,42} Frequent diagnoses in case of antibiotic prescription were tonsillitis, acute bronchitis, URTI, UTI, and otitis. Following the DEGAM guidelines, there is no need for antibiotic prescription in patients with acute bronchitis or URTI because these infections are mostly caused by viruses.^{12-14,31,33} In our study, 237 not indicated antibiotic prescriptions for acute bronchitis and URTI were identified (34% of all antibiotic prescriptions). One reason for the presumably not indicated prescription of antibiotics may be the doctors' fear of a higher number of complications.⁴³

In our setting, 89% of the patients with tonsillitis received an antibiotic. Only a minority of these patients met criteria for bacterial

TABLE 4 Diagnoses most frequently treated with antibiotics and respective prescriptions

Diagnoses (n)	Antibiotic prescription n (%)	Most frequently prescribed antibiotics		Consistent with recommendations
		Name	n (%)	
Tonsillitis (201)	179 (89.1)	Amoxicillin	134 (66.7)	No
		Phenoxymethylpenicillin	25 (12.4)	Possibly yes, first choice
		Others	20 (10.0)	-
Acute bronchitis (191)	142 (74.4)	Amoxicillin	85 (44.5)	No
		Doxycyclin	25 (13.1)	No
		Roxithromycin	15 (7.9)	No
		Others	17 (8.9)	-
URTI (1271)	95 (7.5)	Amoxicillin	51 (4.0)	No
		Clarithromycin	16 (1.3)	No
		Cefuroxime	11 (0.9)	No
		Others	17 (1.3)	-
UTI (96)	70 (72.9%)	Sulfamethoxazole + Trimethoprim	25 (26.0)	No
		Ciprofloxacin	22 (22.9)	No
		Others	23 (24.0)	-
Otitis media (143)	60 (42.0)	Amoxicillin	53 (37.1)	Yes, second choice, or delayed prescription
		Others	7 (4.9)	-

Abbreviations: URTI, upper respiratory tract infections; UTI, urinary tract infections.

infections such as cervical lymph node swelling or fever, which indicate GAS infection and should lead to antibiotic treatment according to current guidelines.³³ However, it is difficult to capture the parameters of the Centor- or McIsaac Score from our data, because “no cough” was not documented and probably other symptoms were often not coded separately. Even if we assume that antibiotic treatment was indicated, amoxicillin was prescribed in 67% of cases, which is neither first nor second line antibiotic.³³

Almost half of the patients with UTI received ciprofloxacin or cotrimoxazol (sulfamethoxazole and trimethoprim). These drug therapies are not recommended in the guidelines anymore,³⁴ but are generally often prescribed in primary care settings, possibly because they were still among the second-line antibiotics in 2009.⁴⁴ In none of the recorded UTI cases, fever was documented. Fever could be a sign for a pyelonephritis, in which case ciprofloxacin would be the preferred treatment option.³⁴ Changes in the resistance situation or updates to the benefit-risk profiles mean that antibiotic guidelines are constantly updated. Neugebauer et al show that 23% of physicians have no access to a database of national guidelines⁴⁵ and instead use their usual medications.

In our observation, patients with otitis media received amoxicillin in 88% of antibiotic prescriptions. Following DEGAM guidelines, amoxicillin is the agent of choice for treating otitis media when antibiotic therapy is indicated, but the guideline emphasizes the principle of reluctant prescription when no risk factors are present.^{35,46} However, beyond refugee healthcare, the concept of reluctant or delayed prescription of antibiotics is often not followed by general practitioners. Petruscke et al. suggest that main reasons for abstaining from delayed prescription were impending weekend or a hard-to-estimate disease

course (44%), as well as the patient's demand for an antibiotic prescription (30%), unfamiliar patients (15%), or language barriers and mental limitations of patients (13%).⁴⁷

In refugee healthcare, studies revealed that physicians rate language barriers as the strongest obstacle for treating a patient in an optimal fashion,⁴⁸⁻⁵¹ followed by cultural differences, psychic trauma, and lacking cooperation of the patients.⁴⁸ Language barriers are also associated with higher healthcare costs⁵² and lower treatment quality²⁵⁻²⁸ and pose the largest impediment for refugees when they seek medical advice.^{50,53,54} In our cohort, the frequent and presumably not indicated prescription of amoxicillin may arise because antibiotics are freely available in some countries and may be a “go-to medicine” for many migrants seeking medical help. It is conceivable that patients might ask for it directly and doctors might give in because of the perceived pressure and the lack of communication possibilities. In general, in our setting of healthcare provision in initial reception centers, cultural and language barriers as well as changing medical staff with limited time resources who are caring for a permanently fluctuating group of refugees may have contributed to the high rate of antibiotic prescriptions.

Our study has several limitations. First, the retrospective design prohibited systematic screening for antibiotic-relevant symptoms in all patients. Diagnoses were mainly based on clinical judgment of the experienced general practitioners. It is possible that the doctors' documentation was incomplete and some antibiotic-related symptoms were not recorded. In addition, a bias among doctors cannot be excluded. However, the representativity and size of the patient sample is a strength of the study.

5 | SUMMARY

Taken together, these data show a rate between 8% and 75% of inappropriate prescriptions of antibiotics in a large current cohort of refugees newly arriving in Germany. This corresponds to inappropriate prescriptions in general population in several high-middle income countries.⁵⁵⁻⁵⁸ Especially in the context of common viral infections such as acute bronchitis, the observed high rate in the prescription of antibiotics appears to be associated with the risks of unnecessary costs, side effects, and the emergence of antimicrobial resistance. In cases of tonsillitis or UTI, where antibiotics are generally appropriate, substances were frequently prescribed that were not or no longer in accordance with the guidelines. Unfortunately, we cannot assess the extent to which infectious diseases with resistant microbial strains occur among refugees and migrants based on our data. This aspect calls for further investigation.

In our setting of healthcare provision in initial reception centers, cultural and language barriers as well as changing medical staff with limited time resources may have contributed to the high rate of inappropriate prescription of antibiotics. Healthcare providers caring for refugees should be aware of this problem and be particularly trained in antibiotic stewardship. We hope to contribute to developing regimens for stringent antibiotic stewardship in the particularly vulnerable population of migrants and refugees in the current situation.

ACKNOWLEDGMENTS

The authors acknowledge the efforts and help of Malteser Friedland, the social services office at Friedland transit camp, and the Landesaufnahmebehörde Niedersachsen (Migration Authority of Lower Saxony). The DICTUM-Friedland study was funded by the Robert Bosch Foundation, the state of Lower Saxony (the subsidy guidelines for the healthcare region), and the European Social Fund (the subsidy guidelines for social innovation). The funding body played no role in the design of the study and collection, analysis, and interpretation of data. CH received funding from the Young Academy through the Hannover Medical School, Germany. AJ received a maternity leave grant from the German Centre for Infection Research at Hannover Medical School - Young Academy. Open Access funding enabled and organized by Projekt DEAL.

CONFLICT OF INTEREST

The authors declare they have no conflict of interest.

ETHICS STATEMENT

The Research Ethics Board of the Göttingen University Medical Center approved the study (Ethics Approval No. 18/5/16) as part of the DICTUM-Friedland study.⁵⁹ The study site in Celle was separately approved by the Research Ethics Board of Hannover Medical School (Ethics Approval No. 3217-2016).

ORCID

Evelyn Kleinert  <https://orcid.org/0000-0002-8951-9874>

REFERENCES

1. The UN Refugee Agency. *Global Trends: Forced Displacement in 2018*; Geneva; 2019. <https://www.unhcr.org/figures-at-a-glance.html>.
2. Bundesamt für Migration und Flüchtlinge 2018. *Das Bundesamt in Zahlen*. Nürnberg; 2018.
3. Mews C, Pruskil S, Kloppe T, Wilsdorf S, Scherer M. Einsatz von Videodolmetschen in der ambulanten Versorgung in Hamburg – eine Bedarfsanalyse. *Z Allg Med*. 2017;93(11):461-465.
4. Alberer M, Wendeborn M, Löscher T, Seilmaier M. Erkrankungen bei Flüchtlingen und Asylbewerbern: Daten von drei verschiedenen medizinischen Einrichtungen im Raum München aus den Jahren 2014 und 2015. *Dtsch Med Wochenschr*. 2016;141:e8-e15.
5. Kleinert E, Mueller F, Furaijat G, et al. Does refugee status matter? Medical needs of newly arrived asylum seekers and resettlement refugees - a retrospective observational study of diagnoses in a primary care setting. *Confl Health*. 2019;13(39):1-9. <https://conflictandhealth.biomedcentral.com/articles/10.1186/s13031-019-0223-z>.
6. Schwabe U, Paffrath D, Ludwig W-D, Klauber J, eds. *Arzneiverordnungs-Report 2017*. Berlin: Springer; 2017.
7. Gradl G, Teichert M, Kieble M, Werning J, Schulz M. Comparing outpatient oral antibiotic use in Germany and The Netherlands from 2012 to 2016. *Pharmacoepidemiol Drug Saf*. 2018;27(12):1344-1355. <https://doi.org/10.1002/pds.4643>.
8. Glaeske G, Hoffmann F, Koller D, Tholen K, Windt R. *Faktencheck Gesundheit: Antibiotika-Verordnungen bei Kindern*. Gütersloh: Bertelsmann-Stiftung; 2012. faktencheck-gesundheit.de.
9. Brauer R, Ruigómez A, Downey G, et al. Prevalence of antibiotic use: a comparison across various European health care data sources. *Pharmacoepidemiol Drug Saf*. 2016;25(suppl 1):11-20. <https://doi.org/10.1002/pds.3831>.
10. Smith DRM, Dolk FCK, Smieszek T, Robotham JV, Pouwels KB. Understanding the gender gap in antibiotic prescribing: a cross-sectional analysis of English primary care. *BMJ Open*. 2018;8(2):e020203. <https://doi.org/10.1136/bmjopen-2017-020203>.
11. Schröder W, Sommer H, Gladstone BP, et al. Gender differences in antibiotic prescribing in the community: a systematic review and meta-analysis. *J Antimicrob Chemother*. 2016;71(7):1800-1806. <https://doi.org/10.1093/jac/dkw054>.
12. Zweigner J, Meyer E, Gastmeier P, Schwab F. Rate of antibiotic prescriptions in German outpatient care – are the guidelines followed or are they still exceeded? *GMS Hyg Infect Control*. 2018;1-8.
13. Little P, Stuart B, Moore M, et al. Amoxicillin for acute lower-respiratory-tract infection in primary care when pneumonia is not suspected: a 12-country, randomised, placebo-controlled trial. *Lancet Infect Dis*. 2013;13(2):123-129. [https://doi.org/10.1016/S1473-3099\(12\)70300-6](https://doi.org/10.1016/S1473-3099(12)70300-6).
14. Butler CC, Hood K, Kelly MJ, et al. Treatment of acute cough/lower respiratory tract infection by antibiotic class and associated outcomes: a 13 European country observational study in primary care. *J Antimicrob Chemother*. 2010;65(11):2472-2478. <https://doi.org/10.1093/jac/dkq336>.
15. Kohut MR, Keller SC, Linder JA, et al. The inconvincible patient: how clinicians perceive demand for antibiotics in the outpatient setting. *Fam Pract*. 2020;37(2):276-282. <https://doi.org/10.1093/fampra/cmz066>.
16. Richter-Kuhlmann E. Global und Interdisziplinär: Kampf gegen Antibiotikaresistenzen. *Dtsch Arztebl*. 2017;114(5):A210-A211.
17. Dobson EL, Klepser ME, Pogue JM, et al. Outpatient antibiotic stewardship: interventions and opportunities. *J Am Pharm Assoc*. 2003;57(4):464-473. <https://doi.org/10.1016/j.japh.2017.03.014>.
18. CDC. *The Core Elements of Hospital Antibiotic Stewardship Programs*. Atlanta: US Department of Health and Human Services; 2019.
19. Drekonja DM, Filice GA, Greer N, et al. Antimicrobial stewardship in outpatient settings: a systematic review. *Infect Control Hosp Epidemiol*. 2015;36(2):142-152. <https://doi.org/10.1017/ice.2014.41>.

20. Salm F, Ernsting C, Kuhlmeier A, Kanzler M, Gastmeier P, Gellert P. Antibiotic use, knowledge and health literacy among the general population in Berlin, Germany and its surrounding rural areas. *PLoS ONE*. 2018;13(2):e0193336. <https://doi.org/10.1371/journal.pone.0193336>.
21. DAK-Gesundheit. *Antibiotika-Report 2014: Eine Wunderwaffe wird stumpf: Folger der Über- und Fehlversorgung*. Hamburg; 2014. <https://www.dak.de/dakonline/live/dak/download/antibiotika-report-2014-1486100.pdf>. Accessed April 30, 2019.
22. Schneider S, Salm F, Schröder C, Ludwig N, Hanke R, Gastmeier P. Antibiotikaeinnahme und Resistenzentwicklung - Wissen, Erfahrungen und Einnahmeverhalten innerhalb der deutschen Allgemeinbevölkerung. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. 2016;59(9):1162-1170. <https://doi.org/10.1007/s00103-016-2417-5>.
23. McNulty CAM, Boyle P, Nichols T, Clappison P, Davey P. The public's attitudes to and compliance with antibiotics. *J Antimicrob Chemother*. 2007;60(suppl 1):i63-i68. <https://doi.org/10.1093/jac/dkm161>.
24. Heß B. SoKo Analysis for First Half of 2018 2019; <http://www.bamf.de/DE/DasBAMF/Forschung/Ergebnisse/Kurzanalysen/kurzanalysen-node.html>. Accessed June 14, 2019.
25. Ngo-Metzger Q, Sorkin DH, Phillips RS, et al. Providing high-quality care for limited English proficient patients: the importance of language concordance and interpreter use. *J Gen Intern Med*. 2007;22(suppl 2):324-330. <https://doi.org/10.1007/s11606-007-0340-z>.
26. Divi C, Koss RG, Schmaltz SP, Loeb JM. Language proficiency and adverse events in US hospitals: a pilot study. *Int J Qual Health Care*. 2007;19(2):60-67. <https://doi.org/10.1093/intqhc/mzl069>.
27. Wilson E, Chen AHM, Grumbach K, Wang F, Fernandez A. Effects of limited English proficiency and physician language on health care comprehension. *J Gen Intern Med*. 2005;20(9):800-806. <https://doi.org/10.1111/j.1525-1497.2005.0174.x>.
28. Weinick RM, Krauss NA. Racial/ethnic differences in children's access to care. *Am J Pub Health*. 2000;90(11):1771-1774.
29. Happle C, Dopfer C, Ernst D, et al. Pediatric healthcare utilization in a large cohort of refugee children entering Western Europe during the migrant crisis. *IJERPH*. 2019;16(22):1-11. <https://doi.org/10.3390/ijerph16224415>.
30. Wetzke M, Happle C, Vakilzadeh A, et al. Healthcare utilization in a large cohort of asylum seekers entering Western Europe in 2015. *IJERPH*. 2018;15(10):2163. <https://doi.org/10.3390/ijerph15102163>.
31. Deutsche Gesellschaft für Allgemeinmedizin und Familienmedizin. *Leitlinie Husten*; 2014.
32. Anderson J, Paterek E. *Tonsillitis*. Treasure Island, FL: StatPearls; 2020.
33. Deutsche Gesellschaft für Allgemeinmedizin und Familienmedizin. *Halsschmerzen*. Stand Oktober 2009. Düsseldorf; 2009. DEGAM-Leitlinie.
34. Deutsche Gesellschaft für Allgemeinmedizin und Familienmedizin. *Brennen beim Wasserlassen: S3-Leitlinie und Anwenderversion der S3-Leitlinie Harnwegsinfektionen*; 2018.
35. DEGAM. *Ohrenschmerzen*; 2014.
36. Bundesamt für Migration und Flüchtlinge. *Das Bundesamt in Zahlen 2015: Asyl, Migration und Integration*; 2016.
37. Steppuhn H, Buda S, Wienecke A, et al. Zeitliche trends in der Inzidenz und Sterblichkeit respiratorischer Krankheiten von hoher Public-Health-Relevanz in Deutschland. *Journal of Health Monitoring* 2017. Berlin: Robert Koch-Institut; 2017. https://www.rki.de/DE/Content/Gesundheitsmonitoring/JoHM/JoHM_node.html.
38. Wissenschaftliches Institut der AOK. *Arzneiverbrauch nach Altersgruppen 2017: Verordnungen, definierte Tagesdosen, Umsätze aufgliedert nach Alters- und Geschlechtsstruktur der gesetzlichen Krankenkassen*; 2018.
39. Rattay P, Butschalowsky H, Rommel A, et al. Inanspruchnahme der ambulanten und stationären medizinischen Versorgung in Deutschland: Ergebnisse der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1). *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. 2013;56:832-844. <https://doi.org/10.1007/s00103-013-1665-x>.
40. Laubach W, Brähler E. Körperliche Symptome und Inanspruchnahme ärztlicher Versorgung. *Dtsch Med Wschr*. 2001;126:T1-T7.
41. Holstiege J, Garbe E. Systemic antibiotic use among children and adolescents in Germany: a population-based study. *Eur J Pediatr*. 2013;172(6):787-795. <https://doi.org/10.1007/s00431-013-1958-y>.
42. Dolk FCK, Pouwels KB, Smith DRM, Robotham JV, Smieszek T. Antibiotics in primary care in England: which antibiotics are prescribed and for which conditions? *J Antimicrob Chemother*. 2018;73(2):ii2-ii10. <https://doi.org/10.1093/jac/dkx504>.
43. Gulliford MC, Moore MV, Little P, et al. Safety of reduced antibiotic prescribing for self limiting respiratory tract infections in primary care: cohort study using electronic health records. *BMJ*. 2016;354:i3410. <https://doi.org/10.1136/bmj.i3410>.
44. Dicheva S. *Harnwegsinfekte bei Frauen. BARMER GEK Arzneimittelreport 2015*. Berlin: BARMER GEK; 2015.
45. Neugebauer M, Ebert M, Vogelmann R. Lack of information and provision of information at the workplace as potential reasons for inappropriate antibiotic therapy in Germany. *Z Evid Fortbild Qual Gesundheitswes*. 2019;144-145:35-41. <https://doi.org/10.1016/j.zefq.2019.06.002>.
46. Simon A, Tenenbaum T, Huppertz HI, et al. Diagnose und Therapie von Atemwegsinfektionen (ohne ambulant erworbene Pneumonie) bei ambulant behandelten Kindern ohne schwerwiegende Grunderkrankung. *Monatsschr Kinderheilkd*. 2017;165(8):711-724. <https://doi.org/10.1007/s00112-017-0257-5>.
47. Petruscke I, Salm F, Schneider S, Gastmeier P, Genischen J. *Antibiotikaspezifisches Verschreibungsverhalten von Hausärzten - Ergebnisse einer Querschnittsbefragung im Rahmen des RAI-Projekts*. Frankfurt am Main: 50. Kongress der Deutschen Gesellschaft für Allgemeinmedizin; 2016. http://www.rai-projekt.de/fileadmin/rai/downloads/DEGAM_2016_Petruscke.pdf.
48. Fölsch UR, Hasenfuß G, Spies H-F, Wesiack W, Faulbaum F. Flucht und Migration: Eine Herausforderung für die Medizin in Deutschland. *Internist (Berl)*. 2016;57(8):822-830. <https://doi.org/10.1007/s00108-016-0103-1>.
49. Nesterko Y, Glaesmer H. Verständigung mit Patienten mit Migrationshintergrund aus der Sicht von Hausärzten: the quality of communication between family practitioners and their migrant patients. *Z Allg Med*. 2015;91(12):506-511.
50. Bermejo I, Hölzel LP, Kriston L, Härter M. Subjektiv erlebte Barrieren von Personen mit Migrationshintergrund bei der Inanspruchnahme von Gesundheitsmaßnahmen. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz*. 2012;55(8):944-953. <https://doi.org/10.1007/s00103-012-1511-6>.
51. Karger A, Lindtner-Rudolph H, Mroczynski R, Ziem A, Joksimovic L. Wie fremd ist mir der Patient? Erfahrungen, Einstellungen und Erwartungen von Ärztinnen und Ärzten bei der Versorgung von Patientinnen und Patienten mit Migrationshintergrund. *Z Psychosom Med Psychother*. 2017;63:280-296.
52. Bischoff A, Denhaerynck K. What do language barriers cost? An exploratory study among asylum seekers in Switzerland. *BMC Health Serv Res*. 2010;10:248. <https://doi.org/10.1186/1472-6963-10-248>.
53. Schröder H, Zok K, Faulbaum F. Gesundheit von Geflüchteten in Deutschland - Ergebnisse einer Befragung von Schutzsuchenden aus Syrien, Irak und Afghanistan. *WldO Monitor*. 2018;15(1):1-20.
54. van Loenen T, van den Muijsenbergh M, Hofmeester M, et al. Primary care for refugees and newly arrived migrants in Europe: a qualitative study on health needs, barriers and wishes. *Eur J Public Health*. 2018; 28(1):82-87. <https://doi.org/10.1093/eurpub/ckx210>.

55. White AT, Clark CM, Sellick JA, Mergenhausen KA. Antibiotic stewardship targets in the outpatient setting. *Am J Infect Control*. 2019;47(8): 858-863. <https://doi.org/10.1016/j.ajic.2019.01.027>.
56. Dekker ARJ, Verheij TJM, van der Velden AW. Inappropriate antibiotic prescription for respiratory tract indications: most prominent in adult patients. *Fam Pract*. 2015;32(4):401-407. <https://doi.org/10.1093/fampra/cmz019>.
57. Fleming-Dutra KE, Hersh AL, Shapiro DJ, et al. Prevalence of inappropriate antibiotic prescriptions among US ambulatory care visits, 2010-2011. *Jama*. 2016;315(17):1864-1873. <https://doi.org/10.1001/jama.2016.4151>.
58. Butt AA, Navasero CS, Thomas B, et al. Antibiotic prescription patterns for upper respiratory tract infections in the outpatient Qatari population in the private sector. *Int J Infect Dis*. 2017;55:20-23. <https://doi.org/10.1016/j.ijid.2016.12.004>.
59. Furajjat G, Kleinert E, Simmenroth A, Müller F. Implementing a digital communication assistance tool to collect the medical history of refugee

patients: DICTUM Friedland - an action-oriented mixed methods study protocol. *BMC Health Serv Res*. 2019;19(1):103. <https://doi.org/10.1186/s12913-019-3928-1>.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Kleinert E, Hillermann N, Jablonka A, Happel C, Müller F, Simmenroth A. Prescription of antibiotics in the medical care of newly arrived refugees and migrants. *Pharmacoepidemiol Drug Saf*. 2021;30:1074–1083. <https://doi.org/10.1002/pds.5254>