

## Recombinant Plasmids Capable of Replication in *B. subtilis* and *E. coli*

J. Kreft, K. Bernhard, and W. Goebel

Institut für Genetik und Mikrobiologie, Universität Würzburg,  
Röntgenring 11, D-8700 Würzburg, Federal Republic of Germany

**Summary.** The plasmid *pBC16* (4.25 kbases), originally isolated from *Bacillus cereus*, determines tetracycline resistance and can be transformed into competent cells of *B. subtilis*. A miniplasmid of *pBC16* (*pBC16-1*), 2.7 kb) which has lost an *EcoRI* fragment of *pBC16* retains the replication functions and the tetracycline resistance. This plasmid which carries only one *EcoRI* site has been joined *in vitro* to *pBS1*, a cryptic plasmid previously isolated from *B. subtilis* and shown to carry also a single *EcoRI* site (Bernhard et al., 1978). The recombinant plasmid is unstable and dissociates into the plasmid *pBS161* (8.2 kb) and the smaller plasmid *pBS162* (2.1 kb). Plasmid *pBS161* retains the tetracycline resistance. It possesses a single *EcoRI* site and 6 *HindIII* sites. The largest *HindIII* fragment of *pBS161* carries the tetracycline resistance gene and the replication function. After circularization *in vitro* of this fragment a new plasmid, *pBS161-1* is generated, which can be used as a *HindIII* and *EcoRI* cloning vector in *Bacillus subtilis*.

Hybrid plasmids consisting of the *E. coli* plasmids *pBR322*, *pWL7* or *pAC184* and different *HindIII* fragments of *pBS161* were constructed *in vitro*. Hybrids containing together with the *E. coli* plasmid the largest *HindIII* fragment of *pBS161* can replicate in *E. coli* and *B. subtilis*. In *E. coli* only the replicon of the *E. coli* plasmid part is functioning whereas in *B. subtilis* replication of the hybrid plasmid is under the control of the *Bacillus* replicon. The tetracycline resistance of the *B. subtilis* plasmid is expressed in *E. coli*, but several antibiotic resistances of the *E. coli* plasmids (ampicillin, kanamycin and chloramphenicol) are not expressed in *B. subtilis*. The hybrid plasmids seem to be more unstable in *B. subtilis* than in *E. coli*.

### Introduction

Recently, we have reported on the isolation and characterization of a large number of plasmids from *B.*

*subtilis* and *B. cereus* (Bernhard et al., 1978). Among those a rather small plasmid, *pBC16*, originally isolated from a tetracycline resistant strain of *B. cereus* could be transformed into *B. subtilis*, where it is stably maintained rendering the transformants resistant to tetracycline. The molecular properties of this plasmid (size, copy number, sites for restriction enzymes, genetic marker) made it appear a suitable vector for gene cloning in *B. subtilis*. We have therefore started experiments to insert fragments from other *B. subtilis* and *E. coli* DNA molecules into this plasmid and to analyze the resultant recombinant DNAs.

Other vector plasmids suitable for the application of this new genetic method in *B. subtilis* have been recently reported (S. Ehrlich, 1977). These plasmids derive from *Staphylococcus aureus*.

Like *pBC16* they seem to be suitable vectors for gene cloning in *B. subtilis* (S. Ehrlich, personal communication; P. Lovett, personal communication). We have made the experience, however, that they seem to be considerably more unstable in the new host than *pBC16*. This report describes experiments, which demonstrate that *pBC16* and some of its *in vitro* constructed derivatives can be used for cloning of homologous and heterologous DNA in *Bacillus subtilis*.

### Materials and Methods

**Bacterial Strains.** *E. coli* 5K,  $r^-$ ,  $m^+$  was obtained from S. Glover. JC 1569  $recA^-$ ,  $thy^+$ ,  $arg^-$ ,  $his^-$ ,  $met^-$ ,  $leu^-$ ,  $Sm^R$  harbouring plasmid pAC184 was a gift from Cohen and Chang. *E. coli* DP9,  $polA^-$ ,  $r^-$ ,  $m^-$  was provided by Ph. Kourilsky. *B. subtilis* 168  $trp^-$  (DSM 402) was from the Deutsche Sammlung von Mikroorganismen, *B. subtilis* BR151-CM1  $trp^-$ ,  $lys^-$ ,  $met^-$ ,  $spo^-$  was kindly provided by P. Lovett. The other strains used here, were described before (Bernhard et al., 1978; Kollek et al., 1978).

**Source of Reagents.** All reagents if not otherwise stated were from Merck, Germany. The antibiotics used were a gift from Bayer, Germany. Sodium dodecyl sulfate (SDS), ethidium bromide, polyethylenglycol type 6000, lysozyme, pancreatic RNase and pronase P from *S. griseus* were obtained from Serva, Germany.

**Enzymes.** The restriction enzyme *EcoRI* was provided by H. Mayer (Braunschweig), *HindIII* and T4 DNA ligase was purchased from Bio Labs (USA).

**Growth of Strains, Cell Lysis and Purification of Plasmid DNA** were performed as previously described (Mayer et al., 1977; Bernhard et al., 1978).

**Cleavage of DNA and Electrophoresis in Agarose Gels.** Digestion with *EcoRI* was carried out in 100 mM Tris-HCl pH 7.5, 50 mM NaCl, 5 mM MgCl<sub>2</sub> at 37° C for 30 min, with *HindIII* in 10 mM Tris-HCl pH 7.5, 25 mM NaCl, 10 mM MgCl<sub>2</sub> at 37° C for 60 min. The reactions were stopped by heating to 68° C for 8 min or by addition of 100 mM EDTA. Electrophoresis was performed on 1% agarose slab gels in 36 mM Tris-HCl, 30 mM NaH<sub>2</sub> PO<sub>4</sub>, 10 mM EDTA, pH 7.5 (Meyers et al., 1975)

**Construction of Hybrid Plasmids and Transformation.** The *in vitro* construction of hybrid plasmids (Cohen et al., 1973) using T4 DNA ligase, transformation of *E. coli* and of *B. subtilis* have been described (Goebel and Bonewald, 1975; Cahn and Fox, 1968). Transformants of *E. coli* were first selected on nutrient broth (ENB) plates containing 100 µg/ml ampicillin (in the case of *pBR322* or *pWL7* hybrids) or 100 µg/ml chloramphenicol (for *pAC184* hybrids). Growing colonies were then tested for sensitivity to 5–50 µg/ml tetracycline (*pBR322* and *pAC184* hybrids), to 100 µg/ml kanamycin (for *pWL7* hybrids). In addition *pWL7* hybrids (Km<sup>S</sup>) were tested for sensitivity against 5–50 µg/ml tetracycline.

**Determination of the Intracellular β-Lactamase Activity.** The enzyme was assayed with hydroxylamine/ammonium ferric sulphate (Dale and Smith, 1971). 1 unit corresponds to the amount of enzyme which catalyses the hydrolysis of one µmole ampicillin/min at 30° C in 0.025 M Tris-HCl, pH 7.4.

## Results

### *In vitro* Construction of Recombinant Plasmids from *B. subtilis* Plasmid *pBS1* and *B. cereus* Plasmid *pBC16*


The tetracycline resistance plasmid *pBC16* (4.2 kb) was previously isolated from *B. cereus* and transformed into *B. subtilis* 168 (Bernhard et al., 1978). Restriction endonuclease *EcoRI* cleaves this plasmid into two fragments of 2.7 kb and 1.5 kb. The cryptic plasmid *pBS1* (7.8 kb) of *B. subtilis* carries a single *EcoRI* site. To construct *in vitro* recombinant plasmids carrying both replicons, *pBC16* was either completely or partially digested by *EcoRI* and ligated, using T4 DNA ligase, with *EcoRI* cleaved *pBS1*. The ligated mixture was transformed into *B. subtilis* 168 made competent as described (Cahn and Fox, 1968). Tetracycline resistant (Tc<sup>R</sup>) colonies were obtained with a frequency of about 10<sup>-6</sup>. Covalently-closed circular (CCC) DNAs were isolated from several

of the obtained clones. None of them represented the expected hybrid plasmid consisting of both replicons. Instead, the CCC-DNAs isolated from several Tc<sup>R</sup> clones could be identified as:

*a) pBC16-1:* Some of tetracycline resistant cells harbour one small plasmid of 2.7 kb, which carries a single *EcoRI* site. This plasmid, which can be also obtained by transforming *B. subtilis* with *pBC16*, completely digested by *EcoRI* and then ligated *in vitro*, represents the circularized large *EcoRI* fragment of *pBC16* (Fig. 1). This mini *pBC16* plasmid, named *pBC16-1*, thus carries the tetracycline resistance gene and the replication functions of *pBC16*.


*b) pBC16-2:* Other Tc<sup>R</sup>-transformants harbour two plasmids of 5.8 kb and 4.2 kb. The smaller one is identical to *pBC16*. Transformation of the isolated larger plasmid yields tetracycline resistant colonies which harbour again both plasmids. Cleavage of the mixture of both plasmids by *EcoRI* (Fig. 1) yields only the two fragments which are also obtained after cleavage of *pBC16* DNA alone. Since the size difference between *pBC16* and the larger plasmid, *pBC16-2*, is 1.5 kb and hence identical to the small *EcoRI* fragment of *pBC16*, we conclude that plasmid *pBC16-2* carries a second small *EcoRI* fragment of *pBC16* inserted into *pBC16*. This plasmid is, however, unstable and eliminates at a rather high rate one of the two identical fragments.

*c) pBS161 and pBS162:* A large number of the tetracycline resistant colonies harbour two plasmids of 8.2 kb (*pBS161*) and 2.1 kb (*pBS162*) (Fig. 1). Plasmid *pBS161* alone but not *pBS162* yields also tetracycline resistant colonies when transformed into competent *B. subtilis* cells. This plasmid being larger in size than *pBS1* but smaller than the expected sum of *pBS1* and *pBC16* or one of its two *EcoRI* fragments, carries only one *EcoRI* site (Fig. 1) but 6 *HindIII* sites. Comparison of the *HindIII* patterns of *pBS1* and *pBS161* reveals that fragments B, C, E, F and G are identical with *pBS1* (Fig. 2 and Fig. 4). Fragment D of *pBS1* is lost, whereas fragment A' has obviously incorporated additional DNA from *pBC16* including the gene determining tetracycline resistance. The smaller plasmid *pBS162* being present in the original clones carries also one *EcoRI* and one *HindIII* site. Since it represents a replicating DNA element it must possess one of the replicon functions of the two parent plasmids *pBC16* or *pBS1*. The sum of the sizes of *pBS161* and *pBS162* is 10.3 kb which is roughly identical to the sum of *pBS1* and the larger *EcoRI* fragment of *pBC16* (10.5 kb). It appears therefore rather likely that a hybrid plasmid consisting of the large *EcoRI*




A B C D E F G H I K L M

**Fig. 1A-M.** Plasmids obtained after transformation of *B. subtilis* 168 with in vitro recombinant DNA consisting of *pBC16* and *pBS1*. Plasmid *pBC16* and *pBS1* were cleaved by *EcoRI* and ligated with T4 DNA ligase as described (Cohen et al., 1973). Transformation into *B. subtilis* was performed by the procedure of Cahn and Fox, 1968.  $Tc^R$ -colonies were picked, their covalently-closed circular DNAs isolated and analysed on a 1% agarose gel (2 mA/cm, 17 h). **A** *pBS1*, **B** *pBS1* cleaved with *EcoRI*, **C** *pBC16*, **D** *pBC16* cleaved with *EcoRI*, **E** *pBC16-1*, **F** *pBC16-1* cleaved with *EcoRI*, **G** *pBC16* and *pBC16-2*, **H** *pBC16* and *pBC16-2* cleaved with *EcoRI*, **J** *pBS161* and *pBS162*, **K** *pBS161* and *pBS162* cleaved with *EcoRI*, **L** *pBS161*, **M** *pBS161* cleaved with *EcoRI*



A B C D E F

**Fig. 2A-F.** Comparison of the *HindIII* and *HindIII/EcoRI* patterns of *pBS1* and *pBS161*. Covalently closed DNA of *pBS1* and *pBS161* was cleaved with *HindIII* and *HindIII+EcoRI*, respectively. The fragments were separated by electrophoresis on a 1% agarose gel. **A** *pBS1* CCC-DNA, **B** *pBS1* cleaved with *HindIII*, **C** *pBS1* cleaved with *HindIII* and *EcoRI*, **D** *pBS161* cleaved with *HindIII* and *EcoRI*, **E** *pBS161* cleaved with *HindIII*, **F** *pBS161* CCC-DNA.



A B C D E

**Fig. 3A-E.** Gel electrophoresis of plasmid *pBS161-1*, obtained by cleaving of *pBS161* with *HindIII* and transformation of the religated DNA into *B. subtilis* 168. **A** Covalently closed *pBS161-1* DNA, **B** *pBS161-1* DNA cleaved with *EcoRI*, **C** with *HindIII*, **D** with *EcoRI* and *HindIII* (the small fragment of 300 bp can not be seen on this gel). **E** *pBS161* cleaved with *HindIII*




Fig. 4. Genealogy of the *B. subtilis* vector plasmids *pBC16*, *pBC16-1*, *pBS161* and *pBS161-1*

fragment of *pBC16*, which carries the tetracycline resistance and the replication functions of *pBC16*, and *pBS1* was formed in vitro. However, after transformation in *B. subtilis* the recombinant plasmid which consists now of two replicons is unstable and dissociates nonreciprocally into *pBS161* and *pBS162*. *pBS161* was cleaved with *HindIII* and the fragments were recircularized in vitro with DNA ligase. After transformation in *B. subtilis* *Tc<sup>R</sup>*-colonies are obtained, that carry a new plasmid *pBS161-1* which represents the circularized *HindIII*-A fragment of *pBS161* (Fig. 3) which indicates that this fragment carries the tetracycline resistance determinant and the replication function of *pBS161*. This fragment has retained in addition one *EcoRI* site. Figure 4 summarizes the formation of the various derivatives of *pBC16*.


#### Construction of Hybrid Plasmids Consisting of *HindIII* Fragments of *pBS161* and *E. coli* Plasmids *pBR322*, *pWL7* or *pAC184*

Plasmid *pBS161* was digested by restriction endonuclease *HindIII* and ligated, using T4 DNA ligase, with the *E. coli* plasmids *pBR322* (Boyer et al., 1977), *pWL7*

(Kollek et al., 1978), or *pAC184* (S. Cohen and A. Chang, unpublished), which were also linearized by *HindIII*. These *E. coli* plasmids were used as vectors since insertion of a *HindIII* fragment into their single *HindIII* sites inactivates the tetracycline resistance (in the case of *pBR322* and *pAC184*) or the kanamycin resistance (in the case of *pWL7*), but leaves another antibiotic resistance determined by these plasmids intact which can then be used for selection of transformants (*Ap<sup>R</sup>* for *pBR322* and *pWL7*, and *Cm<sup>R</sup>* for *pAC184*). The in vitro ligated mixtures were transformed into *E. coli* 5K ( $\tau^-$ ,  $m^+$ ) and hybrid clones were selected by inactivation of the corresponding marker. When clones, obtained after transformation with ligated *HindIII* cleaved *pBR322* and *HindIII* fragments of *pBS161* were tested for inactivation of tetracycline resistance, two different types of clones were obtained: Clones which were completely sensitive to tetracycline and clones which still retained a low level of tetracycline resistance. Whereas strains carrying the original plasmid *pBR322* are resistant to 100  $\mu$ g/ml tetracycline, the upper limit of resistance of the latter clones is 15  $\mu$ g/ml.

Both, tetracycline sensitive and low-level tetracycline resistant colonies contained hybrid plasmids, as





**Fig. 5A-M.** Agarose gel electrophoresis of *Hind*III-cleaved hybrid plasmids consisting of *E. coli* plasmid *pBR322* and *Hind*III fragments of *pBS161*. Covalently closed hybrid DNA was isolated from the indicated strains and cleaved with *Hind*III. **A** *pBS161*-CCC-DNA, **B** *pBS161* cleaved with *Hind*III, **C** CCC-*pJK1* DNA from *E. coli* 5K, **D** cleaved with *Hind*III, **E** *pJK1* DNA from *E. coli* MM383 (*polA12*) cleaved with *Hind*III, **F** *pJK1* DNA from *B. subtilis* 168 cleaved with *Hind*III, **G** *pBR322* cleaved with *Hind*III. **H** CCC-*pBR322*, **J** CCC-*pJK2* DNA from *E. coli* 5K, **K** cleaved with *Hind*III. **L** CCC-*pJK3* DNA from *E. coli* 5K. **M** cleaved with *Hind*III

**Table 1.** Characterization of hybrid plasmids consisting of *E. coli* vectors *pBR322* or *pWL7* and various *Hind*III fragments of *B. subtilis* plasmid *pBS161*

Plasmid	Molecular weight ( $\times 1000$ b.p.)	<i>E. coli</i> vector	<i>Hind</i> III fragments of <i>pBS161</i> inserted into the <i>E. coli</i> vector	Transformation rate <sup>b</sup> in <i>B. subtilis</i> 168
<i>pJK1</i>	9.8	<i>pBR322</i>	A, B, E	$2.6 \times 10^{-5}$
<i>pJK2</i>	11.5 <sup>a</sup>	<i>pBR322</i>	A	$1.4 \times 10^{-5}$
<i>pJK3</i>	7.6	<i>pBR322</i>	A	$5.3 \times 10^{-5}$
<i>pJK4</i>	5.6	<i>pBR322</i>	B	0
<i>pJK101</i>	15.75	<i>pWL7</i>	A	$2.4 \times 10^{-5}$
<i>pBR322</i>	3.9	—	—	0
<i>pWL7</i>	12.1	—	—	0
<i>pBC16</i>	4.25	—	—	$2.0 \times 10^{-5}$
<i>pBC16-1</i>	2.7	—	—	$4.5 \times 10^{-5}$
<i>pBS161</i>	8.2	—	—	$1.9 \times 10^{-5}$

<sup>a</sup> This plasmid contains two molecules of *pBR322* linked to one *Hind*III-A fragment

<sup>b</sup> Transformation rate is defined as the ratio of  $Tc^R$  transformants per  $10^8$  cells and  $3 \mu\text{g}$  of DNA used in the transformation assay

demonstrated by cleaving the isolated CCC-DNA with *Hind*III (Fig. 5). Most plasmids isolated from clones which retained the low tetracycline resistance had the largest *Hind*III fragment (*Hind*III-A) of *pBS161* incorporated into *pBR322* (Table 1). Low-level tetracycline resistant colonies were also obtained when *pWL7* plasmid ( $Km^R$ ,  $Ap^R$ ) was used as vector for the cloning of the *Hind*III fragment of *pBS161*. The  $Tc^R$  colonies carried hybrid plasmids which had also *Hind*III fragment A incorporated into *pWL7*. Since plasmid *pWL7* does not determine tetracycline resis-

tance, this result indicates that the low tetracycline resistance is expressed by the cloned *Hind*III-A fragment of *pBS161*. One clone, also resistant to low concentrations of tetracycline, contained a hybrid plasmid, *pJK4*, which had fragment *Hind*III-B of *pBS161* incorporated into *pBR322* (Table 1). Since the *Hind*III site in *pBR322* inactivates the promoter of the  $Tc^R$  gene, but not the structural gene it was assumed that the tetracycline resistance of *pJK4* may be caused by the replacement of the inactive own promoter by a *B. subtilis* promoter located on fragment *Hind*III-B. Plasmid *pJK4* was therefore cleaved by *Hind*III, religated with DNA ligase in vitro and transformed into *E. coli* 5K. Ampicillin resistant transformants were obtained, which were either  $Tc^R$  or  $Tc^S$ . The  $Tc^S$  transformants contained *pJK4* with *Hind*III-B inserted in the opposite direction as in the original *pJK4*, whereas  $Tc^R$  transformants contained either *pJK4* in the original orientation or *pBR322* only (Fig. 6).

Purified hybrid plasmids obtained in the former cloning experiments in *E. coli* were used to transform competent *B. subtilis* cells. Selection for transformed colonies was performed on tetracycline containing plates.  $Tc^R$  transformants were obtained with all hybrid plasmids containing fragment *Hind*III-A from *pBS161*. Tetracycline resistance of these transformed *B. subtilis* strains is high and comparable to the tetracycline resistance ( $100 \mu\text{g}/\text{ml}$ ) of the parent plasmid *pBS161*. CCC-DNA isolated from the *B. subtilis* clones is identical to the hybrid plasmid DNA isolated from *E. coli* and used for the transformation, i.e. the plasmid DNA in both hosts remains identical (Fig. 5). Similar results were obtained with hybrid




Fig. 6A-G. Recloning of *pJK4* hybrid DNA. Hybrid *pJK4* DNA from low-level *Tc<sup>R</sup>* colonies of *E. coli* 5K was isolated. The DNA was cleaved with *Hind*III and religated with T4 DNA ligase. The DNA was transformed into *E. coli* 5K, *Tc<sup>R</sup>* and *Tc<sup>S</sup>*-colonies were obtained. Cleared lysates of *Tc<sup>R</sup>* and *Tc<sup>S</sup>* colonies were prepared and analysed by agarose gel electrophoresis. **A, B** cleared lysates from *Tc<sup>S</sup>*-clones containing *pJK4* DNA with *Hind*III-B fragment inserted into *pBR322* in the opposite orientation as in the original *pJK4* DNA. **C-E** cleared lysates from *Tc<sup>R</sup>*-clones containing *pBR322* DNA only, **F** cleared lysate from *Tc<sup>R</sup>*-clone containing the original *pJK4* DNA, **G** *pJK4* DNA as control

plasmids of *pWL7* having *Hind*III fragment A incorporated (data not shown).

Transformation of *B. subtilis* BR151-CM1 a non-sporulating derivative of *B. subtilis* 168 (kindly provided by P. Lovett) with *pBC16-1* and the hybrid plasmid *pJK101* leads also to tetracycline resistant colonies. The transformation rate is comparable to that of *B. subtilis* 168, indicating that this nonsporulating *B. subtilis* mutant can be also used as a host for recombinant DNA.

#### Replication Control of Hybrid Plasmids

The *E. coli* plasmids used as vectors for the described cloning experiments are *ColE1* derivatives. Their replication is therefore dependent on high dosis of DNA polymerase I (Kingsbury and Helinski, 1970) and continues in the presence of chloramphenicol (Clewell, 1972). To test which replicon of the obtained hybrid plasmids capable of replication in both hosts is functional, the following experiments were performed:

a) Transformation of *pJK1-3* (see Table 1) in *E. coli* DP9 *polA<sup>-</sup>*, *r<sup>-</sup>*, *m<sup>-</sup>* does not yield transformants. Transformants of MM383, *polA12* (carrying a temperature sensitive DNA polymerase I) obtained with this plasmid are unable to grow at 43° C on tetracycline and/or ampicillin containing plates.

b) Addition of chloramphenicol to a culture of *E. coli* harbouring *pJK1* leads to an amplification of this plasmid. However, the copy number of the hybrid plasmid increases only 3-5 fold after treatment with this antibiotic. Treatment of *B. subtilis* 168 harbouring *pJK1* with chloramphenicol leads to a complete stop of plasmid synthesis (Fig. 7). The experiment was performed with chloramphenicol concentrations between 5 and 150 µg/ml. The results indicate that in *E. coli* the *ColE1* replicon but not the *Bacillus subtilis* replicon is active whereas in *B. subtilis* only the *B. subtilis* but not the *ColE1* replicon is functioning.


#### Differences in Gene Expression of the Hybrid Plasmids in *E. coli* and *B. subtilis*

As described in the preceding section the gene determining resistance to tetracycline in *B. subtilis* is also expressed in *E. coli*, although the level of resistance in the new host is lower. Experiments to insert *in vitro* the *Eco*RI fragment of pML21 conferring resistance to kanamycin in *E. coli* (Hershfield et al., 1974) into *Eco*RI-cleaved plasmid *pBC16* of *B. subtilis* were performed. The ligated mixture which contained according to the analysis on an agarose gel at least 80% ligated DNA was used to transform *B. subtilis*. Although a large number of *Tc<sup>R</sup>*-transformants were obtained none of them were resistant to kanamycin.

The hybrid plasmids *pJK1-3* and *pJK101* which determine in *E. coli* resistance to ampicillin and tetracycline, showed resistance to tetracycline only, after transformation into *B. subtilis*. The lack of expression of the ampicillin resistance gene of *E. coli* in *B. subtilis* is not caused by a modification of the hybrid DNA in *B. subtilis*, since these plasmids isolated from *B. subtilis* and retransformed into *E. coli* lead again to *Ap<sup>R</sup>* and *Tc<sup>R</sup>* colonies.

To test whether  $\beta$ -lactamase of *pJK2* is synthesized in *B. subtilis* but not excreted, *B. subtilis* harbouring *pJK2* was analysed for intracellular  $\beta$ -lactamase activity by the method of Dale and Smith, 1971. No  $\beta$ -lactamase could be detected (Table 2).

Similar experiments were performed with hybrid plasmids consisting of *pBS161-1* (fragment *Hind*III-A of *pBS1*) and the *E. coli* vector plasmid *pAC184* deter-



**Fig. 7 A and B.** Cesium chloride-ethidium bromide gradients of DNA from *E. coli* 5K, *pJK3* and *B. subtilis* 168, *pJK3* synthesized in the absence and presence of chloramphenicol (Cm). Cultures of *E. coli* 5K and *B. subtilis* 168, carrying the hybrid plasmid *pJK3* were labeled at a cell density of  $2 \times 10^8$ /ml with  $10 \mu\text{Ci/ml}$   $^3\text{H}$ -thymidine (spec. act.  $24.3 \text{ Ci/mol}$ ). **A** *E. coli* 5K, *pJK3* without Cm, **A'** *E. coli* 5K, *pJK3* with  $160 \mu\text{g/ml}$  Cm (**A'**) **B** *B. subtilis*, *pJK3* without Cm (**B'**) **B** *B. subtilis*, *pJK3* with  $50 \mu\text{g/ml}$  Cm (other concentrations of Cm – 15 to  $150 \mu\text{g/ml}$  – gave similar results)


**Table 2.** Intracellular  $\beta$ -lactamase activity in *E. coli* 5K and *B. subtilis* 168 harbouring hybrid plasmid *pJK2*

Bacterial strain	Protein concentration (mg/ml)	Specific activity (units/mg protein) <sup>a</sup>
<i>E. coli</i> 5K <i>pJK2</i>	32	7.3
<i>B. subtilis</i> 168 <i>pJK2</i>	9	0
<i>B. subtilis</i> 168	9	0

<sup>a</sup> One enzyme unit hydrolyses one  $\mu\text{mole}$  Ampicillin/min at  $30^\circ \text{C}$  in 25 mM Tris-HCl pH 7.4

mining resistance to chloramphenicol and tetracycline (S. Cohen and A. Chang, unpublished). Transformation of these plasmids into competent cells of *B. subtilis* 168 leads to tetracycline resistant but chloramphenicol sensitive clones, whereas the same DNA transformed in *E. coli* yields  $Tc^R$  and  $Cm^R$  colonies.

After cultivation of *B. subtilis*-clones, which harbour hybrid plasmids consisting of DNA from *E. coli* and *B. subtilis* (*pJK1-3*), over a longer period of time it was observed that the hybrid plasmids were unstable in this host. CCC-DNA from *B. subtilis* harbouring such hybrids was now isolated, which represents smaller plasmids (Fig. 8). They have lost various parts of the *E. coli* DNA. Some have retained essentially only the genetic information of the *B. subtilis* replicon, i.e. the capability of autonomous replication and the tetracycline resistance of *B. subtilis*. A similar instability of these hybrids in *E. coli* has not been observed.



**Fig. 8 A–G.** Covalently closed DNA from *E. coli* and *B. subtilis* clones harbouring hybrid plasmids *pJK1-3* after cultivation for a longer period of time. CCC-DNA was either purified by CsCl-ethidium bromide gradient centrifugation (**A**, **C**, **E**) or cleared lysates were directly applied to the gel (**B**, **D**, **F**). **A** *pJK1* from *E. coli* 5K, **B** *pJK1* from *B. subtilis* 168, **C** *pJK2* from *E. coli* 5K, **D** *pJK2* from *B. subtilis* 168, **E** *pJK3* from *E. coli* 5K, **F** *pJK3* from *B. subtilis* 168. **G** *pBR322*

## Discussion

The plasmid *pBC16*, which was originally isolated from *Bacillus cereus* has been successfully used as a cloning vector in *B. subtilis*. The in vitro recombination experiments described in this report indicate, that the two important vector properties of this plasmid, namely the tetracycline resistance and the capability

of autonomous replication are located on an *EcoRI* fragment of 2.7 kb, which upon circularization results in a new autonomously replicating plasmid *pBC16-1*, which appears to be a suitable *EcoRI* vector for cloning of genes in *B. subtilis*.

Joining of this plasmid to another cryptic extrachromosomal element from *B. subtilis*, *pBS1*, does not lead to a stable bifunctional replicon as expected. Instead, the recombinant DNA dissociates in all clones studied into two new autonomously replicating elements *pBS161* and *pBS162*. The reason for this apparent instability of a recombinant DNA consisting of two functional replicons in *B. subtilis* is unknown. In *E. coli* such bifunctional replicons have been constructed (Cabello et al., 1976; Kollek et al., 1978; Mayer et al., 1977) and shown to be stable.

The tetracycline resistance remains, after dissociation, on the larger plasmid, *pBS161*, which carries only one *EcoRI* site and can be also used as *EcoRI* vector for *B. subtilis*. The plasmid has in addition six *HindIII* sites from *pBS1*. The largest *HindIII* fragment carries the tetracycline resistance and the genetic information required for autonomous replication. Upon circularization of this fragment an autonomously replicating plasmid *pBS161-1* is generated which represents a *HindIII* vector for *B. subtilis*.

Recombinant plasmids consisting of this *HindIII* fragment and an *E. coli* plasmid like *pBR322*, *pWL7* or *pAC184* are capable of stable replication in *B. subtilis* and *E. coli*. The dependency of replication of these recombinant plasmids on *polA* in *E. coli* and their inability to replicate in *B. subtilis* in the presence of chloramphenicol indicate that only the homologous replicon is functioning in the corresponding host.

The investigation of the gene expression of the recombinant DNAs in *E. coli* and *B. subtilis* indicates a rather interesting difference between *E. coli* and *B. subtilis*. Whereas the tetracycline resistance of the *B. subtilis* plasmid can be expressed in *E. coli*—although at a lower level than in the natural host—several antibiotic resistances from *E. coli* (kanamycin-, ampicillin- and chloramphenicol resistance) are not expressed in *B. subtilis*. Trivial explanations for this lack in heterologous gene expression in *B. subtilis* could be ruled out: Modification of the recombinant plasmid in *E. coli* or *B. subtilis* does not occur. The orientation of insertion of the fragment is of no influence in any case. It could also be shown that no TEM- $\beta$ -lactamase determined by the *E. coli* gene is synthesized inside the *B. subtilis* cell.

It remains to be seen whether this lack of heterologous gene expression is a general phenomenon in *B. subtilis*. Since gene expression in the opposite

tion, i.e. *B. subtilis* genes in *E. coli*, is possible (Ehrlich et al., 1976; Mahler and Halvorson, 1977) and the expression of other heterologous genes from Gram-positive and Gram-negative bacteria in *E. coli* has been repeatedly demonstrated (for a summary, see Vosberg, 1978), one may speculate that Gram-positive bacteria may have a more limited potential for expression of genetic information from Gram-negative than vice versa. There is also evidence that the heterologous DNA from the recombinant plasmids is more unstable in *B. subtilis* than in *E. coli*. The reason for this difference is also unknown.

It should be emphasized that not only the wild-type strain of *B. subtilis*, but also a non-sporulating mutant of it yields comparable transformation rates with the *Bacillus subtilis* vector plasmid alone and recombinant DNA constructed with it. This makes *B. subtilis* a rather safe host system for gene cloning.

While this manuscript was in preparation, we learned that similar results concerning the lack of expression of *E. coli* genes in *B. subtilis* were obtained by S.D. Ehrlich (PNAS, in press) using plasmids from *S. aureus* as cloning vehicle.

*Acknowledgements.* We thank S.D. Ehrlich for sending us his manuscript prior to publication. S. Cohen is thanked for providing the plasmid *pAC184*, Ph. Kourilsky for the *E. coli* strain DP9 *polA*<sup>-</sup>, *r*<sup>-</sup>, *m*<sup>-</sup> and P. Lovett for *B. subtilis* BR151-CM1.

We are thankful to Miss M. Vogel and Miss B. Schelle for skillful technical assistance.

This work was supported by a grant from the Deutsche Forschungsgemeinschaft (SFB 105-A1)

## References

- Bernhard, K., Schrempf, H., Goebel, W.: Bacteriocin and antibiotic resistance in *Bacillus cereus* and *Bacillus subtilis*. *J. Bact.* **133**, 897–903 (1978)
- Boyer, H.W., Betlack, M., Bolivar, F., Rodriguez, R.L., Heyneker, H.L., Shine, J., Goodman, H.M.: Recombinant molecules: Impact on science and society (Beers, R.F. and Bassett, E.B., eds.) pp. 9–20. New York: Raven Press 1977
- Cabello, F., Timmis, K., Cohen, S.N.: Replication control in a composite plasmid constructed by in vitro linkage of two distinct replicons. *Nature (Lond.)* **259**, 285–290 (1976)
- Cahn, F., Fox, M.: Fractionation of transformable bacteria from competent cultures of *B. subtilis* on renografin gradients. *J. Bact.* **95**, 869–875 (1968)
- Clewell, D.B.: Nature of ColE1 plasmid replication in *Escherichia coli* in the presence of chloramphenicol. *J. Bact.* **110**, 667–676 (1972)
- Cohen, S.N., Chang, A.C.Y., Boyer, H.W., Helling, R.B.: Construction of biologically functional bacterial plasmids in vitro. *Proc. nat. Acad. Sci. (Wash.)* **70**, 3240–3244 (1972)
- Dale, J.W., Smith, J.T.: The purification and properties of the  $\beta$ -lactamase specified by the resistance factor R-1818 in *E. coli* and *P. mirabilis*. *Biochem. J.* **123**, 493–500 (1971)
- Ehrlich, S.D.: Replication and expression of plasmids from *Staphylococcus aureus* in *Bacillus subtilis*. *Proc. nat. Acad. Sci. (Wash.)* **74**, 1680–1682 (1977)

- Ehrlich, S.D., Brusztyn-Pettegrew, H., Stroynowski, I., Lederberg, J.: Expression of the thymidylate synthetase gene of the *Bacillus subtilis* bacteriophage Phi-3-T in *Escherichia coli*. Proc. nat. Acad. Sci. (Wash.) **73**, 4145–4149 (1976)
- Goebel, W., Bonewald, R.: Class of small multicopy plasmids originating from the mutant antibiotic resistance factor R1*drd*-19B2. J. Bact. **123**, 658–665 (1975)
- Hershfield, V., Boyer, H.W., Yanofsky, C., Lovett, M., Helinski, D.R.: Plasmid ColE1 as a molecular vehicle for cloning and amplification of DNA. Proc. nat. Acad. Sci. (Wash.) **71**, 3455–3459 (1974)
- Kingsbury, D., Helinski, D.R.: DNA polymerase as a requirement for the maintenance of the plasmid colicinogenic factor E1. Biochem. biophys. Res. Commun. **41**, 1538–1544 (1970)
- Kollek, R., Oertel, W., Goebel, W.: Isolation and characterization of the minimal fragment required for autonomous replication (“basic replicon”) of a copy mutant (*pKN102*) of the antibiotic resistance factor R1. Molec. gen. Genet. **162**, 51–57 (1978)
- Mahler, I., Halvorson, H.O.: Transformation of *Escherichia coli* and *Bacillus subtilis* with a hybrid plasmid molecule. J. Bact. **131**, 374–377 (1977)
- Mayer, H., Luibrand, G., Goebel, W.: Replication of the mini-R1 plasmid Rsc11 and Rsc11 hybrid plasmids. Molec. gen. Genet. **152**, 145–152 (1977)
- Meyers, J., Sanchez, D., Elwell, L., Falkow, S.: Simple agarose gel electrophoretic method for the identification and characterization of plasmid deoxyribonucleic acid. J. Bact. **127**, 1529–1537 (1975)
- Vosberg, H.P.: Molecular cloning of DNA. An introduction into techniques and problems. Hum. Genet. **40**, (1978)

Communicated by F. Kaudewitz