

8. Literatur

- Alberola-Ila, J., Takaki, S., Kerner, J.D. & Perlmutter, R.M.** 1997. Differential signaling by lymphocyte antigen receptors. *Annu. Rev. Immunol.* **15**: 125-154.
- Ardouin, L. et al.** 1999. Crippling of CD3-zeta ITAMs does not impair T cell receptor signaling. *Immunity* **10**: 409-420.
- Arpaia, E. et al.** 1994. Defective T cell receptor signaling and CD8⁺ thymic selection in humans lacking zap-70 kinase. *Cell* **76**: 947-958.
- Aruffo, A. & Seed, B.** 1987. Molecular cloning of a CD28 cDNA by a high-efficiency COS cell expression system. *Proc. Natl. Acad. Sci. USA* **84**: 8573.
- Avraham, A. et al.** 1998. Co-stimulation-dependent activation of a JNK-kinase in T lymphocytes. *Eur. J. Immunol.* **28**: 2320-2330.
- Bischof, A. et al.** Autonomous induction of proliferation, JNK and NFκB activation in primary resting T-cells by mobilized CD28. *Eur. J. Immunol.* : submitted.
- Boise, L.H. et al.** 1995. CD28 costimulation can promote T cell survival by enhancing the expression of Bcl-XL. *Immunity* **3**: 87-98.
- Boucher, L.M. et al.** 1995. CD28 signals through acidic sphingomyelinase. *J. Exp. Med.* **181**: 2059-2068.
- Brennan, P. et al.** 1997. Phosphatidylinositol 3-kinase couples the interleukin-2 receptor to the cell cycle regulator E2F. *Immunity* **7**: 679-689.
- Brennan, P., Babbage, J.W., Thomas, G. & Cantrell, D.** 1999. p70(s6k) integrates phosphatidylinositol 3-kinase and rapamycin- regulated signals for E2F regulation in T lymphocytes. *Mol. Cell. Biol.* **19**: 4729-4738.
- Bretscher, P. & Cohn, M.** 1970. A theory of self-nonsel self discrimination. *Science* **169**: 1042-1048.
- Brunner, M.C. et al.** 1999. CTLA-4-Mediated inhibition of early events of T cell proliferation. *J. Immunol.* **162**: 5813-5820.
- Cai, Y.C. et al.** 1995. Selective CD28pYMNM mutations implicate phosphatidylinositol 3-kinase in CD86-CD28-mediated costimulation. *Immunity* **3**: 417-426.
- Calvo, C.R., Amsen, D. & Kruisbeek, A.M.** 1997. Cytotoxic T lymphocyte antigen 4 (CTLA-4) interferes with extracellular signal-regulated kinase (ERK) and Jun NH2-terminal kinase (JNK) activation, but does not affect phosphorylation of T cell receptor zeta and ZAP70. *J. Exp. Med.* **186**: 1645-1653.
- Cantrell, D.** 1996. T cell antigen receptor signal transduction pathways. *Annu. Rev. Immunol.* **14**: 259-274.

- Caplan, S., Zeliger, S., Wang, L. & Baniyash, M.** 1995. Cell-surface-expressed T-cell antigen-receptor zeta chain is associated with the cytoskeleton. *Proc. Natl. Acad. Sci. USA* **92**: 4768-4772.
- Chambers, C.A. & Allison, J.P.** 1999. Costimulatory regulation of T cell function. *Curr. Opin. Cell. Biol.* **11**: 203-210.
- Chan, A.C. et al.** 1994. ZAP-70 deficiency in an autosomal recessive form of severe combined immunodeficiency. *Science* **264**: 1599-1601.
- Chan, A.C. & Shaw, A.S.** 1996. Regulation of antigen receptor signal transduction by protein tyrosine kinases. *Curr. Opin. Immunol.* **8**: 394-401.
- Chan, G. & Ochi, A.** 1995. Sphingomyelin-ceramide turnover in CD28 costimulatory signaling. *Eur. J. Immunol.* **25**: 1999-2004.
- Chen, C.Y., Del Gatto-Konczak, F., Wu, Z. & Karin, M.** 1998. Stabilization of interleukin-2 mRNA by the c-Jun NH₂-terminal kinase pathway. *Science* **280**: 1945-1949.
- Cheng, A.M. et al.** 1995. Syk tyrosine kinase required for mouse viability and B-cell development. *Nature* **378**: 303-306.
- Cohen, G.B., Ren, R. & Baltimore, D.** 1995. Modular binding domains in signal transduction proteins. *Cell* **80**: 237-248.
- Collins, T.L., Deckert, M. & Altman, A.** 1997. Views on Vav. *Immunol. Today* **18**: 221-225.
- Crooks, M.E. et al.** 1995. CD28-mediated costimulation in the absence of phosphatidylinositol 3-kinase association and activation. *Mol. Cell. Biol.* **15**: 6820-6828.
- Danielian, S. et al.** 1992. Both T cell receptor (TcR)-CD3 complex and CD2 increase the tyrosine kinase activity of p56lck. CD2 can mediate TcR-CD3-independent and CD45-dependent activation of p56lck. *Eur. J. Immunol.* **22**: 2915-2921.
- DeBell, K.E. et al.** 1992. Microfilament assembly modulates phospholipase C-mediated signal transduction by the TCR/CD3 in murine T helper lymphocytes. *J. Immunol.* **149**: 2271-2280.
- DeSilva, D.R. et al.** 1997. The p38 mitogen-activated protein kinase pathway in activated and anergic Th1 cells. *Cell. Immunol.* **180**: 116-123.
- Elder, M.E. et al.** 1994. Human severe combined immunodeficiency due to a defect in ZAP-70, a T cell tyrosine kinase. *Science* **264**: 1596-1599.
- Fargnoli, J. et al.** 1995. Syk mutation in Jurkat E6-derived clones results in lack of p72syk expression. *J. Biol. Chem.* **270**: 26533-26537.
- Feder, D. & Bishop, J.M.** 1990. Purification and enzymatic characterization of pp60c-src from human platelets. *J. Biol. Chem.* **265**: 8205-8211.

- Finco, T.S. et al.** 1998. LAT is required for TCR-mediated activation of PLCgamma1 and the Ras pathway. *Immunity* **9**: 617-626.
- Firpo, E.J., Koff, A., Solomon, M.J. & Roberts, J.M.** 1994. Inactivation of a Cdk2 inhibitor during interleukin 2-induced proliferation of human T lymphocytes. *Mol. Cell. Biol.* **14**: 4889-4901.
- Fischer, K.D. et al.** 1998. Vav is a regulator of cytoskeletal reorganization mediated by the T- cell receptor. *Curr. Biol.* **8**: 554-562.
- Franke, T.F. & Cantley, L.C.** 1997. Apoptosis. A Bad kinase makes good. *Nature* **390**: 116-117.
- Franke, T.F., Kaplan, D.R. & Cantley, L.C.** 1997a. PI3K: downstream AKTion blocks apoptosis. *Cell* **88**: 435-437.
- Franke, T.F., Kaplan, D.R., Cantley, L.C. & Toker, A.** 1997b. Direct regulation of the Akt proto-oncogene product by phosphatidylinositol-3,4-bisphosphate. *Science* **275**: 665-668.
- Fraser, J., Irving, B.A., Crabree, G.R. & Weiss, A.** 1991. Regulation of the interleukin-2 gene enhancer activity by the T cell accessory molecule CD28. *Science* **251**: 313.
- Freeman, G.J. et al.** 1995. B7-1 and B7-2 do not deliver identical costimulatory signals, since B7- 2 but not B7-1 preferentially costimulates the initial production of IL- 4. *Immunity* **2**: 523-532.
- Fruman, D.A. et al.** 1999. Impaired B cell development and proliferation in absence of phosphoinositide 3-kinase p85alpha. *Science* **283**: 393-397.
- Garrington, T.P. & Johnson, G.L.** 1999. Organization and regulation of mitogen-activated protein kinase signaling pathways. *Curr. Opin. Cell. Biol.* **11**: 211-218.
- Gelfand, E.W. et al.** 1995. Absence of ZAP-70 prevents signaling through the antigen receptor on peripheral blood T cells but not on thymocytes. *J. Exp. Med.* **182**: 1057-1065.
- Ghaffari-Tabrizi, N. et al.** 1999. Protein kinase Ctheta, a selective upstream regulator of JNK/SAPK and IL-2 promoter activation in Jurkat T cells. *Eur. J. Immunol.* **29**: 132-142.
- Ghiotto-Ragueneau, M. et al.** 1996. Comparison of CD28-B7.1 and B7.2 functional interaction in resting human T cells: phosphatidylinositol 3-kinase association to CD28 and cytokine production. *Eur. J. Immunol.* **26**: 34-41.
- Gibson, S. et al.** 1996a. Functional LCK Is required for optimal CD28-mediated activation of the TEC family tyrosine kinase EMT/ITK. *J. Biol. Chem.* **271**: 7079-7083.
- Gibson, S. et al.** 1996b. The EMT/ITK/TSK (EMT) tyrosine kinase is activated during TCR signaling: LCK is required for optimal activation of EMT. *J. Immunol.* **156**: 2716-2722.

- Gibson, S. et al.** 1998. Efficient CD28 signalling leads to increases in the kinase activities of the TEC family tyrosine kinase EMT/ITK/TSK and the SRC family tyrosine kinase LCK. *Biochem. J.* **330**: 1123-1128.
- Gille, H. et al.** 1995. ERK phosphorylation potentiates Elk-1-mediated ternary complex formation and transactivation. *EMBO J.* **14**: 951-962.
- Han, J. et al.** 1998. Role of substrates and products of PI 3-kinase in regulating activation of Rac-related guanosine triphosphatases by Vav. *Science* **279**: 558-560.
- Hara, T., Fu, S.M. & Hansen, J.A.** 1985. Human T cell activation. II. A new activation pathway used by a major T cell population via a disulfide-bonded dimer of a 44 kilodalton polypeptide (9.3 antigen). *J. Exp. Med.* **161**: 1513-1524.
- Harder, T. & Simons, K.** 1997. Caveolae, DIGs, and the dynamics of sphingolipid-cholesterol microdomains. *Curr. Opin. Cell. Biol.* **9**: 534-542.
- Harder, T. & Simons, K.** 1999. Clusters of glycolipid and glycosylphosphatidylinositol-anchored proteins in lymphoid cells: accumulation of actin regulated by local tyrosine phosphorylation. *Eur. J. Immunol.* **29**: 556-562.
- Haslam, R.J., Koide, H.B. & Hemmings, B.A.** 1993. Pleckstrin domain homology. *Nature* **363**: 309-310.
- Henning, S.W. & Cantrell, D.A.** 1998. GTPases in antigen receptor signalling. *Curr. Opin. Immunol.* **10**: 322-329.
- Heyeck, S.D., Wilcox, H.M., Bunnell, S.C. & Berg, L.J.** 1997. Lck phosphorylates the activation loop tyrosine of the Itk kinase domain and activates Itk kinase activity. *J. Biol. Chem.* **272**: 25401-25408.
- Holdorf, A.D. et al.** 1999. Proline residues in CD28 and the Src homology (SH)3 domain of Lck are required for T cell costimulation. *J. Exp. Med.* **190**: 375-384.
- Huby, R.D., Weiss, A. & Ley, S.C.** 1998. Nocodazole inhibits signal transduction by the T cell antigen receptor. *J. Biol. Chem.* **273**: 12024-12031.
- Hutchcroft, J.E. et al.** 1995. Phorbol ester treatment inhibits phosphatidylinositol 3-kinase activation by, and association with, CD28, a T-lymphocyte surface receptor. *Proc. Natl. Acad. Sci. USA* **92**: 8808-8812.
- Hutchcroft, J.E., Tsai, B. & Bierer, B.E.** 1996. Differential phosphorylation of the T lymphocyte costimulatory receptor CD28. Activation-dependent changes and regulation by protein kinase C. *J. Biol. Chem.* **271**: 13362-13370.
- Jacinto, E., Werlen, G. & Karin, M.** 1998. Cooperation between Syk and Rac1 leads to synergistic JNK activation in T lymphocytes. *Immunity* **8**: 31-41.
- Jenkins, M.K. & Schwartz, R.H.** 1987. Antigen presentation by chemically modified splenocytes induces antigen-specific T cell unresponsiveness in vitro and in vivo. *J. Exp. Med.* **165**: 302-319.

- June, C.H., Bluestone, J.A., Nadler, L.M. & Thompson, C.B.** 1994. The B7 and CD28 receptor families. *Immunol. Today* **15**: 321.
- Jung, S. et al.** 1995. Costimulation requirement for AP-1 and NF-kappa B transcription factor activation in T cells. *Ann. N. Y. Acad. Sci.* **766**: 245-252.
- Kaga, S., Ragg, S., Rogers, K.A. & Ochi, A.** 1998. Activation of p21-CDC42/Rac-activated kinases by CD28 signaling: p21-activated kinase (PAK) and MEK kinase 1 (MEKK1) may mediate the interplay between CD3 and CD28 signals. *J. Immunol.* **160**: 4182-4189.
- Kästner, E.** Der vollständige Vers lautet: Zum neuen Jahr - Wird`s besser? Wird`s schlimmer? - fragt man alljährlich. - Seien wir ehrlich: Leben ist immer lebensgefährlich!
- Kempiak, S.J., Hiura, T.S. & Nel, A.E.** 1999. The Jun kinase cascade is responsible for activating the CD28 response element of the IL-2 promoter: proof of cross-talk with the I kappa B kinase cascade. *J. Immunol.* **162**: 3176-3187.
- King, A.J. et al.** 1998. The protein kinase Pak3 positively regulates Raf-1 activity through phosphorylation of serine 338. *Nature* **396**: 180-183.
- King, P.D. et al.** 1996. CD2 signaling in T cells involves tyrosine phosphorylation and activation of the Tec family kinase, EMT/ITK/TSK. *Int. Immunol.* **8**: 1707-1714.
- King, P.D. et al.** 1997. Analysis of CD28 cytoplasmic tail tyrosine residues as regulators and substrates for the protein tyrosine kinases, EMT and LCK. *J. Immunol.* **158**: 580-590.
- Klasen, S. et al.** 1998. Two distinct regions of the CD28 intracytoplasmic domain are involved in the tyrosine phosphorylation of Vav and GTPase activating protein-associated p62 protein. *Int. Immunol.* **10**: 481-489.
- Kuchroo, V. et al.** 1995. B7-1 and B7-2 costimulatory molecules activate differentially the Th1/Th2 developmental pathways: application to autoimmune disease therapy. *Cell* **80**: 707-718.
- Kupfer, A. & Singer, S.J.** 1989. Cell biology of cytotoxic and helper T cell functions: immunofluorescence microscopic studies of single cells and cell couples. *Annu. Rev. Immunol.* **7**: 309-337.
- Le Good, J.A. et al.** 1998. Protein kinase C isotypes controlled by phosphoinositide 3-kinase through the protein kinase PDK1. *Science* **281**: 2042-2045.
- Leitenberg, D., Boutin, Y., Lu, D.D. & Bottomly, K.** 1999. Biochemical association of CD45 with the T cell receptor complex: regulation by CD45 isoform and during T cell activation. *Immunity* **10**: 701-711.
- Lemmon, M.A., Ferguson, K.M. & Schlessinger, J.** 1996. PH domains: diverse sequences with a common fold recruit signaling molecules to the cell surface. *Cell* **85**: 621-624.

- Lenschow, D.J. et al.** 1995. Differential effects of anti-B7-1 and anti-B7-2 monoclonal antibody treatment on the development of diabetes in the nonobese diabetic mouse. *J. Exp. Med.* **181**: 1145-1155.
- Liao, X.C. et al.** 1997. Itk negatively regulates induction of T cell proliferation by CD28 costimulation. *J. Exp. Med.* **186**: 221-228.
- Liao, X.C. & Littman, D.R.** 1995. Altered T cell receptor signaling and disrupted T cell development in mice lacking Itk. *Immunity* **3**: 757-769.
- Lim, W.A. & Richards, F.M.** 1994. Critical residues in an SH3 domain from Sem-5 suggest a mechanism for proline-rich peptide recognition. *Nat. Struct. Biol.* **1**: 221-225.
- Liu, K.Q., Bunnell, S.C., Gurniak, C.B. & Berg, L.J.** 1998. T cell receptor-initiated calcium release is uncoupled from capacitative calcium entry in Itk-deficient T cells. *J. Exp. Med.* **187**: 1721-1727.
- Lu, Y., Phillips, C.A. & Trevillyan, J.M.** 1995. Phosphatidylinositol 3-kinase activity is not essential for CD28 costimulatory activity in Jurkat T cells: studies with a selective inhibitor, wortmannin. *Eur. J. Immunol.* **25**: 533-537.
- Marengere, L.E. et al.** 1997. The SH3 domain of Itk/Emt binds to proline-rich sequences in the cytoplasmic domain of the T cell costimulatory receptor CD28. *J. Immunol.* **159**: 3220-3229.
- Martiny-Baron, G. et al.** 1993. Selective inhibition of protein kinase C isozymes by the indolocarbazole Go 6976. *J. Biol. Chem.* **268**: 9194-9197.
- Mayer, B.J., Ren, R., Clark, K.L. & Baltimore, D.** 1993. A putative modular domain present in diverse signaling proteins. *Cell* **73**: 629-630.
- Metzler, W.J. et al.** 1997. Solution structure of human CTLA-4 and delineation of a CD80/CD86 binding site conserved in CD28. *Nat. Struct. Biol.* **4**: 527-531.
- Mishell, B.B. & Shiigi, S.M.** 1980. Selected methods in cellular immunology. Freeman and Company, San Fransisco.
- Molina, T.J. et al.** 1992. Profound block in thymocyte development in mice lacking p56lck. *Nature* **357**: 161-164.
- Monks, C.R. et al.** 1998. Three-dimensional segregation of supramolecular activation clusters in T cells. *Nature* **395**: 82-86.
- Monks, C.R. et al.** 1997. Selective modulation of protein kinase C-theta during T-cell activation. *Nature* **385**: 83-86.
- Moran, M. & Miceli, M.C.** 1998. Engagement of GPI-linked CD48 contributes to TCR signals and cytoskeletal reorganization: a role for lipid rafts in T cell activation. *Immunity* **9**: 787-796.
- Mueller, D.L., Jenkins, M.K. & Schwartz, R.H.** 1989. Clonal expansion versus functional clonal inactivation: a costimulatory signalling pathway determines the outcome of T cell antigen receptor occupancy. *Annu. Rev. Immunol.* **7**: 445-480.

- Nagasawa, M. et al.** 1997. Rapid nuclear translocation and increased activity of cyclin-dependent kinase 6 after T cell activation. *J. Immunol.* **158**: 5146-5154.
- Negishi, I. et al.** 1995. Essential role for ZAP-70 in both positive and negative selection of thymocytes. *Nature* **376**: 435-438.
- Nishina, H. et al.** 1997. Impaired CD28-mediated interleukin 2 production and proliferation in stress kinase SAPK/ERK1 kinase (SEK1)/mitogen-activated protein kinase kinase 4 (MKK4)-deficient T lymphocytes. *J. Exp. Med.* **186**: 941-953.
- Nunes, J.A. et al.** 1994. The role of p21ras in CD28 signal transduction: triggering of CD28 with antibodies, but not the ligand B7-1, activates p21ras. *J. Exp. Med.* **180**: 1067-1076.
- Nunes, J.A., Truneh, A., Olive, D. & Cantrell, D.A.** 1996. Signal transduction by CD28 costimulatory receptor on T cells. B7-1 and B7-2 regulation of tyrosine kinase adaptor molecules. *J. Biol. Chem.* **271**: 1591-1598.
- Okkenhaug, K. & Rottapel, R.** 1998. Grb2 forms an inducible protein complex with CD28 through a Src homology 3 domain-proline interaction. *J. Biol. Chem.* **273**: 21194-21202.
- Ostergaard, H.L., Lou, O., Arendt, C.W. & Berg, N.N.** 1998. Paxillin phosphorylation and association with Lck and Pyk2 in anti-CD3- or anti-CD45-stimulated T cells. *J. Biol. Chem.* **273**: 5692-5696.
- Pages, F. et al.** 1994. Binding of phosphatidylinositol-3-OH kinase to CD28 is required for T-cell signalling. *Nature* **369**: 327-329.
- Pai, S.Y., Calvo, V., Wood, M. & Bierer, B.E.** 1994. Cross-linking CD28 leads to activation of 70-kDa S6 kinase. *Eur. J. Immunol.* **24**: 2364-2368.
- Parry, R.V. et al.** 1997. Ligation of the T cell co-stimulatory receptor CD28 activates the serine-threonine protein kinase protein kinase B. *Eur. J. Immunol.* **27**: 2495-2501.
- Pawson, T.** 1995. Protein modules and signalling networks. *Nature* **373**: 573-580.
- Peach, R.J. et al.** 1994. Complementarity determining region 1 (CDR1)- and CDR3-analogous regions in CTLA-4 and CD28 determine the binding to B7-1. *J. Exp. Med.* **180**: 2049-2058.
- Penninger, J.M. & Crabtree, G.R.** 1999. The actin cytoskeleton and lymphocyte activation. *Cell* **96**: 9-12.
- Penninger, J.M. et al.** 1999. The oncogene product Vav is a crucial regulator of primary cytotoxic T cell responses but has no apparent role in CD28-mediated co-stimulation. *Eur. J. Immunol.* **29**: 1709-1718.
- Peterson, E.J., Clements, J.L., Fang, N. & Koretzky, G.A.** 1998. Adaptor proteins in lymphocyte antigen-receptor signaling. *Curr. Opin. Immunol.* **10**: 337-344.

- Prasad, K.V. et al.** 1994. T-cell antigen CD28 interacts with the lipid kinase phosphatidylinositol 3-kinase by a cytoplasmic Tyr(P)-Met-Xaa-Met motif. *Proc. Natl. Acad. Sci. USA* **91**: 2834-2838.
- Raab, M. et al.** 1995. p56Lck and p59Fyn regulate CD28 binding to phosphatidylinositol 3-kinase, growth factor receptor-bound protein GRB-2, and T cell-specific protein-tyrosine kinase ITK: implications for T-cell costimulation. *Proc. Natl. Acad. Sci. USA* **92**: 8891-8895.
- Raab, M., da Silva, A.J., Findell, P.R. & Rudd, C.E.** 1997. Regulation of Vav-SLP-76 binding by ZAP-70 and its relevance to TCR zeta/CD3 induction of interleukin-2. *Immunity* **6**: 155-164.
- Rayter, S.I. et al.** 1992. p21ras mediates control of IL-2 gene promoter function in T cell activation. *EMBO J.* **11**: 4549-4556.
- Reif, K. & Cantrell, D.A.** 1998. Networking Rho family GTPases in lymphocytes. *Immunity* **8**: 395-401.
- Ridgeway, N.D.** 1995. 25-Hydroxycholesterol stimulates sphingomyelin synthesis in Chinese hamster ovary cells. *J. Lipid. Res.* **36**: 1345-1358.
- Rodgers, W. & Rose, J.K.** 1996. Exclusion of CD45 inhibits activity of p56lck associated with glycolipid-enriched membrane domains. *J. Cell. Biol.* **135**: 1515-1523.
- Rodriguez-Palmero, M., Hara, T., Thumbs, A. & Hünig, T.** 1999. Triggering of T-cell proliferation through CD28 induces GATA-3 and promotes T-helper type 2 differentiation in vitro and in vivo. *Eur. J. Immunol.* : in press.
- Rozdzial, M.M., Malissen, B. & Finkel, T.H.** 1995. Tyrosine-phosphorylated T cell receptor zeta chain associates with the actin cytoskeleton upon activation of mature T lymphocytes. *Immunity* **3**: 623-633.
- Rudd, C.E.** 1996. Upstream-downstream: CD28 cosignaling pathways and T cell function. *Immunity* **4**: 527-534.
- Rudd, C.E.** 1999. Adaptors and molecular scaffolds in immune cell signaling. *Cell* **96**: 5-8.
- Rulifson, I.C. et al.** 1997. CD28 costimulation promotes the production of Th2 cytokines. *J. Immuno.* **158**: 658-665.
- Salmon, R.A., Foltz, I.N., Young, P.R. & Schrader, J.W.** 1997. The p38 mitogen-activated protein kinase is activated by ligation of the T or B lymphocyte antigen receptors, Fas or CD40, but suppression of kinase activity does not inhibit apoptosis induced by antigen receptors. *J. Immunol.* **159**: 5309-5317.
- Salojin, K.V., Zhang, J. & Delovitch, T.L.** 1999. TCR and CD28 are coupled via ZAP-70 to the activation of the Vav/Rac-1- /PAK-1/p38 MAPK signaling pathway. *J. Immunol.* **163**: 844-853.

- Schafer, P.H., Wadsworth, S.A., Wang, L. & Siekierka, J.J.** 1999a. p38 alpha mitogen-activated protein kinase is activated by CD28- mediated signaling and is required for IL-4 production by human CD4⁺CD45RO⁺ T cells and Th2 effector cells. *J. Immunol.* **162**: 7110-7119.
- Schafer, P.H. et al.** 1999b. T cell activation signals up-regulate p38 mitogen-activated protein kinase activity and induce TNF-alpha production in a manner distinct from LPS activation of monocytes. *J. Immunol.* **162**: 659-668.
- Scharenberg, A.M. & Kinet, J.P.** 1998. PtdIns-3,4,5-P3: a regulatory nexus between tyrosine kinases and sustained calcium signals. *Cell* **94**: 5-8.
- Schneider, H. et al.** 1995. T cell antigen CD28 binds to the GRB-2/SOS complex, regulators of p21ras. *Eur. J. Immunol.* **25**: 1044-1050.
- Schulze-Koops, H., Lipsky, P.E. & Davis, L.S.** 1998. Human memory T cell differentiation into Th2-like effector cells is dependent on IL-4 and CD28 stimulation and inhibited by TCR ligation. *Eur. J. Immunol.* **28**: 2517-2529.
- Serfling, E., Avots, A. & Neumann, M.** 1995. The architecture of the interleukin-2 promoter: a reflection of T lymphocyte activation. *Biochim. Biophys. Acta* **1263**: 181-200.
- Shahinian, A. et al.** 1993. Differential T cell costimulatory requirements in CD28-deficient mice. *Science* **261**: 609-612.
- Shaw, A.S. & Dustin, M.L.** 1997. Making the T cell receptor go the distance: a topological view of T cell activation. *Immunity* **6**: 361-369.
- Sicheri, F., Moarefi, I. & Kuriyan, J.** 1997. Crystal structure of the Src family tyrosine kinase Hck [see comments]. *Nature* **385**: 602-609.
- Siefken, R. et al.** 1998. A CD28-associated signaling pathway leading to cytokine gene transcription and T cell proliferation without TCR engagement. *J. Immunol.* **161**: 1645-1651.
- Siefken, R., Kurrle, R. & Schwinzer, R.** 1997. CD28-mediated activation of resting human T cells without costimulation of the CD3/TCR complex. *Cell. Immunol.* **176**: 59-65.
- Sieh, M., Bolen, J.B. & Weiss, A.** 1993. CD45 specifically modulates binding of Lck to a phosphopeptide encompassing the negative regulatory tyrosine of Lck. *EMBO J.* **12**: 315-321.
- Simons, K. & Ikonen, E.** 1997. Functional rafts in cell membranes. *Nature* **387**: 569-572.
- Sirim, P., Zeitlmann, L. & Kolanus, W.** 1999. Costimulation through the beta2-integrin LFA-1 amplifies T cell receptor signal transduction by an intracellular crosstalk mechanism. *Immunobiology* **200**: 365-366.

- Slavik, J.M., Hutchcroft, J.E. & Bierer, B.E.** 1999. CD80 and CD86 are not equivalent in their ability to induce the tyrosine phosphorylation of CD28. *J. Biol. Chem.* **274**: 3116-3124.
- Sperling, A.I. et al.** 1998. TCR signaling induces selective exclusion of CD43 from the T cell- antigen-presenting cell contact site. *J. Immunol.* **161**: 6459-6462.
- Stoffel, B. et al.** 1998. Ceramide-independent CD28 and TCR signaling but reduced IL-2 secretion in T cells of acid sphingomyelinase-deficient mice. *Eur. J. Immunol.* **28**: 874-880.
- Su, B. et al.** 1994. JNK is involved in signal integration during costimulation of T lymphocytes. *Cell* **77**: 727-736.
- Su, B. & Karin, M.** 1996. Mitogen-activated protein kinase cascades and regulation of gene expression. *Curr. Opin. Immunol.* **8**: 402-411.
- Suzuki, H. et al.** 1999. Xid-like immunodeficiency in mice with disruption of the p85alpha subunit of phosphoinositide 3-kinase. *Science* **283**: 390-392.
- Tacke, M.** 1995. Expression und Funktion des CD28-Moleküls in der Ratte. Doktorarbeit, Universität Würzburg.
- Tacke, M., Clark, G.J., Dallman, M.J. & Hünig, T.** 1995. Costimulatory function and cellular distribution of rat CD28: Regulated expression during thymocyte maturation. *J. Immunol.* **154**: 5121-5127.
- Tacke, M., Hanke, G., Hanke, T. & Hünig, T.** 1997. CD28-mediated induction of proliferation in resting T-cells in vitro and in vivo without engagement of the T-cell receptor: evidence for functionally distinct forms of CD28. *Eur. J. Immunol.* **27**: 239-247.
- Tao, X., Constant, S., Jorritsma, P. & Bottomly, K.** 1997. Strength of TCR signal determines the costimulatory requirements for Th1 and Th2 CD4+ T cell differentiation. *J. Immunol.* **159**: 5956-5963.
- Teng, J.M. et al.** 1996. Phosphorylation of each of the distal three tyrosines of the CD28 cytoplasmic tail is required for CD28-induced T cell IL-2 secretion. *Tissue Antigens* **48**: 255-264.
- Thomas, M.L.** 1999. The regulation of antigen-receptor signaling by protein tyrosine phosphatases: a hole in the story. *Curr. Opin. Immunol.* **11**: 270-276.
- Thompson, C.B.** 1995. Distinct roles for the costimulatory ligands B7-1 and B7-2 in T helper cell differentiation? *Cell* **81**: 979-982.
- Toullec, D. et al.** 1991. The bisindolylmaleimide GF 109203X is a potent and selective inhibitor of protein kinase C. *J. Biol. Chem.* **266**: 15771-15781.
- Truitt, K.E. et al.** 1995. CD28 delivers costimulatory signals independently of its association with phosphatidylinositol 3-kinase. *J. Immunol.* **155**: 4702-4710.

- Tsuchida, M., Manthei, E.R., Knechtle, S.J. & Hamawy, M.M.** 1999. CD28 ligation induces rapid tyrosine phosphorylation of the linker molecule LAT in the absence of Syk and ZAP-70 tyrosine phosphorylation. *Eur. J. Immunol.* **29**: 2354-2359.
- Tuosto, L. & Acuto, O.** 1998. CD28 affects the earliest signaling events generated by TCR engagement. *Eur. J. Immunol.* **28**: 2131-2142.
- Turner, M. et al.** 1995. Perinatal lethality and blocked B-cell development in mice lacking the tyrosine kinase Syk. *Nature* **378**: 298-302.
- Ueda, Y. et al.** 1995. Both CD28 ligands CD80 (B7-1) and CD86 (B7-2) activate phosphatidylinositol 3-kinase, and wortmannin reveals heterogeneity in the regulation of T cell IL-2 secretion. *Int. Immunol.* **7**: 957-966.
- van Leeuwen, J.E. & Samelson, L.E.** 1999. T cell antigen-receptor signal transduction. *Curr. Opin. Immunol.* **11**: 242-248.
- van Oers, N.S. et al.** 1996. alpha beta T cell development is abolished in mice lacking both Lck and Fyn protein tyrosine kinases. *Immunity* **5**: 429-436.
- Varma, R. & Mayor, S.** 1998. GPI-anchored proteins are organized in submicron domains at the cell surface. *Nature* **394**: 798-801.
- Viola, A. & Lanzavecchia, A.** 1996. T cell activation determined by T cell receptor number and tunable thresholds. *Science* **273**: 104-106.
- Viola, A., Schroeder, S., Sakakibara, Y. & Lanzavecchia, A.** 1999. T lymphocyte costimulation mediated by reorganization of membrane microdomains. *Science* **283**: 680-682.
- Wange, R.L. & Samelson, L.E.** 1996. Complex complexes: signaling at the TCR. *Immunity* **5**: 197-205.
- Ward, S. et al.** 1995. Inhibition of CD28-mediated T cell costimulation by the phosphoinositide 3-kinase inhibitor wortmannin. *Eur. J. Immunol.* **25**: 526-532.
- Ward, S.G., June, C.H. & Olive, D.** 1996. PI 3-kinase: a pivotal pathway in T-cell activation? *Immunol. Today* **17**: 187-197.
- Wardenburg, J.B. et al.** 1996. Phosphorylation of SLP-76 by the ZAP-70 protein-tyrosine kinase is required for T-cell receptor function. *J. Biol. Chem.* **271**: 19641-19644.
- Weiss, A., Koretzky, G., Schatzman, R.C. & Kadlecsek, T.** 1991. Functional activation of the T-cell antigen receptor induces tyrosine phosphorylation of phospholipase C-gamma 1. *Proc. Natl. Acad. Sci. USA* **88**: 5484-5488.
- Weiss, A. & Littman, D.R.** 1994. Signal transduction by lymphocyte antigen receptors. *Cell* **76**: 263-274.
- Werlen, G., Jacinto, E., Xia, Y. & Karin, M.** 1998. Calcineurin preferentially synergizes with PKC-theta to activate JNK and IL-2 promoter in T lymphocytes. *EMBO J.* **17**: 3101-3111.

- Wulfing, C. & Davis, M.M.** 1998. A receptor/cytoskeletal movement triggered by costimulation during T cell activation. *Science* **282**: 2266-2269.
- Wulfing, C., Sjaastad, M.D. & Davis, M.M.** 1998. Visualizing the dynamics of T cell activation: intracellular adhesion molecule 1 migrates rapidly to the T cell/B cell interface and acts to sustain calcium levels. *Proc. Natl. Acad. Sci. USA* **95**: 6302-6307.
- Wunderlich, L., Farago, A., Downward, J. & Buday, L.** 1999. Association of Nck with tyrosine-phosphorylated SLP-76 in activated T lymphocytes. *Eur. J. Immunol.* **29**: 1068-1075.
- Xavier, R. et al.** 1998. Membrane compartmentation is required for efficient T cell activation. *Immunity* **8**: 723-732.
- Xavier, R. & Seed, B.** 1999. Membrane compartmentation and the response to antigen [In Process Citation]. *Curr. Opin. Immunol.* **11**: 265-269.
- Xu, W., Harrison, S.C. & Eck, M.J.** 1997. Three-dimensional structure of the tyrosine kinase c-Src [see comments]. *Nature* **385**: 595-602.
- Yablonski, D., Kuhne, M.R., Kadlecsek, T. & Weiss, A.** 1998. Uncoupling of nonreceptor tyrosine kinases from PLC-gamma1 in an SLP-76- deficient T cell. *Science* **281**: 413-416.
- Yang, W.C., Ghiotto, M., Barbarat, B. & Olive, D.** 1999. The role of Tec protein-tyrosine kinase in T cell signaling. *J. Biol. Chem.* **274**: 607-617.
- Yang, W.C. & Olive, D.** 1999. Tec kinase is involved in transcriptional regulation of IL-2 and IL-4 in the CD28 pathway. *Eur. J. Immunol.* **29**: 1842-1849.
- Zhang, J. et al.** 1999a. p38 mitogen-activated protein kinase mediates signal integration of TCR/CD28 costimulation in primary murine T cells. *J. Immunol.* **162**: 3819-3829.
- Zhang, W. et al.** 1999b. Functional analysis of LAT in TCR-mediated signaling pathways using a LAT-deficient Jurkat cell line. *Int. Immunol.* **11**: 943-950.
- Zhang, W. et al.** 1998. LAT: the ZAP-70 tyrosine kinase substrate that links T cell receptor to cellular activation. *Cell* **92**: 83-92.