

## Moesin: A Cell Membrane Protein Linked with Susceptibility to Measles Virus Infection

LEE M. DUNSTER,\*<sup>1</sup> JÜRGEN SCHNEIDER-SCHAULIES,\* SIEGLINDE LÖFFLER,\* WOLFGANG LANKES,†  
REINHARD SCHWARTZ-ALBIEZ,‡ FRIEDRICH LOTTSPEICH,§ AND VOLKER TER MEULEN\*

\*Institut für Virologie und Immunbiologie, Versbacher Strasse 7, D-97078 Würzburg, Germany; †Medizinische Klinik I, Klinikum der RWTH Aachen, Pauwel Strasse 30, D-52074 Aachen, Germany; ‡Deutsches Krebsforschungszentrum, Im Neuenheimer Feld 280, D-69120 Heidelberg, Germany; and §Max Planck Institut für Biochemie, Am Klopferspitz, D-82152 Martinsried, Germany

Received July 27, 1993; accepted September 17, 1993

Measles virus is a highly contagious virus causing acute and persistent diseases in man, the receptor of which is still not well characterized. We have isolated a monoclonal antibody (mAb), designated mAb 119, which specifically inhibits measles virus infection of susceptible cell lines in a dose-dependent manner. This antibody precipitates a protein with an apparent molecular mass of 75 kDa from <sup>125</sup>I surface-labeled cells and its epitope is present on human peripheral blood mononuclear cells, human cell lines, and the African green monkey cell line Vero. Affinity chromatography of detergent-solubilized cell membrane proteins over a Sepharose column with covalently bound mAb 119 led to the partial purification of the 75-kDa protein. Preincubation of measles virus with this affinity-purified protein inhibited measles virus infection dose dependently. Amino acid microsequencing of this protein revealed its identity with the human membrane-organizing extension spike protein moesin, a protein intra- and extracellularly associated with the plasma membrane of cells. Subsequently, an antibody raised against purified moesin (mAb 38/87) was also found to specifically inhibit measles virus infection of susceptible cells and confirmed our data obtained with mAb 119. Our data suggest that moesin is acting as a receptor for measles virus. © 1994 Academic Press, Inc.

### INTRODUCTION

Measles virus (MV) is a highly successful pathogen and is responsible for the disease measles, classically a childhood community-related disease. The acute disease, which is largely self-limiting, gives life long immunity to MV, a situation which is aided by the existence of a single immunotype of MV. However, life threatening diseases can occur following exposure to MV and include acute measles encephalitis, measles inclusion body encephalitis (MIBE), and subacute sclerosing panencephalitis (SSPE), which target the central nervous system with devastating effect. The tropism of the virus in acute measles is directed largely to epithelial cells of the respiratory tract and cells of the immune system, i.e., lymphocytes and macrophages (for review see Casali *et al.*, 1989). By comparison, in brains of SSPE and MIBE patients MV has been detected mainly in neurons, but also in oligodendrocytes, astrocytes, microglial cells, and endothelial cells (Esiri *et al.*, 1981; Kirk *et al.*, 1991; Haase *et al.*, 1985; Moench *et al.*, 1988; Schneider-Schaulies and ter Meulen, 1992).

As a member of the Morbilliviruses, MV possesses an envelope that surrounds the viral nucleocapsid which contains one copy of a single-stranded negative sense RNA genome. The virus envelope has two virus-encoded glycoproteins inserted into it: the hemagglutinin

(H) protein, which mediates both virus attachment to susceptible cells and the biological activity of hemagglutination, while the fusion (F) protein, in conjunction with MV-H, is responsible for cell entry and fusion (Wild *et al.*, 1991). Natural MV infection is limited to humans, although monkeys can be infected under experimental conditions, thus suggesting the presence of a MV cell receptor that is unique to higher primates.

The primary event for successful infection of a cell by the virus involves attachment to the target cell and subsequent uptake of virus particles. This process is usually, although not exclusively, mediated by a specific virus receptor on the cell surface, which, depending on their distribution, may influence virus tropism (White and Littman, 1989). Viral receptors have been established for a growing number of viruses and recent evidence suggests that many viruses can utilize additional and/or alternative receptors for the attachment to cells (Dveskler *et al.*, 1993; Harouse *et al.*, 1989, 1991; Lentz, 1990; Schneider-Schaulies *et al.*, 1992).

For MV there have been a number of studies performed in an attempt to define its cellular receptor (Krah 1989, 1991; Harrowe *et al.*, 1990, 1992; Yoshikawa *et al.*, 1991). Putative receptors were found to be present on human cells, Vero cells, and monkey erythrocytes and most likely to consist of glycosylated proteins. Identification by gel electrophoresis of the protein species involved in attachment of MV to susceptible cells was, unfortunately, largely inconclusive (Krah,

<sup>1</sup> To whom reprint requests should be addressed.

1991). However, in a recent study, a monoclonal antibody (mAb) was developed against a specific human cell surface protein which was found to inhibit MV infection (Naniche *et al.*, 1992). The mAb immunoprecipitated a protein with the apparent molecular mass of 57/67 kDa. The protein, which was designated gp57/67, was subsequently identified as the membrane co-factor protein or CD46 (Naniche *et al.*, 1993b). However, CD46 is not present on all cells susceptible to MV infection, suggesting the presence of an additional receptor species. In this study we have characterized a protein which is immunoprecipitated by a mAb, which specifically inhibits to a high degree the infection of a wide range of cells by MV. The immunoaffinity-purified protein binds MV and has been identified as the membrane-organizing extension spike protein, moesin. These data implicate moesin as a further receptor species for attachment and uptake of MV into susceptible cells.

## MATERIALS AND METHODS

### Cell culture

Adherent cell lines (human astrocytoma U-251, human cervical carcinoma HeLa, human neuroblastoma SKN-MC, African green monkey kidney Vero) were propagated and maintained in Eagles minimal essential medium (EMEM) supplemented with 5% fetal calf serum. Human monocyte cell line U-937 and primary human peripheral blood mononuclear cells (PBMC) were maintained in RPMI 1640 supplemented with 10% fetal calf serum, 2.0 g/liter sodium bicarbonate, 0.1 g/liter penicillin, 0.1 g/liter streptomycin, 1 mM sodium pyruvate, nonessential amino acids, BME vitamins and 2 mM L-glutamine (all Seromed).

### Viruses

The Edmonston strain of MV was used throughout the experiments with virus stocks prepared according to Schneider-Schaulies *et al.* (1993). Additional viruses (canine distemper virus CDV, poliovirus serotypes 1, 2, and 3, Coxsackie viruses B2 and B4, and Echovirus 11) were propagated as above, except for the freeze-thaw step which was omitted for the enteroviruses.

### Culture and selection of monoclonal antibodies

Monoclonal antibodies were raised against cell surface epitopes by inoculating BALB/c mice intraperitoneally (i.p.) with  $1-2 \times 10^6$  U-251 cells. Four weeks later the mice were boosted with a further i.p. inoculation of U-251 cells and followed 1 month later by a final challenge intravenously, 3-4 days prior to harvesting the spleen. BALB/c spleen cells were fused with the myeloma cell line Sp2/0-Ag14 and seeded into 96-well dishes for selection of hybridomas. Hybridoma cells

were grown in RPMI 1640 medium as described earlier.

Screening of hybridoma cells for those secreting mAbs that inhibit infection by MV was performed in microtiter plates. Supernatants of hybridoma cultures were added to confluent Vero cells ( $3 \times 10^4$  per well) in microtiter plates 30 min prior to infection with MV (m.o.i. = 0.1) and incubated for 2 days at 37°. Inhibition of virus-induced cytopathic effect was observed microscopically. Protein-G column-purified mAbs were stored at -20° at a concentration of 1 mg/ml for use. The mAb 38/87 raised against the membrane-organizing extension spike protein, moesin, was described (Lankes *et al.*, 1988). Antibodies to MHC class I (W6/32) and CD58 (LFA-3) were purchased (Dianova) and mAbs to MV-H (K83), MV-F (A504), and U85 (raised against U-251 cells) were prepared in the laboratory and purified over protein-G columns.

### Plaque formation inhibition test

This test was performed on both Vero and U-251 cell monolayers in six-well plastic dishes ( $1 \times 10^6$  cells per well). Confluent cell monolayers were washed with PBS and 100  $\mu$ l of hybridoma cell supernatant (clarified by centrifugation at 300 g for 5 min) or specific concentrations of purified antibodies were added. After incubation at room temperature for 45 min, the monolayers were washed once with PBS and 100  $\mu$ l of PBS containing 100 PFU of MV was added to each test well. Further incubation at room temperature for 1 hr was followed by washing with PBS prior to overlaying the wells with 5 ml of EMEM supplemented with 1% fetal calf serum and 1% Noble agar (Sigma). Plaques were visualized after 3-5 days of incubation at 37° by staining with neutral red.

### Rosetting assay

Rosetting assays were made with  $10^5$  U-937 cells in 100  $\mu$ l culture medium and incubated for 1 hr at 4° with either mAb 38/87 or mAb 119 or with the control mAbs W6/32 (MHC class 1 specific) and U85 (nonbinding) (all 0.5 mg/ml). After washing with PBS, 50 hemagglutinating units of MV was added followed by incubation for 1 hr at 4°. The cells were washed with PBS and resuspended in 200  $\mu$ l PBS containing monkey erythrocytes at a final concentration of 0.1% and incubated for 1 hr at 37°. Rosettes were defined microscopically as cells with three or more erythrocytes attached to the surface and results normalized to 0% inhibition against the nonbinding mAb U85.

### Inhibition of vaccinia virus MV-H/F-mediated fusion

As an additional rapid screening assay for infection inhibiting mAbs we used a vaccinia virus double recombinant encoding both MV-H and -F proteins (VV-H/F,

Wild *et al.*, 1991; a kind gift from T. F. Wild, Lyon, France). mAbs were screened for their ability to inhibit the rapid and characteristic cell fusion associated with the expression of MV-H and -F proteins. Briefly, cells (U-251, HeLa or Vero) were seeded into 48-well cluster plates ( $1 \times 10^5$  per well) and grown to confluency. The monolayers were washed with PBS and dilutions of mAb supernatant added to each test well and incubated at room temperature for 30 min. After incubation, VV-H/F at an m.o.i. of 0.1 PFU was added and incubation for 18 hr at  $37^\circ$  made before fixing the cell monolayers with 3.5% paraformaldehyde in PBS and determining the inhibition of syncytium formation under the microscope.

#### Flow cytometry analysis

Flow cytometric analysis was performed as described (Schneider-Schaulies *et al.*, 1992). Briefly,  $2 \times 10^5$  cells were incubated for 45 min on ice with 200  $\mu$ l mAb. Cells were washed three times in FACS buffer (PBS containing 0.4% BSA and 0.02% sodium azide) and incubated with 200  $\mu$ l of a 1:100 dilution of fluorescein isothiocyanate-conjugated goat anti-mouse immunoglobulin (Dako) on ice for a further 45 min. After three washes with FACS buffer as before, flow cytometric analysis was performed on a FACScan (Becton Dickinson). For intracellular staining of epitopes, cells were fixed and permeabilized prior to application of the first antibody with 3.7% paraformaldehyde, 0.25% Triton X-100 in PBS for 5 min and unspecific binding was blocked by FACS buffer containing 10% BSA.

#### Surface labeling of cells with $^{125}$ I

Surface iodination of cells ( $1 \times 10^7$  cells/500  $\mu$ l) with 500  $\mu$ Ci  $^{125}$ I was performed using the Iodogen method and the manufacturers protocol (Pierce). Following iodination, the reaction was quenched with 30 mM sodium iodide after which the cells were washed three times with PBS and resuspended in lysis buffer (10 mM Tris, pH 7.4, 150 mM sodium chloride, 0.5% deoxycholate, 0.5% Triton X-100, 0.05% SDS) containing protease inhibitors (10 U aprotinin, 1 mM PMSF). After vigorous mixing for 30 sec and incubation on ice for 15 min, the lysates were centrifuged at 15,000  $g$  for 30 min at  $4^\circ$  and the supernatant was stored at  $-70^\circ$  for use.

#### Immunoprecipitations

Immunoprecipitations were performed according to Schneider-Schaulies *et al.* (1992) with the following modifications. All lysates were precleared prior to immunoprecipitation by the addition of 5  $\mu$ l normal mouse serum and incubated on ice for 1 hr. Mouse immunoglobulins were precipitated by adding 30  $\mu$ l of packed protein A-Sepharose beads (Pharmacia) bound to rabbit anti-mouse immunoglobulins for 30 min on ice fol-

lowed by centrifugation. Precipitates were analyzed by separation on 10% SDS-containing polyacrylamide gels (SDS-PAGE) and exposure of the dried gels to a phosphor imager (Molecular Dynamics) screen for 16 hr.

#### Cell membrane preparation

All procedures for the preparation of cell membranes were performed at  $4^\circ$  in the presence of protease inhibitors to minimize any degradation of sensitive protein structures. Anchorage-dependent cells were removed nonenzymatically from the culture flask with  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  free PBS plus 1 mM EDTA, while suspension cells were pelleted by low speed centrifugation (200  $g$  for 5 min). Cells were washed twice with  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  free PBS before resuspending the pellet in Buffer I ( $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  free PBS, 10% sucrose, 2 mM EDTA, and 1 mM PMSF) and centrifuging for 1 hr at 22,000 rpm in an SW55 rotor (Beckman) to pellet vesicles. The pellet was gently resuspended in hypotonic Buffer II (50 mM phosphate buffer, pH 7.4, 1 mM PMSF) and centrifuged for 30 min at 14,000 rpm in an SW40 rotor (Beckman) to allow for the increased volume of buffer. The resulting pellet was resuspended in lysis buffer (PBS plus 1.5% octylglucoside, 1 mM PMSF) at a ratio of 3 ml/g wet cell mass and incubated on ice for 1 hr. The lysate was centrifuged at 35,000 rpm for 1 hr in an SW55 rotor (Beckman), after which the supernatant was carefully removed and dialyzed overnight against PBS with 0.1% octyl glucoside before aliquoting and storage at  $-70^\circ$ . Total protein concentration was determined using the bicinchoninic acid method (Sigma) with typical yields from 1 g of wet cell mass equalling 10–15 mg/ml.

#### Immunoaffinity purification of proteins

Specific immunoaffinity purification of cell membrane proteins by mAb was achieved using standardized protocols from manufacturers. Protein-G-purified mAb was linked to activated Sepharose 4B beads (Pharmacia) to form an immunoaffinity matrix. Detergent-extracted cell membranes (see above) were then passed over the mAb column at  $4^\circ$  and bound proteins eluted with a sodium chloride step gradient. The eluted fractions were dialyzed against PBS (for biological analysis) or 10 mM Tris (for biochemical analysis).

#### Amino acid sequencing

For direct N-terminal amino acid sequencing, the affinity-purified protein was applied to a 10% SDS-PAGE gel, separated, and analyzed after electroblotting onto Glassybond (Biometra, Göttingen) according to Eckerskorn *et al.*, 1988. For internal sequencing the protein was cleaved in the gel with protease Lys C (Boehringer, Tutzing), the peptides eluted and separated by reversed-phase HPLC on Superspher 60 RP select B,




FIG. 1. Inhibition of MV infection by mAb 119. The reduction in virus titer following pretreatment of cell monolayers with mAb 119 was determined by plaque inhibition assay (100 PFU test). The percentage of infectivity was determined by comparison to mock-treated (PBS) cell monolayers for each virus. An inhibition of infection was specifically detected for MV, but not for CDV, Polio 1, 2, 3, Echo 11, and Coxsackie B2, B4.

125 × 2 mm (Eckerskorn and Lottspeich, 1989). Amino acid sequence analyses were performed on a 477A pulse liquid phase sequencer equipped with a 120A PTH analyzer (Applied Biosystems, Weiterstadt).

## RESULTS

### Isolation of antibodies inhibiting MV infection

BALB/c mice were immunized with human astrocytoma cell line U-251, these cells being highly susceptible to infection with MV strain Edmonston and supporting its replication to high titers ( $10^6$  PFU/ml). Approximately 3000 antibody producing hybridoma clones were assayed for their ability to inhibit MV infection using a rapid infection inhibition test in microtiter plates (see Materials and Methods). One clone which efficiently inhibited MV infection, and also recognized an epitope on the surface of U-251 cells as determined by flow cytometry (not shown), was subcloned three times by limiting dilution until a stable antibody producing clone was isolated. The mAb secreted by this clone was designated mAb 119. In the following experiments the mAb was used as a purified fraction from a protein-G column.

Inhibition of MV infection following treatment of cell monolayers with mAb 119 was found to be specific for MV alone after repeating 100 PFU plaque reduction tests with a number of different viruses. The data in Fig. 1 show that MV infection of Vero cells is inhibited by 92% after pretreatment of the monolayer with mAb

119. A similar level of inhibition of MV infection (90%) by mAb 119 was observed on U-251 cells (data not shown). In comparison, no inhibition of virus infection was seen for the related morbillivirus canine distemper virus (CDV), polio serotypes 1, 2, and 3, Echovirus 11, and Coxsackie viruses B2 and B4 in similar assays. Therefore, mAb 119 exhibits properties characteristic and coincident with a receptor-blocking antibody and represents an ideal instrument with which to identify the cell surface protein involved in this specific interaction.

### Immunoprecipitation of a 75-kDa protein with mAb 119

To determine the molecular mass of the putative MV cell receptor molecule, surface proteins of the human monocyte cell line U-937, which expresses high levels of the mAb 119 epitope, were labeled with  $^{125}$ I. mAb 119 precipitated a protein of an apparent molecular weight of 75 kDa from U-937 cells (Fig. 2, lanes 4 and 5). A protein of the same apparent molecular weight could be precipitated from HeLa, U-251, and Vero cells (data not shown). Because the cells were radiiodinated, the protein precipitated by mAb 119 is expected to be expressed on the surface of cells and hence be able to interact with external ligands. By Western blot analysis of U-937 cell lysates, mAb 119 was unable to




FIG. 2. Immunoprecipitation of  $^{125}$ I-labeled cell surface proteins with mAb 119. Cell surface proteins of U-937 cells were iodinated and cell lysates immunoprecipitated with mAb 119 and an antibody to MHC class I as control. The precipitates were separated by 10% SDS-PAGE and exposed to a phosphor imager screen for 16 hr. Lane 1,  $^{14}$ C-labeled protein markers; lane 2, U-937 cell lysate precipitated with PBS as negative control; lane 3, U-937 cell lysate precipitated with the MHC class I-specific mAb W6/32; lane 4, U-937 cell lysate precipitated with mAb 119 (10  $\mu$ g/ml); lane 5, U-937 cell lysate precipitated with mAb 119 (100  $\mu$ g/ml). The mAb 119 specifically precipitated a molecule with the apparent molecular mass of 75 kDa.

detect the 75-kDa protein, suggesting that protein conformation is vital (data not shown).

#### Affinity purification of putative receptor molecules

In order to identify the protein precipitated by mAb 119 by microsequencing and to further characterize it biologically, it was necessary to obtain large amounts of purified protein. To achieve this, mAb 119 was covalently bound to activated Sepharose to form a matrix for affinity chromatography. Detergent (octyl glucoside) solubilized cell membranes from U-937 cells were circulated over the column for 18 hr at 4° prior to washing the column free of unbound proteins. Elution of bound proteins from the column was achieved using a NaCl step gradient and the eluate was then further concentrated with polyethylene glycol. The proteins were analyzed by SDS-PAGE and stained with Coomassie blue (Fig. 3). The predominant protein species was eluted with 300 mM NaCl and possessed the same apparent molecular mass as the immunoprecipitated (75 kDa) protein (Fig. 3, lanes 4 and 6). The yield of purified protein from the immunoaffinity column was typically 200–250  $\mu$ g per 20 mg total protein loaded.

#### Biological activity of the enriched 75-kDa protein

The immunoaffinity-purified protein likely contains a receptor binding motif for MV and by inference should effectively bind MV in solution, if protein conformation




FIG. 3. Immunoaffinity purification of the 75-kDa protein. Detergent-derived cell lysate from U-937 cells was passed over an immunoaffinity column supporting mAb 119. Proteins binding to the affinity column were eluted by a NaCl step gradient and analyzed by gel electrophoresis and subsequent staining with Coomassie blue. Lane 1, molecular weight markers; lane 2, detergent-solubilized U-937 cell membrane preparation; lane 3, unbound proteins after 16 hr circulation over the column; lane 4, 300 mM NaCl eluate; lane 5, 500 mM NaCl eluate; lane 6, polyethylene glycol-concentrated 300 mM eluate; lane 7, polyethylene glycol-concentrated 500 mM eluate.




FIG. 4. Reduction of the infectivity of MV after incubation with soluble immunoaffinity enriched 75-kDa protein. The immunoaffinity column-enriched 75-kDa protein and bovine serum albumin (BSA) as control were diluted in PBS to yield a range of concentrations between 0 and 2 mg/ml. Dilutions of MV were mixed with a constant protein concentration and incubated for 1 hr on ice, and residual infectivity was determined by the plaque assay. The results obtained were normalized by setting the infectivity obtained in the presence of BSA to 100%. The solid line represents MV titrated against the 75-kDa protein, whereas the broken line represents BSA plus MV.

has been maintained. Therefore, the ability of the enriched 75-kDa protein to bind MV was assayed by infection inhibition tests. MV was titrated against constant concentrations of the 75-kDa protein diluted in PBS and incubated for 1 hr on ice. As control, the 75-kDa protein was replaced with similar concentrations of bovine serum albumin (BSA). After incubation, the residual infectivity was determined by the standard plaque assay. End-point titration of MV against the purified protein yielded an accurate estimate of the degree of virus binding. A reduction in virus infectivity between 0.95 and 1.1  $\log_{10}$  PFU/ml was observed for MV incubated with purified protein of concentration 200 mg/ml (Fig. 4). This represents an overall loss of MV infectivity due to binding to protein of approximately 90%. Lower concentrations of purified protein gave less dramatic reductions in MV infectivity titer (Fig. 4). Incubation of MV with BSA under the same conditions did not reduce the virus infectivity titer. Likewise, similar assays performed with the morbillivirus CDV revealed no reduction in the virus infectivity titer (data not shown).

#### Amino acid sequencing of the isolated protein

N-terminal amino acid microsequencing was performed after separation of the eluted proteins from the mAb 119 immunoaffinity column by SDS-PAGE containing approximately 50–100  $\mu$ g of the 75-kDa protein. The Coomassie-stained 75-kDa band was excised and digested in the gel with protease Lys C and blotted onto Glassybond (Biometra, Göttingen) for sequencing of internal peptides. Three peptides, one at the amino terminus and two internal, were sequenced.




FIG. 5. Dose-dependent inhibition of MV infection after pretreatment of cell monolayers with mAb 119 and mAb 38/87. Vero cell monolayers ( $1 \times 10^6$  cells/well) were incubated with dilutions of mAbs 119 and 38/87 in PBS as indicated and washed with PBS prior to infection with MV. The percentage of inhibition of plaque formation was plotted against the concentration of mAbs 119 (solid line) and 38/87 (broken line). Mean values of three experiments with standard deviations are shown.

Comparison of the sequences of the peptides with the sequences available in the EMBL data base showed that all three peptides were identical to peptides of the cloned human protein moesin (membrane-organizing extension spike protein, Lankes and Furthmayer, 1991). The peptides correspond to amino acids 1–8 (PKTISVRV), 132–137 (KYGDFN), and 500–513 (KDR-SEEERTTEAEK) of human moesin which is comprised of 576 amino acids in total with a calculated molecular mass of 67 kDa.

#### Inhibition of MV infection by moesin-specific antibodies is dose dependent

To confirm the data obtained with mAb 119, the moesin-specific mAb 38/87 was included in subsequent experiments. The inhibition of MV infection following treatment of cell monolayers with mAbs 119 or 38/87 was determined using the plaque inhibition assay. Half-maximal inhibition of MV infection on Vero cells was achieved after treating cell monolayers with approximately 15 µg/ml of mAb 119 (45% plaque inhibition), while maximal inhibition was observed with  $\geq 50$  µg/ml ( $\geq 90\%$  plaque inhibition) of the mAb (Fig. 5). Comparable results were obtained when the assay was performed on other human cell lines (data not shown). A similar dose-dependent curve, albeit much reduced in comparison to the data for mAb 119, was observed with the moesin-specific antibody mAb 38/87. Half-maximal inhibition of MV infection with this antibody was achieved with 30 µg/ml (27% plaque inhibition) while maximal inhibition required 90 µg/ml (54%

plaque inhibition; Fig. 5). Our data show that mAb 119 and mAb 38/87 recognize different epitopes on moesin, mAb 38/87 recognizing a linear epitope detected in Western blot as described (Lankes *et al.*, 1988), whereas mAb 119 recognizes a conformational epitope.

#### Distribution of the epitopes on MV susceptible cell lines and primary cells

The expression of the epitopes recognized by mAbs 119 and 38/87 on the surface of human primary PBMC and various human and nonhuman cell lines was investigated by flow cytometry. The percentage of cells stained with mAb 119 above the negative control, following labeling with a fluorescein-conjugated second antibody, varied between 40 and 98% for the human cells examined (Fig. 6). The African green monkey kidney cells (Vero), which are routinely used to propagate MV, were positive to approximately 50% (Fig. 6). In comparison, mAb 38/87 exhibited a different staining pattern. For each cell line tested, the level of expression of the mAb 38/87 epitope was much reduced, varying between 14 and 38% (Fig. 6). These results correlate well with the reduced ability of mAb 38/87 to inhibit MV infection in the infection inhibition test (Fig. 5). In contrast to the surface staining of cells, every cell




FIG. 6. Distribution of the epitopes of mAbs 119 and 38/87 on the surface of human and monkey cell lines. Flow cytometry was used to determine the distribution of the epitopes recognized by mAb 119 and mAb 38/87 on the surface of a number of different cell lines. The Y axis shows the percentage of cells stained above background, the open boxes representing values for mAb 119 and the shaded boxes mAb 38/87. Peripheral blood mononuclear cells (PBMC) and U-937 monocytes revealed the highest levels of surface moesin expression detected by mAb 119, whereas U-251 cells (astrocytoma), SKN-MC cells (neuroblastoma), and Vero cells (African green monkey kidney) expressed the epitope for mAb 119 to a lesser extent.

of the cell lines investigated so far strongly expresses moesin intracellularly with equal intensities detected with either mAb 119 or mAb 38/87 (data not shown).

#### Inhibition of MV binding to cells

In order to investigate the question of whether moesin is involved in the attachment of MV to cells, mAbs 38/87 and 119 were tested in rosetting assays for their ability to inhibit MV binding to monkey red blood cells. Incubation of U-937 cells with mAb 38/87 led to  $52 \pm 3\%$  reduction in the number of rosettes whilst mAb 119 gave  $67 \pm 5\%$  reduction in rosette formation when compared to control mAbs W6/32 ( $9 \pm 2\%$ ) and non-binding U85 ( $0 \pm 0\%$ ) ( $n = 3$ ).

#### Inhibition of vaccinia virus MV-H/F-mediated fusion


The MV-H and -F glycoproteins are both required for fusion activity of cellular membranes (Wild *et al.*, 1991). In the rapid screening for new antibodies we found that mAb 119 inhibited fusion of recombinant vaccinia virus MV-H/F-infected U-251, HeLa, and Vero cells (not shown). In addition, mAb 38/87 inhibited fusion of vaccinia virus MV-H/F-infected U-251 cells (Fig. 7), HeLa, and Vero cells (not shown). These data suggest that in addition to the epitope recognized by mAb 119, the less accessible cell surface moesin epitope of mAb 38/87 is also functionally active in inhibition assays of MV fusion.

#### Downregulation of surface moesin expression on MV-infected cells

The degree of surface and intracellular expression of moesin in uninfected and persistently MV-infected cells was determined by flow cytometry. Persistently infected U-937 cells revealed a considerable reduction of the signal intensity for surface moesin (Fig. 8A, compare signal 1 and 2). In comparison, the expression of the MV fusion protein (MV-F) was detected exclusively on infected cells (Fig. 8B, signal 1 represents the negative control and signal 2 that of anti-MV-F mAb A504), and the expression of MHC class I molecules was unaltered following infection (Fig. 8C). The intensity of the intracellular expression of moesin was not affected by the infection with MV (not shown). These data suggest that moesin is downregulated from the surface of infected cells, but unaltered intracellularly.

### DISCUSSION

In this report we have described a mAb (mAb 119) that is directed against an epitope expressed on the surface of a number of human and primate cells and inhibits the binding of MV to its cell receptor, MV glycoprotein-mediated fusion and, hence, infection of the



**FIG. 7.** Inhibition of vaccinia virus MV-H/F-mediated fusion by antibodies to moesin. U-251 cells were infected with a vaccinia virus recombinant expressing MV-H and -F (m.o.i. = 0.1) for 24 hr in the presence of 20  $\mu\text{g/ml}$  of the unrelated antibody G26 to lymphocyte function-associated antigen (LFA-3) as negative control (A), 20  $\mu\text{g/ml}$  mAb 38/87 to moesin (B), and 20  $\mu\text{g/ml}$  mAb K83 to MV-H as positive control (C). Extensive fusion with large syncytia was observed exclusively in cultures treated with the control antibody (A). Bar, 100  $\mu\text{m}$ .

normally susceptible cell. The epitope recognized by mAb 119 is widely expressed to high levels on human and primate cells and the antibody was found to specifically inhibit MV infection. The mAb precipitates a 75-kDa protein, which was identified as the human membrane-organizing extension spike protein, moesin. It is this protein which we believe plays an integral role in the binding and uptake of MV to susceptible cells based on the evidence in this study.




FIG. 8. Downregulation of surface moesin expression after infection of cells with MV. The intensity of the surface expression of moesin on uninfected U-937 cells (signal 1) and U-937 cells persistently infected with MV (signal 2) was measured by flow cytometry. Cells were stained with mAb 38/87 to moesin (A), with mAb A504 recognizing MV-F as a control of infection (B), and with mAb W6/32 to MHC class I as a control for the staining procedure (C). The cell number (Y axis) was plotted against the fluorescence intensity (X axis).

Recently, a glycoprotein with the mass of 57/67 kDa was identified with the aid of a monoclonal antibody as a putative MV receptor (Naniche *et al.*, 1992). This molecule has been identified as the human membrane cofactor protein (CD46) present on many cells of human origin (Naniche *et al.*, 1993b). The described antibody, MC120.6, recognizing CD46, inhibits MV binding in a rosetting assay, reduces infection of HeLa cells by MV, and inhibits cellular fusion mediated by vaccinia virus-

expressed MV-H and -F proteins (Naniche *et al.*, 1992). MV infection also led to the downregulation of CD46 on the cell surface and moreover, it was found that the MV-H protein was responsible for this phenomenon (Naniche *et al.*, 1993a). Since specific receptor downregulation following infection has been observed for several viruses with differing receptor molecules, this finding, in addition to our own observation concerning the downregulation of moesin, add support to the hypothesis that CD46 and moesin act as receptors for MV.

CD46 is known to exist in as many as 14 different isoforms through alternative splicing of the cDNA (Russell *et al.*, 1992) and it is unknown whether or not all these isoforms are able to act as receptor for MV. Indeed, the poliovirus receptor is known to be widely distributed throughout human cell types although the virus is able to infect only a select number of cell types (Ziebert and Wimmer, 1992). It is believed that correct post-transcriptional and post-translational modification of the poliovirus receptor by individual cells is necessary to bind virus (Ziebert and Wimmer, 1992). In the case of MV, cells which do not express human CD46 and are not recognized by CD46-specific antibodies, such as some mouse cell lines, are readily infectable with MV (Rager-Zisman *et al.*, 1984) pointing to the existence of further receptor molecules like moesin, homologs of which are expressed by cells of murine origin.

The protein moesin was initially identified as a cell surface component and thought to act as receptor for binding heparin and heparin sulfate (Lankes *et al.*, 1988). Subsequent cloning of its cDNA and further analysis has revealed that moesin is largely localized at the inner side of the plasma membrane (Lankes *et al.*, 1991). Neither a signal sequence or a transmembrane domain could be identified by sequence analysis, suggesting that it is not incorporated into the plasma membrane in the classical way (Lankes *et al.*, 1991). Following the identification of moesin as the protein precipitated by mAb 119, we acquired a mAb (38/87) specific for the protein moesin that was used to characterize this protein (Lankes *et al.*, 1988, 1991). From the outset it was known that these two mAbs recognized very different epitopes of the protein moesin. By FACS analysis, the epitope of mAb 119 was found to be widely distributed on the surface of human and primate cells, whereas the epitope of mAb 38/87 was detected to a much lower degree on the surface of cells (Fig. 7). Importantly, by Western blot analysis, only mAb 38/87 is able to detect moesin, suggesting that mAb 119 recognizes a conformational epitope. Crucial evidence supporting the hypothesis that moesin is acting as a receptor for MV came from the ability of mAb 38/87 to inhibit MV infection of susceptible cells. Moreover, this mAb inhibited cell fusion following infection with recombinant vaccinia-expressed MV-H and -F protein,



and inhibited rosette formation by MV and by flow cytometric analysis was downregulated on MV-infected cells, similar to that observed with the putative MV receptor CD46 (Naniche *et al.*, 1993a). The above criterion fulfill the basic requirements for receptor blocking activity and mirrors the results obtained with mAb 119. The level of infection inhibition by mAb 38/87 was somewhat less than that observed with mAb 119 and can be correlated to the accessibility of the epitopes for each specific antibody on the surface of cells. As mentioned earlier, moesin is mainly localized on the internal side of the cell membrane which becomes apparent when cells are permeabilized prior to staining with antibodies. It is possible that the mode by which mAb 38/87 inhibits MV binding is by blocking MV access to the conformational mAb 119 site following attachment to its cell surface-expressed motif. This is currently being investigated.

A further important observation that points to moesin as a putative receptor for MV is based on the observation that the purified protein binds MV itself. This is not unusual given that, for example, soluble CD4 has been shown to inhibit the infection of certain cell types with HIV (Orloff *et al.*, 1993). Interestingly, in the case of HIV, alternative receptors which are not inhibitable by soluble CD4 can be used in the brain and by muscle cells (Clapham *et al.*, 1989). In addition, the results presented in this study point to the fact that more than one molecule is likely to be acting as receptors for MV. To gain entry into cells, MV utilizes two major protein species, the H and F proteins, and both are required for fusion of cells (Wild *et al.*, 1991). It may be that the initial receptor interaction is mediated by a single specific receptor species with MV-H, followed by contact of the virus with a second receptor (F interaction) to initiate successful infection. Our data show that there is a specific interaction between MV and moesin and that the critical motif is conformational and expressed on the cell surface. The question of whether moesin and CD46 are alternative receptors or part of one receptor complex is currently under investigation.

Moesin, as well as its homologs of the ERM (ezrin-radixin-moesin) gene family have been shown to be localized at cell-to-cell adherens junctions, cleavage furrows, microvilli, and ruffling membranes (Sato *et al.*, 1992). Most intense staining was found at the cleavage furrows in the mitotic phase of the cell cycle and could point to a correlation with the cell cycle dependence of infection with MV. Furthermore, the concentration of ERM proteins at cell-to-cell adhesion junctions and cleavage furrows and their interaction with actin (Sato *et al.*, 1992), mediating the association of actin filaments with the plasma membrane, suggest a direct connection between actin filaments and MV. How moesin is involved in the attachment and uptake

of MV is an intriguing question which is under investigation.

## ACKNOWLEDGMENTS

We thank Sibylle Schneider-Schaulies for helpful discussions and Katerina Pech for technical assistance. This work was supported by grants from the Deutsche Forschungsgemeinschaft and the Bundesministerium für Forschung und Technologie. L. M. Dunster is a research fellow of the Alexander von Humboldt Foundation.

## REFERENCES

- CASALI, P., NAKAMURA, M., and MCCHESENEY, M. B. (1989). Immunosuppression by measles virus. In "Virus Induced Immunosuppression" (S. Spector, M. Bendinelli, and H. Friedmann, Eds.), Chap. 18, pp. 345-373. Plenum Press, NY/London.
- CLAPHAM, P. R., WEBER, J. N., WHITBY, D., MCINTOSH, K., DALGEISH, A. G., MADDON, P. J., DEEN, K. C., SWEET, R. W., and WEISS, R. A. (1989). Soluble CD4 blocks the infectivity of diverse strains of HIV and SIV for T cells and monocytes but not for brain and muscle cells. *Nature* 337, 368-370.
- DVESKLER, G. S., DIEFFENBACH, C. W., CARDELLICHO, C. B., MCCUAIG, K., PENSIERO, M. N., JIANG, G-S., BEAUCHEMIN, N., and HOMES, K. V. (1993). Several members of the mouse carcinoembryonic antigen-related glycoprotein family are functional receptors for the coronavirus mouse hepatitis virus A59. *J. Virol.* 67, 1-8.
- ECKERSKORN, C., and LOTTSPREICH, F. (1989). Internal amino acid sequence analysis of proteins separated by gel electrophoresis after tryptic digestion in a polyacrylamide matrix. *Chromatographia* 29, 92-94.
- ECKERSKORN, C., MEWES, W., GORETZKI, H., and LOTTSPREICH, F. (1988). A new siliconized-glass fiber as a support for protein-chemical analysis of electroblotted proteins. *Eur. J. Biochem.* 176, 509-519.
- ESIRI, M. M., OPPENHEIMER, D. R., BROWNELL, B., and HAIRE, M. (1981). Distribution of measles antigen and immunoglobulin containing cells in the CNS in subacute sclerosing panencephalitis (SSPE) and atypical measles. *J. Neurol. Sci.* 53, 29-43.
- HAASE, A. T., GANTZ, D., EBLE, B., WALKER, D., STOWING, L., VENTURA, P., BLUM, H., WIETGREFE, S., ZUPANICIC, M., and TOURTELLOTTE, W. (1985). Natural history of restricted synthesis and expression of measles virus genes in subacute sclerosing panencephalitis. *Proc. Natl. Acad. Sci. USA* 82, 3020-3024.
- HAROUSE, J. M., BHAT, S., SPITALNIK, S. L., LAUGHLIN, M., STEFANO, K., SILBERBERG, D. H., and GONZALES-SCARANO, F. (1991). Inhibition of entry of HIV-1 in neural cell lines by antibodies against galactosyl ceramide. *Science* 253, 320-323.
- HAROUSE, J. M., KUNSCH, C., HARTLE, H. T., LAUGHLIN, M. A., HOXIE, J. A., WIGDAHL, B., and GONZALES-SCARANO, F. (1989). CD4-independent infection of human neural cells by human immunodeficiency virus type 1. *J. Virol.* 63, 2527-2533.
- HARROWE, G., MITSUHASHI, M., and PAYAN, D. G. (1990). Measles virus-substance P receptor interactions. *J. Clin. Invest.* 85, 1324-1327.
- HARROWE, G., SUDDUTH-KLINGER, J., and PAYAN, D. G. (1992). Measles virus-substance P receptor interaction: Jurkat lymphocytes transfected with substance P receptor cDNA enhance measles virus fusion and replication. *Cell. Mol. Neurobiol.* 12, 397-409.
- KIRK, J., ZHOU, A. L., McQUAID, S., COSBY, S. L., and ALLEN, I. V. (1991). Cerebral endothelial cell infection by measles virus in subacute sclerosing panencephalitis: Ultrastructural and *in situ* hybridization evidence. *Neuropathol. Appl. Neurobiol.* 17, 289-297.
- KRAH, D. L. (1989). Characterization of octyl-glucoside-solubilized cell membrane receptors for binding measles virus. *Virology* 172, 386-390.

- KRAH, D. L. (1991). Receptors for binding measles virus on host cells and erythrocytes. *Microbial Pathogen*. **11**, 221–228.
- LANKES, W. T., and FURTHMAYER, H. (1991). Moesin: A member of the protein 4.1-talin-ezrin family of proteins. *Proc. Natl. Acad. Sci. USA* **88**, 8297–8301.
- LANKES, W. T., GRIESMACHER, A., GRÜNWARD, J., SCHWARTZ-ALBIEZ, R., and KELLER, R. (1988). A heparin-binding protein involved in inhibition of smooth muscle cell proliferation. *Biochem. J.* **251**, 831–842.
- LENTZ, T. L. (1990). The recognition event between virus and host receptor: A target for antiviral agents. *J. Gen. Virol.* **71**, 751–766.
- MOENCH, T. R., GRIFFIN, D. E., OBRICHT, C. R., VAISBERG, A., and JOHNSON, R. T. (1988). Distribution of measles virus antigen and RNA in acute measles with and without neurologic involvement. *J. Infect. Dis.* **158**, 433–442.
- NANICHE, D., WILD, T. F., RABOURDIN-COMBE, C., and GERLIER, D. (1992). A monoclonal antibody recognizes a human cell surface glycoprotein involved in measles virus binding. *J. Gen. Virol.* **73**, 2617–2624.
- NANICHE, D., WILD, T. F., RABOURDIN-COMBE, C., and GERLIER, D. (1993a). Measles virus haemagglutinin induces down-regulation of gp57/67, a molecule involved in virus binding. *J. Gen. Virol.* **74**, 1073–1079.
- NANICHE, D., VARIOR-KISHNAN, G., CERVONI, F., WILD, T. F., ROSSI, B., RABOURDIN-COMBE, C., and GERLIER, D. (1993b). Human membrane cofactor protein (CD 46) acts as a cellular receptor for measles virus. *J. Virol.*, **67**, 6025–6032.
- ORLOFF, S. L., KENNEDY, M. S., BELPERRON, A. A., MADDON, P. J., and McDUGAL, J. S. (1993). Two mechanisms of soluble CD4 (sCD4)-mediated inhibition of human immunodeficiency virus type 1 (HIV-1) infectivity and their relation to primary HIV-1 isolates with reduced sensitivity to sCD4. *J. Virol.* **67**, 1461–1471.
- RAGER-ZISMAN, B., EGAN, J. E., KRESS, Y., and BLOOM, B. R. (1984). Isolation of cold-sensitive mutants of measles virus from persistently infected murine neuroblastoma cells. *J. Virol.* **51**, 845–855.
- RUSSELL, S. M., SPARROW, R. L., MCKENZIE, I. F. C., and PURCELL, D. F. J. (1992). Tissue specific and allelic expression of the complement regulator CD46 is controlled by an alternative splicing. *Eur. J. Immunol.* **22**, 1513–1518.
- SATO, N., FUNAYAMA, N., NAGAFUCHI, A., YONEMURA, S., TSUKITA, S., and TSUKITA, S. (1992). A gene family consisting of ezrin, radixin and moesin. *J. Cell Sci.* **103**, 131–143.
- SCHNEIDER-SCHAULIES, J., SCHNEIDER-SCHAULIES, S., BRINKMANN, R., TAS, P., HALBRÜGGE, M., WALTER, U., HOLMES, C., and TER MEULEN, V. (1992). HIV-1 gp120 receptor on CD4-negative brain cells activates a tyrosine kinase. *Virology* **191**, 765–772.
- SCHNEIDER-SCHAULIES, J., SCHNEIDER-SCHAULIES, S., and TER MEULEN, V. (1993). Differential induction of cytokines by primary and persistent measles infections in human glial cells. *Virology* **195**, 219–228.
- SCHNEIDER-SCHAULIES, S., and TER MEULEN, V. (1992). Molecular aspects of measles virus induced central nervous system diseases. In "Molecular Neurovirology" (R. P. Roos, Ed.), pp. 419–449. Humana Press Inc., Clifton, NJ.
- WHITE, J. M., and LITTMAN, D. R. (1989). Viral receptors of the immunoglobulin superfamily. *Cell* **56**, 725–728.
- WILD, T. F., MALVOISIN, E., and BUCKLAND, R. (1991). Measles virus: Both the haemagglutinin and fusion glycoproteins are required for fusion. *J. Gen. Virol.* **72**, 439–442.
- YOSHIKAWA, Y., YAMANOUCHI, K., TAKASU, T., RAUF, S., and AHMED, A. (1991). Structural homology between hemagglutinin (HA) of measles virus and the active site of long neurotoxins. *Virus Genes* **5**, 57–67.
- ZIEBERT, A., and WIMMER, E. (1992). N-glycosylation of the virus binding domain is not essential for function of the human poliovirus receptor. *J. Virol.* **66**, 7368–7373.