

IX. Literaturverzeichnis

- Aichholz R. und Lorbeer E. (1999) Investigation of combwax of honeybees with high-temperature gas chromatography-chemical ionization mass spectrometry. I. High-temperature gas chromatography. *Journal of Chromatography A* 855: 601-615.
- Aitchison J. (1986) *The Statistical Analysis of Compositional Data*. Chapman and Hall, London New York.
- Arnold G., Queunet B., Cornuet J.-M., Masson C., de Schepper B., Estoup A. und Gasqui P. (1996) Kin recognition in honeybees. *Nature* 370: 498.
- Auger J., Rousset S., Thibout E. und Jaillas B. (1998) Solid-phase microextraction-gas chromatography-direct deposition infrared spectrometry as a convenient method for the determination of volatile compounds from living organisms. *Journal of Chromatography A* 819: 45-50.
- Basson I. und Reynhardt E.C. (1992) The structure and melting of paraffinic Fischer-Tropsch waxes. *Chemical Physics Letters* 198 (3,4): 367-372.
- Basson I. und Reynhardt E.C. (1988) An Investigation of the Structures and Molecular Dynamics of Natural Waxes: I. Beeswax. *Journal of Physiology* 21: 1421-1428.
- Bitterman M.E., Menzel R., Fietz A. und Schäfer S. (1983) Classical conditioning of proboscis extension in honeybees (*Apis mellifera*). *J. Comp. Psychol.* 97 (2): 107-119.
- Blomquist G.J., Tillman J.A., Mpuru S. und Seybold S.J. (1998) The cuticle and cuticular hydrocarbons of insects: Structure, Function and Biochemistry. *in: Pheromone communication in social insects: Ants, Wasps, Bees and Termites*. eds: Vander Meer R.K., Breed M.D., Espelie K.E. und Winston M.L., Westview Press, Boulder.
- Blomquist G.J., Nelson D.R. und de Renobales M. (1987). Chemistry, Biochemistry and Physiology of Insect Cuticular Lipids. *Archives of Insect Biochemistry and Physiology* 6: 227-265.
- Blomquist G.J., Chu A.J. und Remaley S. (1980) Biosynthesis of Wax in the Honeybee *Apis mellifera* L. *Insect Biochem.* 10: 313-321.
- Blomquist G.J. und Ries M.K. (1979) The Enzymatic Synthesis of Wax Esters by a Microsomal Preparation from the Honeybee *Apis mellifera* L. *Insect Biochem.* 9: 183-188.
- Blum M.S. und Fales H.M. (1988) Eclectic chemisociality of the honeybee: A wealth of behaviors, pheromones and exocrine glands. *J. Chem. Ecol.* 14 (11): 2099-2107.
- Blume A. (1996) Properties of lipid vesicles: FTIR spectroscopy and fluorescence probe studies. *Curr. Opin. Colloid. Interface S.* 1(1): 64-77.

- Breed M.D., Leger E.A., Pearce A.N. und Wang Y.J. (1998) Comb wax effects on the ontogeny of honey bee nestmate recognition. *Animal Behaviour* 55: 13-20.
- Breed M.D., Garry M.F., Pearce A.N., Hibbard B.E., Bjostad L.B. und Page jr. R.E. (1995a) The role of wax comb in honey bee nestmate recognition. *Animal Behaviour* 50 (2): 489-496
- Breed M.D., Page jr. R.E., Hibbard B.E. und Bjostad L.B. (1995b) Interfamily variation in comb wax hydrocarbons produced by honey bees. *J. Chem. Ecol.* 21 (9): 1329-1338.
- Breed M.D. und Stiller T.M. (1992) Honey Bee, *Apis mellifera*, Nestmate Discrimination: Hydrocarbon Effects and the Evolutionary Implications of Comb Choice. *Animal Behaviour* 43: 875-883.
- Brewster D. (1815) Experiments on the Depolarisation of Light: On Affection of Light Transmitted Through Crystallized Bodies. *Philos. Trans. R. Soc. Lond.* 104: 187-211.
- Brohmer P. (1988) Fauna von Deutschland. 17. Auflage Quelle und Meyer Verlag, Heidelberg Wiesbaden.
- Buckner J.S. (1993) Cuticular Polar Lipids of Insects. *in: Insect Lipids: Chemistry, Biochemistry and Biology.* eds: Stanley-Samuelson D.W. und Nelson D.R., University of Nebraska Press, Lincoln.
- Butler C.G., Callow R.K., Greenway A.R. und Simpson J. (1974) Movement of the pheromone, 9-oxodec-2-enoic acid, applied to the body surfaces of honeybees (*Apis mellifera*). *Ent. exp. appl.* 17: 112-116.
- Butler C.G., Fletcher D.J.C. und Watler D. (1969) Nest-Entrance Marking with Pheromones by the Honeybee *Apis mellifera* L., and by a Wasp, *Vespula vulgaris* L. *Animal Behaviour* 17: 142-147.
- Butler C.G., Callow R.K., Johnston F.R.S. und Johnston N.C. (1961) The isolation and synthesis of queen substance, 9-oxodec-trans-2-enoic acid, a honeybee pheromone. *Proc. Roy. Soc. B* 155: 417-432.
- Cameron D.G., Casal H.L. und Mantsch H.H. (1980) Characterization of the pretransition in 1,2-dipalmitoyl-sn-glycero-3-phosphocholine by Fourier Transform Infrared Spectroscopy. *Biochemistry* 19: 3665-3672.
- Campoy I. (1996) FTIR studies of thermotropic main chain polyesters. Part1. Poly(decamethylene-4,4'-diphenoxy terephthalate). *Spectrochimica Acta* 52: 619-627.
- Carlson D.A., Roan C-S., Yost R.A. und Hector J. (1989) Dimethyl disulfide derivatives of long chain alkenes, alkadienes and alkatrienes for gas chromatography-mass spectrometry. *Analyt. Chem.* 61: 1564-1571.
- Chauvin R. (1962) Sur l'épagine et sur les glandes tarsales d'Arnhart. *Insectes sociaux* 9 (1):

- Darchen R. (1962) Observation directe du développement d'un rayon de cire. Le role des chaines d'abeilles. *Insectes sociaux* 9: 103-120.
- Davidson B.C. und Hepburn H.R. (1986) Transformation of the Acylglycerols in Comb Construction by Honeybees. *Naturwissenschaften* 73: 159-160.
- Dönhoff E. (1855) Über den gelben Farbstoff des Wachses. *Bienenzeitung (Nördlingen)* 11: 179.
- Downs S.G. und Ratnieks F.L. (1999) Recognition of conspecifics by honeybee guards uses nonheritable cues aquired in the adult stage. *Animal Behaviour* 58: 643-648.
- Esch H. (1961) Über die Schallerzeugung beim Werbetanz der Honigbiene. *Z. Vergl. Physiol.* 45: 1-11.
- Esslen J. und Kaissling K-E. (1976) Zahl und Verteilung antennaler Sensillen bei der Honigbiene (*Apis mellifera* L.). *Zoomorphologie* 83: 227-251.
- Ferguson A.W. und Free J.B. (1980) Queen pheromone transfer within honeybee colonies. *Physiol. Entomol.* 5: 359-366.
- Francis B.R., Blanton W.E., Littlefield J.L. und Nunamaker R.A. (1989) Hydrocarbons of the Cuticle and Hemolymph of the Adult Honey Bee (Hymenoptera: Apidae). *An. Entomol. Soc. Am.* 82 (4): 486-494.
- Francis B.R., Blanton W.E. und Nunamaker R.A. (1985) Extractable surface hydrocarbons of workers and drones of the genus *Apis*. *Journal of Apicultural Research* 24 (1): 13-16.
- Francis G.W. und Veland K. (1981) Alkylthiolation for the determination of double-bond positions in linear alkenes. *Journal of Chromatography* 219: 379-384.
- Free J.B. (1987) Pheromones of Social Bees. Chapman and Hall, London.
- Free J.B. (1983) The social organisation of honey bees. Arc & Throstle Press, Todmorden.
- Free J.B. und Williams I.H. (1974) Factors determining food storage and brood rearing in honeybee (*Apis mellifera* L.) comb. *J. Entomol.* A 49: 47-63.
- Freudenstein H. (1961) Beobachtungen über die Färbung der Waben durch die Honigbiene. *Zoologischer Beitrag* 7: 311-319.
- Fröhlich B., Tautz J. und Riederer M. (2000a) Chemometric classification of comb and cuticular waxes of the honeybee *Apis mellifera carnica*. *J. Chem. Ecol.* 26 (1): 123-137.
- Fröhlich B., Riederer M. und Tautz J. (2000b) Comb wax discrimination by honeybees tested with the proboscis extension reflex. *J. exp. Biol.* (im Druck).
- Gamboa, G.J., Reeve, H.K. und Pfennig D.W. (1986). The evolution and ontogeny of nestmate recognition in social wasps. *Ann .Rev. Entomol.* 31: 431-454.

- Getz W.M. und Page R.E.jr. (1991) Chemosensory kin-communication systems and kin recognition in honey bees. *Ethology* 87: 298-315.
- Getz W.M., Brückner D. und Smith K.B. (1988) Variability of chemosensory stimuli within honeybee (*Apis mellifera*) colonies: Differential conditioning assay for discrimination cues. *Journal Chem. Ecol.* 14 (1): 253-264.
- Getz W.M. und Smith K.B. (1986) Honey bee kin recognition: learning self and nestmate phenotypes. *Animal Behaviour* 34: 1617-1626.
- Getz W.M., Brückner D. und Smith K.B. (1986) Conditioning honeybees to discriminate between heritable odors from full and half sisters. *J. Comp. Physiol. A* 159: 251-256.
- Ghisalberti E.L. (1979) Propolis: A Review. *Bee World* 60: 59-84.
- Gibbs A. (1995) Physical properties of insect cuticular hydrocarbons: model mixtures and lipid interactions. *Comp. Biochem. Physiol.* 112B: 667-672.
- Gibbs A. und Pomonis J.G. (1995) Physical properties of insect cuticular hydrocarbons: The effects of chain length, methyl-branching and unsaturation. *Comp. Biochem. Physiol.* 112B: 243-249.
- Gibbs A., Kuenzli M. und Blomquist G.J. (1995) Sex- and age-related changes in the biophysical properties of cuticular lipids of the housefly, *Musca domestica*. *Archives of Insect Biochemistry and Physiology* 29: 87-97.
- Gibbs A. und Crowe J.H. (1991) Intra-individual variation in cuticular lipids studied using fourier transform infrared spectroscopy. *J. Insect Physiol.* 37(10): 743-748.
- Gibbs A., Mousseau T.A. und Crowe J.H. (1991) Genetic and acclimatory variation in biophysical properties of insect cuticular lipids. *Proc. Natl. Acad. Sci. USA* 88: 7257-7260.
- Hepburn H.R. (1998) Reciprocal interactions between honeybees and combs in the integration of some colony functions in *Apis mellifera* L. *Apidologie* 29: 47-66.
- Hepburn H.R., Bernard R.T.F., Davidson B.C., Muller W.J., Lloyd P., Kurstjens S.P. und Vincent S.L. (1991) Synthesis and Secretion of Beeswax in Honeybees. *Apidologie* 22: 21-36.
- Hepburn H.R. und Kurstjens S.P. (1988) The combs of honeybees as composite materials. *Apidologie* 19 (1): 25-36.
- Hepburn H.R. und Muller W.J. (1988) Wax Secretion in Honeybees. Thermal Integrity of the Festoon. *Naturwissenschaften* 75: 628-629.
- Hepburn H.R. (1986) Honeybees And Wax. An Experimental Natural History. Springer Verlag, Berlin Heidelberg New York.
- Hepburn H.R. und Kurstjens S.P. (1984) On the Strength of Propolis (Bee Glue).

- Hepburn H.R., Armstrong E. und Kurstjens S. (1983) The Ductility of Native Beeswax is Optimally Related to Honeybee Colony Temperature. *South African Journal of Science* 79: 416-417.
- Howard R. W. (1993) Cuticular Hydrocarbons and Chemical Communication. *in: Insect Lipids: Chemistry, Biochemistry and Biology.* eds: Stanley-Samuels D.W. und Nelson D.R., University of Nebraska Press, Lincoln.
- Howard R. W., Blomquist G.J. (1982) Chemical Ecology and Biochemistry of Insect Hydrocarbons. *Ann. Rev. Entomol.* 27: 149-172.
- Huber F. (1814) Nouvelles observations sur les abailles. 2. Übersetzung (1926) Dadant, Hamilton.
- Hüsing J.O. und Nitschmann J. (1987) Lexikon der Bienenkunde. Ehrenwirth-Verlag, München.
- Joerges J., Küttner A., Galizia C.G. und Menzel R. (1997) Representation of odours and odour mixtures visualized in the honeybee brain. *Nature* 387: 285-288.
- Jordan R., Tischer J., und Illner E. (1940) Vergleich des von Altbienen erzeugten Wachses mit "Jungfernwachs" und gewöhnlichem Bienenwachs. *Z. Vergl. Physiol.* 28: 353-357.
- Iyengar B.T.R und Schlenk H. (1968) Melting points of synthetic wax esters. *Lipids* 4 (1): 28-30.
- Jyothi J.V.A. und Reddy C.C. (1993) Physical and chemical properties of the comb of the giant honeybee, *Apis dorsata* F. *Geobios* 20: 269-271.
- Kirchner W.H. (1993) Vibrational Signals in the Tremble Dance of the Honeybee, *Apis mellifera*. *Behavioral Ecology and Sociobiology* 33: 169-172.
- Kirsch T. (1997) Charakterisierung der kutikulären Barriereigenschaften ausgewählter Baumarten. Dissertation Universität Würzburg.
- Koeniger N. und Veith H.J. (1984) Spezifität eines Brutpheromons und Bruterkennung bei der Honigbiene (*Apis mellifera* L.). *Apidologie* 15 (2): 205-210.
- Koidsumi K. (1957) Antifungal Action of Cuticular Lipids in Insects. *J. Insect. Physiol.* 1: 40-51.
- Kollatukudy P.E. (1967) Mechanisms of synthesis of waxy esters in brocolli (*Brassica oleracea*). *Biochem.* 6: 2705-2715.
- Kratky O. (1937) Die Orientierung der Moleküle des Bienenwachses und die Festigkeit der Waben. *Naturwissenschaften* 25: 239.
- Kurstjens S.P., Hepburn H.R., Schoening F.R.L. und Davidson B.C. (1985) The Conversion of Wax Scales into Comb Wax by African Honeybees. *J. Comp. Physiol. B* 156: 95-

- Lachenbruch P.A. und Clarke W.R. (1980) Discriminant Analysis and its Application in Epidemiology. *Methods of Information in Medicine* 19: 220-226.
- Lachenbruch P.A. und Goldstein M. (1979) Discriminant analysis. *Biometrics* 35: 69-85.
- Lahav S., Soroker V., Hefetz A. und Vander Meer R.K. (1999) Direct behavioral evidence for hydrocarbons as ant recognition discriminators. *Naturwissenschaften* 86: 246-249.
- Lambremont E.N. und Wykle R.L. (1979) Wax Synthesis by an Enzyme System from the Honeybee. *Comp. Biochem. Physiol.* 63 b: 131-135.
- Larsson K. (1986) Physical properties- structural and physical characteristics. *in: The Lipid Handbook*. eds: Gunstone F.D., Harwood J.L. und Padley F.B., Chapman and Hall, London New York.
- Le Conte Y., Sreng L. und Poitout S.H. (1995) Brood pheromone can modulate the feeding behavior of *Apis mellifera* workers (Hymenoptera: Apidae). *J. Econ. Entomol.* 88 (4): 799-804.
- Le Conte Y., Sreng L. und Trouiller J. (1994) The Recognition of Larvae by Worker Honeybees. *Naturwissenschaften* 81: 462-465.
- Le Conte Y., Arnold G., Trouiller J., Masson C. und Chappe B. (1990) Identification of a Brood Pheromone in Honeybees. *Naturwissenschaften* 77: 334-336.
- Lensky Y. und Cassier P. (1992) Control of Swarming by Queen Bee Pheromones. *Bee Science* 2 (1): 7-11.
- Lensky Y., Cassier P., Rosa S. und Grandperrin D. (1991) Induction of balling in worker honeybees (*Apis mellifera* L.) by "stress" pheromone from Koschewnikow glands of queen bees: Behavioural, structural and chemical study. *Comp. Biochem. Physiol.* 100A (3): 585-594.
- Lensky Y., Slabetzki Y. (1981) The Inhibiting Effect of the Queen Bee (*Apis mellifera* L.) Foot-Print Pheromone on the Construction of Swarming Queen Cups. *J. Insect Physiol.* 27 (5): 313-323.
- LeRoux J.H. (1970) Fischer-Tropsch waxes. V. Study of branching and its effects on crystallinity using an improved infra-red method. *J. appl. Chem.* 19 (230): 203-207.
- Le Roux J.H. (1969) Fischer –Tropsch waxes. II. Crystallinity and physical properties. *J. appl. Chem.* 19 (March): 86-88.
- Lewis C.T. (1962) Diffusion of oil films over insects. *Nature* 193: 904.
- Lockey K.H. (1988) Lipids of the Insect Cuticle: Origin, Composition and Function. *Comp. Biochem. Physiol.* 89 b (4): 595-645.
- Lockey K.H. (1980) Insect Cuticular Hydrocarbons. *Comp. Biochem. Physiol.* 65 b: 457-462.

- Lorenzi M.C., Begnères A.G., Clément J.-L. und Turillazzi S. (1997) *Polistes biglumis bimaculatus* epicuticular hydrocarbons and nestmate recognition. *Insectes Sociaux* 44: 123-138.
- Lourens J.A.J. und Reynhardt E.C. (1979) NMR investigation in Fischer-Tropsch waxes. *J. Phys. D: Appl. Phys.* 12: 1963-1972.
- Maier I. und Fieber M. (1988) Retention Characteristics of Volatile Compounds on Tenax TA. *J. High Resol. Chrom. Commun.* 11: 566-576.
- Mantsch H.H. und McElhaney R.N. (1991) Phospholipid phase transitions in model and biological membranes as studied by infrared spectroscopy. *Chemistry and Physics of Lipids* 57: 213-226.
- Marcucci M.C. (1995) Propolis: Chemical Composition, Biological Properties and Therapeutic Activity. *Apidologie* 26: 83-99.
- Markstädter C. (1994) Untersuchungen zur jahreszeitlichen Entwicklung der kutikulären Wachse von *Fagus sylvatica* L. Dissertation Universität Kaiserslautern.
- Marsen H. (1962) Infrarotspektroskopische Untersuchungen zur Temperaturabhängigkeit der Kristallinität von Esterwachsen. *Fette, Seifen, Anstrichmittel* 64 (1): 41-44.
- Marsen H. (1961) Infrarotspektroskopische Untersuchungen zur Temperaturabhängigkeit der Kristallinität von Kohlenwasserstoff-Wachsen. *Fette, Seifen, Anstrichmittel* 63 (1): 42-48.
- Martin H. und Lindauer M. (1966) Sinnesphysiologische Leistungen beim Wabenbau der Honigbiene. *Z. Vergl. Physiol.* 53: 372-404.
- McDaniel C.A., Howard R.W., Blomquist G.J. und Collins A.M. (1984) Hydrocarbons of the Cuticle, Sting Apparatus and Sting Shaft of *Apis mellifera* L. Identification and Preliminary Evaluation as Chemotaxonomic Characters. *Sociobiology* 8 (3): 287-298.
- Menzel R., Greggers U. und Hammer M. (1993) Functional Organization of Appetitive Learning and Memory in a Generalist Pollinator, the Honey Bee. *in: Insect Learning: Ecology and Evolutionary Perspectives.* eds: Papaj D.R. und Lewis A.C., Chapman and Hall, New York London.
- Merk S. (1998) FTIR-spektroskopische Untersuchungen zum Phasenverhalten und zur Kristallinität pflanzlicher kutikulärer Wachse. Dissertation Universität Würzburg.
- Merk S., Riederer M. und Blume A. (1998) An FTIR spectroscopic method for the determination of physical parameters for plant cuticular waxes. *Planta* 204: 44-53.
- Meyer W. (1954) Die "Kittharzbiene" und ihre Tätigkeiten. *Zeitschrift für Bienenforschung* 2 2(7): 185-200.

- Michelsen A., Kirchner W.H. und Lindauer M. (1986) Sound and Vibrational Signals in the Dance Language of the Honeybee, *Apis mellifera*. *Behav. Ecol. Sociobiol.* 18: 207-212.
- Morabito P.L., Ormand J.R. und Nestrick T.J. (1990) Dynamic vapor generator for moderate and low volatility compounds. *Chemosphere* 21: 991-997.
- Moritz R.F.A. und Southwick E.E. (1992) Bees as Superorganisms. An Evolutionary Reality. Springer Verlag, Berlin Heidelberg New York.
- Moritz R.F.A. und Crewe R.M. (1991) The volatile emission of honeybee queens (*Apis mellifera* L.). *Apidologie* 22: 205-212.
- Müssbichler A. (1952) Die Bedeutung äußerer Einflüsse und der Corpora allata bei der Afterweiselentstehung von *Apis mellifera*. *Z. Vergl. Physiol.* 34: 207-221.
- Nachtigall W. und Kresling B. (1992) Bauformen der Natur: Teil II.: Technische Biologie und Bionik von Platten- und Faltkonstruktionen. *Naturwissenschaften* 79: 251-259.
- Nass R., Markstädter C., Hauke V. und Riederer, M. (1998) Quantitative gas chromatographic analysis of plant cuticular waxes containing long-chain aldehydes. *Phytochem. Analysis* 9: 112-118.
- Naumann K., Winston M.L., Slessor K.N., Prestwich G.D. und Latli B. (1992) Intra-nest transmission of aromatic honey bee queen mandibular gland pheromone components: movement as a unit. *Can. Entomol.* 124: 917-934.
- Naumann K. (1991) Grooming behaviors and the translocation of queen mandibular gland pheromone on worker honey bees (*Apis mellifera* L.). *Apidologie* 22: 523-531.
- Naumann K., Winston M.L., Slessor K.N., Prestwich G.D. und Webster F.X. (1991) Production and transmission of honey bee queen (*Apis mellifera* L.) mandibular gland pheromone. *Behav. Ecol. Sociobiol.* 29: 321-332.
- Nelson D.R. und Blomquist G.J. (1995) Insect Waxes. *in: Waxes: Chemistry, Molecular Biology and Functions*. Volume 6. ed: Hamilton R. J., The Oily Press, Dundee.
- Omholt S.W. (1988) Drone production in honeybee colonies: Controlled by a longlasting pheromone from the drones? *J. theor. Biol.* 134: 309-318.
- Page jr. R.E., Metcalf R.A., Metcalf R.L., Erickson jr. E.H. und Lampman R.L. (1991) Extractable Hydrocarbons and Kin Recognition in Honeybee (*Apis mellifera* L.). *J. Chem. Ecol.* 17 (4): 745-756.
- Page R.E., Blum M.S. und Fales H.M. (1988) o-Aminoacetophenone, a Pheromone that Repels Honeybees (*Apis mellifera* L.). *Experientia* 44: 270-271.
- Pain J., Barbier M., Bogdanovsky D. und Lederer E. (1962) Chemistry and biological activity of the secretions of queen and worker honeybees (*Apis mellifica* L.). *Comp. Biochem. Physiol.* 6: 233-241.

- Pankiw.T., Page jr. R.E., und Fondrk M.K. (1998) Brood pheromone stimulates pollen foraging in honey bees (*Apis mellifera*). *Behav. Ecol. Sociobiol.* 44: 193-198.
- Pankiw T., Winston M.L., Plettner E., Slessor K.N., Pettis J.S. und Taylor O.R. (1996) Mandibular gland components of european and africanized honey bee queens (*Apis mellifera* L.). *J. Chem. Ecol.* 22 (4): 605-615.
- Pawliszyn J. (1997) Solid Phase Microextraction: Theory and Practice. Wiley-VCH, New York Chichester Weinheim Brisbane Singapore Toronto.
- Petersen H. (1997) Pheromone im Bienenvolk. *Die Neue Bienenzucht - Norddeutsche Imkerzeitung* 5: 135-136.
- Philipp P.W. (1928) Das Kittharz, seine Herkunft und Verwendung im Bienenhaushalt. *Biologisches Zentralblatt* 48: 705-715.
- Piek T. (1964) Synthesis of Wax in the Honeybee (*Apis mellifera* L.). *J. Insect Physiol.* 10: 563-572.
- Plettner E., Otis G.W., Wimalaratne P.D.C., Winston M.L., Slessor K.N., Pankiw T. und Punchihewa P.W.K. (1997) Species- and caste-determined mandibular gland signals in honeybees (*Apis*). *J. Chem. Ecol.* 23 (2): 363-377.
- Plettner E., Slessor K. N., Winston M. L. und Oliver J. E. (1996) Caste-Selective Pheromone Biosynthesis in Honeybees. *Science* 271: 1851-1853.
- Post D.C., Page jr R.E. und Erickson E.H. (1987) Honeybee (*Apis mellifera* L.) queen feces: Source of a pheromone that repels worker bees. *J. Chem. Ecol.* 13 (3): 583-591.
- Reid R.C., Prausnitz J.M., und Sherwood T.K. (1988) The properties of gases and liquids. 4. edition, McGraw-Hill Book Company, New York.
- Reyment R.A. (1989) Compositional data analysis. *Terra Review* 1: 29-34.
- Reynhardt E.C. und Riederer M. (1994) Structures and molecular dynamics of plant waxes II: Cuticular waxes from leaves of *Fagus sylvatica* L. and *Hordeum vulgare* L. *European Biophysics Journal* 23: 59-70.
- Riederer M. und Markstädter C. (1996) Cuticular waxes: a critical assessment of current knowledge. in: Plant Cuticles: an Integrated Functional Approach. ed: Kerstiens G., BIOS Scientific Publishers, Oxford.
- Rinderer T.E. (1981) Volatiles from empty comb increase hoarding by the honey bee. *Animal Behaviour* 29: 1275-1276.
- Robinson G.E. (1996) Chemical Communication in Honeybees. *Science* 271: 1824-1825.
- Rohrseitz K. (1998) Biophysikalische und ethologische Aspekte der Tanzkommunikation der Honigbienen (*Apis mellifera carnica* Pollm.). Dissertation Universität Würzburg.

- Sandeman D.C., Tautz J. und Lindauer M. (1996) Transmission of Vibration across Honeycombs and its Detection by Bee Leg Receptors. *J. Exp. Biol.* 199: 2585-2594.
- Schoene K., Steinhanses J. und König A. (1990) Sorption isotherms of organic vapours on Tenax. *Journal of Chromatography* 514: 279-286.
- Schoening F.R.L. (1980) The X-Ray Diffraction Pattern and Deformation Texture of Beeswax. *S. A. Journal of Science* 76: 262-265.
- Schreiber L und Schönherr J. (1993) Mobilities of organic compounds in reconstituted cuticular wax of barley leaves: Determination of diffusion coefficients. *Pestic. Sci.* 38: 353-361.
- Schwedt G. (1996) Taschenatlas der Analytik. 2. Auflage, Georg Thieme Verlag, Stuttgart New York.
- Seeley T.D. (1998) Thoughts on information and integration in honey bee colonies. *Apidologie* 29: 67-80.
- Seeley T.D., Kühnholz S. und Weidenmüller A. (1996) The honey bee's tremble dance stimulates additional bees to function as nectar receivers. *Behav. Ecol. Sociobiol.* 39: 419-427.
- Seeley T.D. (1979) Queen substance dispersal by messenger workers in honeybee colonies. *Behav. Ecol. Sociobiol.* 5: 391-415.
- Shearer D.A. und Boch R. (1965) 2-Heptanone in the Mandibular Gland Secretion of the Honeybee. *Nature* 206: 530.
- Slessor K.N., Kaminski L-A., King G.G.S. und Winston M.L. (1990) Semiochemicals of the honeybee queen mandibular glands. *J. Chem. Ecol.* 16 (3): 851-860.
- Slessor K.N., Kaminski L-A., King G.G.S., Bordens J.H. und Winston M.L. (1988) Semiochemical Basis of the Retinue Response to Queen Honey Bees. *Nature* 332: 354-356.
- Small D.M. (1986) Handbook of Lipid Research Volume 4: The Physical Chemistry of Lipids. Plenum Press, New York London.
- Smith B.H. und Breed M.D. (1995) The chemical basis for nestmate recognition and mate discrimination in social insects. *in: Chemical Ecology of Insects II*, eds: Carde R.T. und Bell W.J., Chapman and Hall, New York.
- StatSoft Inc. (1994) Handbook of Statistica for Windows. Copyright 1984-1994 by StatSoft Inc., Tulsa.
- Steinhoff H. (1948) Untersuchungen über die Haftfähigkeit von Duftstoffen am Bienenkörper. *Z. Vergl. Physiol.* 31: 38-57.
- Stránský K. und Streibl M. (1971) On natural waxes XVIII.: Simple alkyl esters of the wax of

- Stránský K., Streibl M. und Kubelka V. (1971) On natural waxes XXV.: Complex esters of the wax of the honeybee (*Apis mellifera* L.) *Coll.Czech.Chem.Com.* 36: 2281-2297.
- Stránský K., Streibl M. und Sorm F. (1966) Über Naturwachse IV.: Über einen neuen Typ verzweigter Paraffine aus dem Wachs der Honigbiene (*Apis mellifera* L.). *Coll.Czech.Chem.Com.* 31: 4694-4702.
- Streibl M., Stránský K. und Sorm F. (1966) Über einige neue Kohlenwasserstoffe im Wachs der Honigbiene (*Apis mellifica* L.). *Fette, Seifen, Anstrichmittel* 68 (10): 799-805.
- Tautz J. und Lindauer M. (1997) Honeybees establish specific sites on the comb for their waggle dances. *J. Comp. Physiol.* 180: 537-539.
- Tautz J., Rohrseitz K. und Sandemann D.C. (1996) One-Strided Waggle Dance in Bees. *Nature* 382: 32.
- Tietz V. (2000) Zulassungsarbeit in Vorbereitung, Universität Würzburg.
- Tischer J. (1962) Über die Herkunft der gelben Farbstoffe des Bienenwachses. *Hoppe-Seyler's Z. Physiol. Chem.* 267: 14-22.
- Trouiller J., Arnold G., Chappe B., Le Conte Y. und Masson C. (1992) Semiochemical basis of infestation of honey bee brood by *Varroa jacobsoni*. *J. Chem. Ecol.* 18 (11): 2041-2053.
- Tulloch A.P. (1980) Beeswax-Composition and Analysis. *Bee World* 61: 47-62.
- Tulloch A.P. (1971) Beeswax: Structure of the Esters and Their Component Hydroxy Acids and Diols. *Chem. Phys. Lipids* 6: 235-265.
- Ullmann G., Schmidt H., Brotz W., Michalczyk G., Payer W., Dietsche W., Hohner G. und Wildgruber J. (1983) Wachse. in: Ullmanns Encyklopädie der technischen Chemie. Band 24, 4. Auflage, Verlag Chemie, Weinheim.
- Vallet A., Cassier P. und Lensky Y. (1991) Ontogeny of the fine structure of the mandibular glands of the honeybee (*Apis mellifera* L.) workers and the pheromonal activity of 2-Heptanone. *J. Insect. Physiol.* 37 (11): 789-804.
- Vansell G.H. und Bisson C.S. (1935) Origin of Color in Western Beeswax. *J. Econ. Entomol.* 28: 1001-1002.
- Vareschi E. (1971) Duftunterscheidung bei der Honigbiene - Einzelzell-Ableitungen und Verhaltensreaktionen. *Z. vergl. Physiol.* 75: 143-173.
- Vejrosta J., Mikesova M. und Filip P. (1990) Sorption of primary n-alkanols on Tenax. *J. of Chromatography* 518: 234-237.
- Vejrosta J., Mikesova M. und Drozd J. (1989) Sorption of n-alkanes on Tenax. *J. of Chromatography* 464: 394-399.

- Velthuis H.H.W. (1990) Chemical signals and dominance communication in the honeybee *Apis mellifera* (Hymenoptera: Apidae). *Entomol. Gener.* 15 (2): 83-90.
- Verheijen-Voogd C. (1959) How Worker Honeybees Perceive the Presence of Their Queen. *Z. Vergl. Physiol.* 41: 27-582.
- von Frisch K. + (bearb. von Martin Lindauer) (1993) Aus dem Leben der Bienen. 10. Auflage, Springer Verlag, Berlin Heidelberg New York.
- Voogd S. (1955) Inhibition of Ovary Development in Worker Bees by Extraction Fluid of the Queen. *Experientia* 11 (5): 181-182.
- Watmough J. (1997) A general model of pheromone transmission within honey bee hives. *J. theor. Biol.* 189: 159-170.
- Whiffler L.A. und Hepburn H.R. (1991) Inhibition of queen cell construction in the Cape Honeybee, *Apis mellifera capensis*. *Apidologie* 22: 229-236.
- Winston M.L., Higo H.A., Colley S.J., Pankiw T. und Slessor K.N. (1991) The role of queen mandibular pheromone and the colony congestion in honey bee (*Apis mellifera* L.) reproductive swarming (Hymenoptera: Apidae). *J. Insect Behav.* 4(5): 649-660.
- Zerbi G., Gallino G., Del Fanti N. und Bains L. (1989) Structural depth profiling in polyethylene films by multiple internal reflection infra-red spectroscopy. *Polymer* 30: 2324-2327.