

Preface

I would like to take opportunity to thank a number of people without whom the writing of this thesis would not have been possible.

First of all, I am very grateful to Priv-Doz. Dr. *Fritz Boege* for providing excellent working conditions and for his continued support.

I wish to thank Dr.med. *Maxwell Sehested* and Dr.med. *Peter Buhl Jensen* for giving me the opportunity to continue the work on the etoposide resistant cell line.

Many thanks to Prof. *Marie-Christine Dabauvalle* for introducing me to the centrosomes isolation methods.

Thanks is also due to Dr. *Herman Kneitz* for showing me the lymphocyte isolation methods.

In Würzburg the Red Cross provided buffy coats for the experiments.

The KiS1- topoisomerase II α antibody was kindly provided from Prof. Dr. med *H.Kreipe*.

Dr. *Winfried Ness* and Dr. *Jochen Seufert* are thanked for their help.

Dr. *Tobias Straub* is thanked for helping with PCR, and *Hans Ullrich Barthelmes* for helping with the immunofluorescence shown in Fig 7.

I want to thank *Claudia Volff*, *Elisabeth Höfer* and *Silke Feineis*. I have benefitted much from their excellent technical skills.

I am grateful to all the people in *Hormonlabor* for their unlimited help and support.

Finally, I want to thank Prof. *Ulrich Scheer* for his co-supervising.