

Literatur

- Adam, D. (1990). Klonierung und Charakterisierung des melanominduzierenden Gens von *Xiphophorus* und Analyse des Mechanismus der onkogenen Aktivierung. Dissertation, Universität München
- Adam, D., Dimitrijevic, N. & Schartl, M. (1993). Tumor suppression in *Xiphophorus* by an accidentally acquired promoter. *Science* **259**, 816-819
- Adam, D., Mäueler, W. & Schartl, M. (1991). Transcriptional activation of the melanoma inducing *Xmrk* oncogene in *Xiphophorus*. *Oncogene* **6**, 73-80
- Ahuja, M.R. (1979). On the nature of genetic change as an underlying cause for the origin of neoplasms. In: Antiviral mechanisms in the control of neoplasia. pp 17-37, Chandra, P. (ed), Plenum Press, New York
- Ahuja, M.R., Schwab, M. & Anders, F. (1980). Linkage between a regulatory locus for melanoma cell differentiation and an esterase locus in *Xiphophorus*. *J. Hered.* **71**, 403-407
- Ahuja, M.R. & Anders, F. (1976). A genetic concept of the origin of cancer, based in part upon studies of neoplasms in fishes. *Prog. Exp. Tumor Res.* **20**, 380-397
- Altschmied, J., Hornung, U., Schlupp, I., Gadau, J., Kolb, R. & Schartl, M. (1997a). Isolation of DNA suitable for PCR for field and laboratory work. *BioTechniques* **23**, 228-229
- Altschmied, J., Ditzel, L. & Schartl, M. (1997b). Hypomethylation of the *Xmrk* oncogene promoter in melanoma cells of *Xiphophorus*. *Biol. Chem.* **378**, 1457-1466
- Amor, M., Parker, K., Globerman, H., New, M. & White, P. (1988). Mutation in the *CYP21B* gene (Ile-172->Asn) causes steroid 21-hydroxylase deficiency. *Proc. Natl. Acad. Sci. USA* **85**, 1600-1604
- Anders, A., Petry, H., Fleming, C., Petry, K., Brix, P., Luke, W., Groger, H., Schneider, E., Kiefer, J. & Anders, F. (1994). Increasing melanoma incidence: putatively explainable by retrotransposons. Experimental contributions of the Xiphophorine Gordon-Kosswig melanoma system. *Pigm. Cell Res.* **7**, 433-450
- Anders, A., Gröger, H., Anders, F., Zechel, C., Smith, A. & Schlatterer, B. (1991). Discrimination of initiating and promoting carcinogens in fish. *Ann. Rech. Vet.* **22**, 273-294
- Anders, A., Anders, F. & Klinke, K. (1973). Regulation of gene expression in the Gordon-Kosswig melanoma system. In: Genetics and Mutagenesis of Fish. pp 33-63, Schroeder, J.H. (ed), Springer, New York
- Anders, F. (1991). Contributions of the Gordon-Kosswig melanoma system to the present concept of neoplasia. *Pigm. Cell Res.* **3**, 7-29
- Anders, F., Schartl, M., Barnekow, A. & Anders, A. (1984). *Xiphophorus* as an in vivo model for studies on normal and defective control of oncogenes. *Adv. Cancer Res.* **42**, 191-275
- Anders, F., Diehl, H. & Scholl, E. (1980). Differentiation of normal melanophores and neoplastically transformed melanophores in the skin of fishes of genus *Xiphophorus*. In: The

Skin of Vertebrates. pp 211-218, Spearman, R.I.C. & Riley, P.A. (eds), Linnean Society Symposium Series 9

Anderson, J., Burns, H.D., Enriquez-Harris, P., Wilkie, A.O.M. & Heath, J.K. (1998). Apert syndrome mutations in fibroblast growth factor receptor 2 exhibit increased affinity for FGF ligand. *Hum. Mol. Genet.* **7**, 1475-1483

Aroian, R.V., Koga, M., Mendel, J.E., Ohshima, Y. & Sternberg, P.W. (1990). The *let-23* gene necessary for *Caenorhabditis elegans* vulval induction encodes a tyrosine kinase of the EGF receptor subfamily. *Nature* **348**, 693-699

Bachman, K.E., Herman, J.G., Corn, P.G., Merlo, A., Costello, J.F., Cavenee, W.K., Baylin, S.B. & Graff, J.R. (1999). Methylation-associated silencing of the tissue inhibitor of metalloproteinase-3 gene suggest a suppressor role in kidney, brain, and other human cancers. *Cancer Res.* **59**, 798-802

Baltimore, D. (1981). Gene conversion: some implications for immunoglobulin genes. *Cell* **24**, 592-594

Baudler, M., Duschl, J., Winkler, C., Schartl, M. & Altschmied, J. (1997). Activation of transcription of the melanoma inducing *Xmrk* oncogene by a GC box element. *J. Biol. Chem.* **272**, 131-137

Bishop, J.M. (1987). The molecular genetics of cancer. *Science* **235**, 305-311

Blin, N. & Stafford, D.W. (1976). A general method for isolation of high molecular weight DNA from eukaryotes. *Nucl. Acids. Res.* **3**, 2303-2308

Bullock, W.O., Fernandez, J.M. & Short, J.M. (1987). XL1-Blue: a high efficiency plasmid transforming recA Escherichia coli strain with b-galactosidase selection. *Bio Techniques* **5**, 376-378

Burgdorf, C., Welzel, K., Hasenbank, R., Zehetner, G., Weis, S. & Lehrach, H. (1998). Gridded genomic libraries of different chordate species: a reference library system for basic and comparative studies of chordate genomes. *Genomics* **52**, 230-232

Call, K.M., Glaser, T., Yto, C.Y., Buckler, A.J., Pelletier, J., Haber, D.A., Rose, E.A., Kral, A., Yeger, H., Lewis, W.H., Jones, C. & Housman, D.E. (1990). Isolation and characterization of a zinc finger polypeptide gene at the human chromosome 11 Wilm's tumor locus. *Cell* **60**, 509-520

Callaghan, T., Antczak, M., Flickinger, T., Raines, M., Myers, M. & Kung H.J. (1993). A complete description of the EGF-receptor exon structure: implication in oncogenic activation and domain evolution. *Oncogene* **8**, 2939-2948

Campbell, L., Potter, A., Ignatius, J., Dubowitz, V. & Davies, K. (1997). Genomic variation and gene conversion in spinal muscular atrophy: implications for disease process and clinical phenotype. *Am. J. Hum. Genet.* **61**, 40-50

Clifford, R. & Schüpbach, T. (1994). Molecular analysis of the *Drosophila* EGF receptor homolog reveals that several genetically defined classes of alleles cluster in subdomains of the receptor protein. *Genetics* **137**, 531-550

Collins, F.S. (1992). Positional cloning: Let's not call it reverse anymore. *Nature Genet.* **1**, 3-6

- Coughlan, T., Schartl, M., Hope, I. & Stewart, A. (1999). PCR based sex test for *Xiphophorus maculatus*. *J. Fish Biol.* **54**, 218-222
- Chrysogelos, S.A. (1993). Chromatin structure of the EGFR gene suggests a role for intron 1 sequences in its regulation in breast cancer cells. *Nucl. Acids Res.* **21**, 5736-5741
- Deeb, S.S., Alvarez, A., Malkki, M. & Motulsky, A.G. (1995). Molecular patterns and sequence polymorphisms in the red and green visual pigment genes of Japanese men. *Hum. Genet.* **95**, 501-506
- Del Sal, G., Manfioletti, G. & Schneider, C. (1988). One tube plasmid DNA minipreparation suitable for sequencing. *Nucl. Acids. Res.* **16**, 9878
- Dimitrijevic, N., Winkler, C., Wellbrock, C., Gómez, A., Duschl, J., Altschmied, J. & Schartl, M. (1998). Activation of the *Xmrk* proto-oncogene of Xiphophorus by overexpression and mutational alterations. *Oncogene* **16**, 1681-1690
- Donis-Keller, H., Green, P., Helms, C., Cartinhour, S., Weiffenbach, B., Stephens, K., Keith, T.P., Bowden, D.W., Smith, D.R., Lander, E.S., et al. (1987). A genetic linkage map of the human genome. *Cell* **51**, 319-337
- Dryja, T.P., Rapaport, J.M., Joyce, J.M. & Petersen, R.A. (1986). Molecular detection of deletions involving band q14 of chromosome 13 in retinoblastomas. *Proc. Natl. Acad. Sci. USA* **83**, 7391-7394
- Dunn, I.S. & Blattner, F.R. (1987). Charons 36 to 40: Multi enzyme, high capacity, recombination deficient replacement vectors with polylinkers and polystuffers. *Nucl. Acids Res.* **15**, 2677-2698
- Dzwillo, M. & Zander, C.D. (1967). Geschlechtsbestimmung bei Zahnkarpen (pisces). *Mitt. Hamburg Zool. Mus. Inst.* **64**, 47-162
- Esteller, M., Sanchez-Cespedes, M., Rosell, R., Sidransky, D., Baylin, S.B. & Herman, J.G. (1999). Detection of aberrant promoter hypermethylation of tumor suppressor genes in serum DNA from non-small cell lung cancer patients. *Cancer Res.* **59**, 67-70
- Feinberg, A.P. & Vogelstein, B. (1983). A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* **132**, 6-13
- Flickinger, T.W., Maihle, N.J. & Kung, H.J. (1992). An alternatively processed mRNA from the Avian *c-erbB* gene encodes a soluble, truncated form of the receptor that can block ligand-dependent transformation. *Mol. Cell. Biol.* **12**, 883-893
- Förnzler, D. (1994). Untersuchungen zur Regulation des *Xmrk*-Onkogens von Xiphophorus: Isolierung molekularer Marker für das Regulatorgen *R*, Methylierungsstatus, Charakterisierung des neuen Genlocus *D*. Dissertation, Universität Würzburg
- Förnzler, D., Altschmied, J., Nanda, I., Kolb, R., Baudler, M., Schmid, M. & Schartl, M. (1996). The *Xmrk* oncogene promoter is derived from a novel amplified locus of unusual organization. *Genome Res.* **6**, 102-113
- Förnzler, D., Wittbrodt, J. & Schartl, M. (1991). Analysis of an Esterase linked to a locus involved in the regulation of the melanoma oncogene and isolation of polymorphic marker sequences in *Xiphophorus*. *Biochem. Genet.* **29**, 509-524

- Franklin, G.C., Donovan, M., Adam, G.I., Holmgren, L., Pfeifer-Ohlsson, S. & Ohlsson, R. (1991). Expression of the human PDGF-B gene is regulated by both positively and negatively acting cell type-specific regulatory elements located in the first intron. *EMBO J.* **10**, 1365-1373
- Friend, S.H., Bernards, R., Rogelj, S., Weinberg, R.A., Rapaport, J.M., Albert, D.M. & Dryja, T.P. (1986). A human DNA segment with properties of the gene that predisposes to retinoblastoma and osteosarcoma. *Nature* **323**, 643-646
- Gessler, M., Poustka, A., Cavenee, W., Neve, R.L., Orkin, S.H. & Bruns, G. (1990). Homozygous deletion in Wilm's tumors of a zinc-finger gene identified by chromosome jumping. *Nature* **343**, 774-778
- Gonzalez-Zulueta, M., Bender, C.M., Yang, A.S., Nguyen, T., Beart, R.W., Van Tornout J.M. & Jones, P.A. (1995). Methylation of the 5' CpG island of the p16/CDKN2 tumor suppressor gene in normal and transformed human tissues correlates with gene silencing. *Cancer Res.* **55**, 4531-4535
- Gordon, M. (1948). Effects of five primary genes on the site of melanomas in fishes and the influence of two color genes on their pigmentation. In: The biology of melanomas. pp 216-268, New York Academy of Science, New York
- Gordon, M. (1927). The genetics of a viviparous top-minnow Platypoecilus; the inheritance of two kinds of melanophores. *Genetics* **12**, 253-283
- Gordon, H. & Gordon, M. (1957). Maintenance of polymorphism by potentially injurious genes in eight natural populations of the platyfish, *Xiphophorus maculatus*, *J. Genetics* **55**, 1-44
- Grimm, T., Müller, B., Dreier, M., Kind, E., Bettecken, T., Meng, G. & Müller, C.R. (1989). Hot spot of recombination within DDXS164 in the Duchenne muscular dystrophy gene. *Am. J. Hum. Genet.* **45**, 368-372
- Gruis, N.A., van der Velden, P.A., Sandkuijl, L.A., Prins, D.E., Weaver-Feldhaus, J., Kamb, A., Bergman, W. & Frants, R.R. (1995). Homozygotes for CDKN2 (p16) germline mutation in Dutch familial melanoma kindreds. *Nature Genet.* **10**, 351-353
- Gutbrod, H. (1996). Versuch zur Isolierung molekularer Markersequenzen für den Regulatorlocus *R* von Xiphophorus mit der RAPD-Technik. Diplomarbeit, Universität Würzburg
- Gutbrod H. and Schartl M. (1999). Intragenic sex-chromosomal cross-overs of Xmrk oncogene alleles affect pigment pattern formation and the severity of melanoma in Xiphophorus. *Genetics*, **151**, 773-783
- Hanada, M., Delia, D., Aiello, A., Stadtmauer, E. & Reed, J.C. (1993). bcl-2 gene hypomethylation and high-level expression in B-cell chronic lymphocytic leukemia. *Blood* **82**, 1820-1828
- Häussler, G. (1928). Über Melanombildung bei Bastarden von *Xiphophorus maculatus* var. *rubra*. *Klin. Wochenschr.* **7**, 1561-1562
- Henikoff, S. & Matzke, M.A. (1997). Exploring and explaining epigenetic effects. *Trends Genet.* **13**, 293-295
- Hess, J.F., Schmid, C.W. & Shen, C.K. (1984). A gradient of sequence divergence in the human adult alpha-globin duplication units. *Science* **226**, 67-70

- Higashi, Y., Tanae, A., Inoue, H., Hiromasa, T. & Fujii-Kuriyama, Y. (1988). Aberrant splicing and missense mutations cause steroid 21-hydroxylase P-450 (C21) deficiency in humans: possible gene conversion products. *Proc. Natl. Acad. Sci. USA.* **85**, 7486-7490
- Holliday, R. (1964). A mechanism for gene conversion in fungi. *Genet. Res.* **78**, 273-287
- Hussussian, C.J., Struewing, J.P., Goldstein, A.M., Higgins, P.A., Ally, D.S., Sheahan, M.D., Clark, W.H. Jr., Tucker, M.A. & Dracopoli, N.C. (1994). Germline *p16* mutations in familial melanoma. *Nature Genet.* **8**, 15-21
- Jansen, R. & Ledley, F.D. (1990). Disruption of phase during PCR amplification and cloning of heterozygous target sequences. *Nucl. Acids. Res.* **18**, 5153-5156
- Kallman, K.D. (1975). The platyfish, *Xiphophorus maculatus*. In: Handbook of Genetics. pp 81-132, King, R.C. (ed), Plenum Press, New York
- Kallman, K.D. (1970). Sex determination and the restriction of pigment pattern to the X and Y chromosomes in populations of a poeciliid fish, *Xiphophorus maculatus*, from the Sibun and Belize Rivers of British Honduras. *Zoologica* **55**, 1-16
- Kallman, K.D. & Schreibman, M.P. (1971). The origin and possible genetic control of new, stable pigment patterns in the Poeciliid fish *Xiphophorus maculatus*. *J. Exp. Zool.* **176**, 147-168
- Kamb, A. (1995). Cell-cycle regulators and cancer. *Trend Genet.* **11**, 136-140
- Kazianis, S. (1995). A molecular approach to the study of the Xiphophorus melanoma model systems . Dissertation, Universität New York
- Kazianis, S., Gutbrod, H., Nairn, R.S., McEntire, B.B., Della Coletta, L., Walter, R.B., Borowsky, R.L., Woodhead, A.D., Setlow, R.B., Schartl, M. & Morizot, D.C. (1998). Localization of a *CDKN2* gene in linkage group V of *Xiphophorus* fishes defines it as a candidate for the *DIFF* tumor suppressor. *Genes, Chromosomes & Cancer* **22**, 210-220
- Kazianis, S., Morizot, D.C., McEntire, B.B., Nairn, R.S. & Borowsky, R.L. (1996). Genetic mapping in Xiphophorus hybrid fish: assignment of 43 AP-PCR/RAPD and isozyme markers to multipoint linkage groups. *Genome Res.* **6**, 280-289
- Kim, Y.S., Kim, M.H., Choi, Y.K., Kim, C.H. & Lee, D.S. (1994). Allelic divergence in the human insulin gene provides evidence for intragenic recombination events in the non-coding regions: evidence for existence of new alleles. *Mol. Gen. Genet.* **245**, 146-151
- Kmiec, E.B., Angelides, K.J. & Holloman, W.K. (1985). Left-handed DNA and the synaptic pairing reaction promoted by *Ustilago rec1* protein. *Cell* **40**, 139-145
- Kobori, J.A., Strauss, E., Minard, K. & Hood, L. (1986). Molecular analysis of the hotspot of recombination in the murine major histocompatibility complex. *Science* **234**, 173-179
- Konopka, J.B., Watanabe, S.M. & Witte, O.N. (1984). An alteration of the human c-abl protein in K562 leukemia cells unmasks associated tyrosine kinase activity. *Cell* **37**, 1035-1042
- Kosswig, C. (1939). Die Geschlechtsbestimmung beim Zahnkarpfen. *Univ. Fen. Falk. Yeni. Seri.* **4**, 91-144
- Kosswig, C. (1928). Über Kreuzungen zwischen den Teleostiern *Xiphophorus helleri* und *Platypoecilus maculatus*. *Z. Indukt. Abstammungs- Vererbungsl.* **47**, 150-158

- Lafuse, W.P. & David, C.S. (1986). Recombination hotspots within the I region of the mouse H-2 complex map to the E beta and E alpha genes. *Immunogenetics* **24**, 352-360
- Lander, E.S., Green, P., Abrahamson, J., Barlow, A., Daly, M.J., Lincoln, S.E. & Newburg, L. (1987). Mapmaker: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* **1**, 174-181
- Lax, I., Johnson, A., Howk, R., Sap, J., Bellot, F., Winkler, M., Ullrich, A., Vennstrom, B., Schlessinger, J. & Givol, D. (1988). Chicken epidermal growth factor (EGF) receptor: cDNA cloning, expression in mouse cells, and differential binding of EGF and transforming growth factor alpha. *Mol. Cell. Biol.* **8**, 1970-1978
- Lee, H. & Maihle, N.J. (1998). Isolation and characterization of four alternate *c-erbB3* transcripts expressed in ovarian carcinoma-derived cell lines and normal human tissues. *Oncogene* **16**, 3243-3252
- Lehrman, M.A., Schneider, W.J., Sudhof, T.C., Brown, M.S., Goldstein, J.L. & Russell, D.W. (1985). Mutation in LDL receptor: Alu-Alu recombination deletes exons encoding transmembrane and cytoplasmic domains. *Science* **227**, 140-146
- Li, Y., Nichols, M.A., Shay, J.W. and Xiong, Y. (1994). Transcriptional repression of the D-type cyclin-dependent kinase inhibitor p16 by the retinoblastoma susceptibility gene product pRb. *Cancer Res.* **54**, 6078-6082
- Liskay, R.M., Letsou, A. & Stachelek, J.L. (1987). Homology requirement for efficient gene conversion between duplicated chromosomal sequences in mammalian cells. *Genetics* **115**, 161-167
- Lo Ten Foe, J.R., Kwee, M.L., Rooimans, M.A., Oostra, A.B., Veerman, A.J., van Weel, M., Pauli, R.M., Shahidi, N.T., Dokal, I., Roberts, I., Altay, C., Gluckman, E., Gibson, R.A., Mathew, C.G., Arwert, F. & Joenje, H. (1997). Somatic mosaicism in Fanconi anemia: molecular basis and clinical significance. *Eur. J. Hum. Genet.* **5**, 137-148
- Lois, A.F., Cooper, L.T., Geng, Y., Nobori, T. & Carson, D.A. (1995). Expression of the p16 and p15 cyclin-dependent kinase inhibitors in lymphocyte activation and neuronal differentiation. *Cancer Res.* **55**, 4010-4013
- Longley, B.J. Jr., Metcalfe, D.D., Tharp, M., Wang, X., Tyrrell, L., Lu, S.Z., Heitjan, D. & Ma, Y. (1999). Activating and dominant inactivating c-KIT catalytic domain mutations in distinct clinical forms of human mastocytosis. *Proc. Natl. Acad. Sci. USA* **96**, 1609-1614
- Maeda, N. & Smithies, O. (1986). The evolution of multigene families: human haptoglobin genes. *Annu. Rev. Genet.* **20**, 81-108
- Maekawa, T., Imamoto, F., Merlino, G.T., Pastan, I. & Ishii, S. (1989). Cooperative function of two separate enhancers of the human epidermal growth factor receptor proto-oncogene. *J. Biol. Chem.* **264**, 5488-5494
- Mäuelter, W., Schartl, A. & Schartl, M. (1993). Different expression patterns of oncogenes and proto-oncogenes in hereditary and carcinogen-induced tumors of Xiphophorus. *Int. J. Cancer* **55**, 288-296
- Maihle, N.J., Flickinger, T.W., Raines, M.A., Sanders, M.L. & Kung, H.-J. (1991). Native avian *c-erbB* gene expresses a secreted protein product corresponding to the ligand-binding domain of the receptor. *Proc. Natl. Acad. Sci. USA* **88**, 1825-1829

- Manly, K.F. (1993). A Macintosh program for storage and analysis of experimental mapping data. *Mamm. Genome* **4**, 303-313
- McCormack, W.T., Tjoelker, L.W. & Thompson, C.B. (1991). Avian B-cell development: generation of an immunoglobulin repertoire by gene conversion. *Annu. Rev. Immunol.* **9**, 219-241
- Meselson, M. S. & Radding, C.M. (1975). A general model for genetic recombination. *Proc. Natl. Acad. Sci. USA* **72**, 358-361
- Meyer, A., Morrissey, J.M. & Schartl, M. (1994). Recurrent origin of a sexually selected trait in *Xiphophorus* fishes inferred from a molecular phylogeny. *Nature* **368**, 539-542
- Morizot, D.C., Nairn, R.S., Walter, R.B. & Kazianis, S. (1998). The linkage map of *Xiphophorus* fishes. *ILAR* **39**, 237-248
- Morizot, D.C., Slaugenhaupt, S.A., Kallman, K.D. & Chakravarti, A. (1991). Genetic linkage map of fishes of the genus *Xiphophorus*. *Genetics* **127**, 399-410
- Morizot, D.C. & Siciliano, M.J. (1983). Linkage group V of platyfishes and swordtails of the genus *Xiphophorus* (Poeciliidae): Linkage of loci for malate dehydrogenase-2 and esterase-1 and esterase-4 with a gene controlling the severity of hybrid melanomas. *J. Natl. Cancer Inst.* **71**, 809-813
- Morris, D.W. & Dutra, J.C. (1997). Identification of a MMTV insertion mutation within the coding region of the Fgf-3 protooncogene. *Virology* **238**, 161-165
- Murray, N.E., Brammar, W.J. & Murray, K. (1977). Lamboid phages that simplify the recovery of in vitro recombinants. *Mol. Gen. Genet.* **150**, 53-61
- Nairn, R.S., Kazianis, S., McEntire, B.B., Della Coletta, L., Walter, R.B. & Morizot, D.C. (1996). A CDKN2-like polymorphism in *Xiphophorus* linkage group V is associated with UV-B-induced melanoma formation in platyfish-swordtail hybrids. *Proc. Natl. Acad. Sci. USA* **93**, 13042-13047
- Nanda, I., Weis, S., Förnzler, D., Altschmied, J., Schartl, M. & Schmid, M. (1996). Clustered organization and conservation of the *Xiphophorus maculatus* D locus, which includes two distinct gene sequences. *Chromosoma* **105**, 242-249
- Nathans, J., Piantanida, T.P., Eddy, R.L., Shows, T.B. & Hogness, D.S. (1986). Molecular genetics of inherited variation in human color vision. *Science* **232**, 203-210
- Nobori, T., Miura, K., Wu D.J., Lois, A., Takabayashi, K. & Carson, D.A. (1994). Deletions of the cyclin-dependent kinase 4 inhibitor gene in multiple human cancers. *Nature* **368**, 753-756
- Okamoto, A., Demetrick, D.J., Spillare, E.A., Hagiwara, K., Hussain, S.P., Bennett, W.P., Forrester, K., Gerwin, B., Serrano, M., Beach, D.H. & Harris, C.C. (1994). Mutations and altered expression of p16INK4 in human cancer. *Proc. Natl. Acad. Sci. USA* **91**, 11045-11049
- Otsen, M., den Bieman, M., Kuiper, M.T.R., Pravenec, M., Kren, V., Kurtz, T.W., Jacob, H.J., Lankhorst, A. & van Zutphen, B.F.M. (1996). Use of AFLP markers for gene mapping and QTL detection in the rat. *Genomics* **37**, 289-294

- Pagany, M. (1999). Das Melanommodellsystem Xiphophorus: "Charakterisierung von Antigenen im malignen Melanom und Untersuchungen zur genomischen Instabilität". Dissertation, Universität Würzburg
- Parry, D., Bates, S., Mann, D.J. & Peters, G. (1995). Lack of cyclin D-Cdk complexes in Rb-negative cells correlates with high levels of p16INK4/MTS1 tumour suppressor gene product. *EMBO J.* **14**, 503-511
- Penhoat, A., Naville, D., Jaillard, C., Durand, P. & Begeot, M. (1997). Presence of multiple functional polyadenylation signals in the 3'-untranslated region of human corticotropin receptor cDNA. *Biochim. Biophys. Acta* **1356**, 249-252
- Petch, L.A., Harris, J., Raymond, V.W., Blasband, A., Lee, D.C. & Earp, H.S. (1990). A truncated, secreted form of the epidermal growth factor receptor is encoded by an alternatively spliced transcript in normal rat tissue. *Mol. Cell. Biol.* **10**, 2973-2982
- Pierotti, M.A., Santoro, M., Jenkins, R.B., Sozzi, G., Bongarzone, I., Grieco, M., Monzini, N., Miozzo, M., Herrmann, M.A., Fusco, A., Hay, J.D., Della Porta, G. & Vecchio, G. (1992). Characterization of an inversion on the long arm of chromosome 10 juxtaposing D10S170 and RET and creating the oncogenic sequence RET/PTC. *Proc. Natl. Acad. Sci. USA* **89**, 1616-1620
- Pittman, D.L. & Schimenti, J.C. (1998). Recombination in the mammalian germ line. In: Meiosis and Gametogenesis. pp 2-35, Handel, M.A. (ed), Academic Press, San Diego
- Ponder, B.A. (1990). Inherited predisposition to cancer. *Trends Genet.* **6**, 213-218
- Postlethwait, J.H., Johnson, S.L., Midson, C.N., Talbot, W.S., Gates, M., Ballinger, E.W., Africa, D., Andrews, R., Carl, T., Eisen, J.S., Horne, S., Kimmel, C.B., Hutchinson, M., Johnson, M. & Rodriguez, A. (1994). A genetic linkage map for the Zebrafish. *Science* **264**, 699-703
- Rabbitts, T.H. (1994). Chromosomal translocations in human cancer. *Nature* **372**, 143-149
- Raulf, F. (1991). Untersuchungen zur Struktur, Expression und Evolution von c-src: Proto-Onkogen und aktiviertes Onkogen in Xiphophorus (Teleostei: Poeciliidae). Dissertation, Ludwig-Maximilians-Universität, München
- Razin, A. & Cedar, H. (1991). DNA methylation and gene expression. *Microbiol. Rev.* **55**, 451-458
- Reeves, R.H., Crowley, M.R., O'Hara, B.F. & Gearhart, J.D. (1990). Sex, strain, and species differences affect recombination across an evolutionary conserved segment of mouse chromosome 16. *Genomics* **8**, 141-148
- Reiter, J.L. & Maihle, N.J. (1996). A 1.8 kb alternative transcript from the human epidermal growth factor receptor gene encodes a truncated form of the receptor. *Nucl. Acids Res.* **24**, 4050-4056
- Rich, A., Nordheim, A., Wang, A.H.-J. (1984). The chemistry and biology of left-handed Z-DNA. *Ann. Rev. Biochem.* **53**, 791-846
- Rosen, D. (1979). Fishes from uplands and intermountain basins of Guatemala. Revisionary studies and comparative geography. *Bull. Am. Mus. Nat. Hist.* **162**, 267-367
- Rouppé van der Voort, J.N., van Zandvoort, P., van Eck, H.J., Folkertsma, R.T., Hutten, R.C., Draaijstra, J., Gommers, F.J., Jacobsen, E., Helder, J. & Bakker, J. (1997). Use of

allele specificity of comigrating AFLP markers to align genetic maps from different potato genotypes. *Mol. Gen. Genet.* **255**, 438-447

Rubnitz, J. & Subramani, S. (1986). Extrachromosomal and chromosomal gene conversion in mammalian cells. *Mol. Cell. Biol.* **6**, 1608-1614

Saiki, R.K., Gelfand, D.H., Stoffel, B., Scharf, S.J., Higushi, R., Horn, G.T., Mullis, K.B. & Ehrlich, H.A. (1988). Primer directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science* **239**, 487-490

Sambrook, J., Fritsch, E.F. & Maniatis, T. (1989). Molecular cloning - a laboratory manual. Cold Spring Harbor Laboratory Press, Cold Spring Harbor

Sanger, F., Nicklen, S. & Coulson, A. R. (1977). DNA sequencing with chain terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**, 5463-5467

Schartl, M. (1990). Homology of melanoma-inducing loci in the genus *Xiphophorus*. *Genetics* **126**, 1083-1091

Schartl, M. (1988). A sex chromosomal restriction-fragment-length marker linked to melanoma-determining *Tu* loci in *Xiphophorus*. *Genetics* **119**, 679-685

Schartl, M., Hornung, U., Gutbrod, H., Volff, J.N. & Wittbrodt, J. (1999). Melanoma loss-of-function mutants in *Xiphophorus* caused by *Xmrk*-oncogene deletion and gene-disruption by a transposable element. *Genetics*, im Druck

Schwab, M. (1982). Genome organization in *Xiphophorus* (Poeciliidae; Teleostei). *Mol. Gen. Genet.* **188**, 410-417

Senterre-Lesenfants, S., Alag, A.S. & Sobel, M.E. (1995). Multiple mRNA species are generated by alternate polyadenylation from the human calmodulin-I gene. *J. Cell Biochem.* **58**, 445-454

Setlow, R.B., Woodhead, A.D. & Grist, E. (1989). Animal model for ultraviolet radiation-induced melanoma: Platfish-swordtails hybrid. *Proc. Nat. Acad. Sci. USA* **86**, 8922-8926

Shaw, C.R. & Prasad, R. (1970). Starch gel electrophoresis of enzymes - a compilation of recipes. *Biochem. Genet.* **4**, 297-320

Sheffield, V.C., Beck, J.S., Kwitek, A.E., Sandstrom, D.W. & Stone, E.M. (1993). The sensitivity of single-strand conformation polymorphism analysis for the detection of single base substitutions. *Genomics* **16**, 325-332

Shinagawa, T., Ishiguro, N., Horiuchi, M., Matsui, T., Okada, K. & Shinagawa, M. (1997). Deletion of *c-myb* exon 9 induced by insertion of repeats. *Oncogene* **14**, 2775-2783

Shiroishi, T., Sagai, T., Hanzawa, N., Gotoh, H. & Moriwaki, K. (1991). Genetic control of sex-dependent meiotic recombination in the major histocompatibility complex of the mouse. *EMBO J.* **10**, 681-686

Shiroishi, T., Hanzawa, N., Sagai, T., Ishiura, M., Gojobori, T., Steinmetz, M. & Moriwaki, K. (1990). Recombinational hotspot specific to female meiosis in the mouse major histocompatibility complex. *Immunogenetics* **31**, 79-88

Siciliano, M.J. & Shaw, C.R. (1976). Separation and visualization of enzymes on gels. In: Chromatographic and Electrophoretic Techniques. pp 185-209, Smith (ed), Heinemann, London

- Siciliano, M.J., Morizot, D.C. & Wright, D.A. (1976). Factors responsible for platyfish-swordtail hybrid melanoma - many or few? In: Melanomas: Basic Properties and Clinical Behaviour. pp 47-54, Riley, V. (ed), Karger, Basel
- Simons, G., van der Lee, T., Diergaard, P., van Daelen, R., Groenendijk, J., Frijters, A., Buschges, R., Hollricher, K., Topsch, S., Schulze-Lefert, P., Salamini, F., Zabeau, M. & Vos, P. (1997). AFLP-based fine mapping of the *Mlo* gene to a 30 kb DNA segment of the barley genome. *Genomics* **44**, 61-70
- Slightom, J.L., Blechl, A.E. & Smithies, O. (1980). Human fetal G γ - and A γ -globin genes: complete nucleotide sequences suggest that DNA can be exchanged between these duplicated genes. *Cell* **21**, 627-638
- Southern, E.M. (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* **98**, 503-517
- Stark, G.R. & Wahl, G.M. (1984). Gene amplification. *Annu. Rev. Biochem.* **53**, 447-491
- Steinmetz, M., Stephen, D. & Lindahl, K.F. (1986). Gene organization and recombinational hotspots in the murine major histocompatibility complex. *Cell* **44**, 895-904
- Stringer, J.R. (1985). Rekombination between poly[d(GT)-d(CA)] sequences in simian virus 40-infected cultured cells. *Mol. Cell. Biol.* **5**, 1247-1259
- Szostak, J.W., Orr-Weaver, T.L., Rothstein, R.J. & Stahl, F.W. (1983). The double-strand-break repair model for recombination. *Cell* **33**, 25-35
- Tam, S.W., Shay, J.W. & Pagano, M. (1994). Differential expression and cell cycle regulation of the cyclin-dependent kinase 4 inhibitor p16^{INK4}. *Cancer Res.* **54**, 5816-5820
- Tracy, R.B., Baumohl, J.K. & Kowalczykowski, S.C. (1997). The preference for GT-rich DNA by the yeast Rad51 protein defines a set of universal pairing sequences. *Genes Dev.* **11**, 3423-3431
- Ullrich, A. & Schlessinger, J. (1990). Signal transduction by receptors with tyrosine kinase activity. *Cell* **61**, 203-212
- Urabe, K., Kimura, A., Harada, F., Iwanaga, T. & Sasazuki, T. (1990). Gene conversion in steroid 21-hydroxylase genes. *Am. J. Hum. Genet.* **46**, 1178-1186
- van Endert, P.M., Lopez, M.T., Patel, S.D., Monaco, J.J. & McDevitt, H.O. (1992). Genomic polymorphism, recombination, and linkage disequilibrium in human major histocompatibility complex - encoded antigen-processing genes. *Proc. Natl. Acad. Sci. USA* **89**, 11594-11597
- Varras, M.N., Sourvinos, G., Diakomanolis, E., Koumantakis, E., Flouris, G.A., Lekka-Katsouli, J., Michalas, S. & Spandidos, D.A. (1999). Detection and clinical correlations of ras gene mutations in human ovarian tumors. *Oncology* **56**, 89-96
- Vielkind, J.R., Kallman, K.D. & Morizot, D.C. (1989). Genetics of melanomas in *Xiphophorus* fishes. *J. Aqua. Anim. Health* **1**, 69-77
- Vielkind, U. (1976). Genetic control of cell differentiation in platyfish-swordtail melanomas. *J. Exp. Zool.* **196**, 197-204

- Vielkind, U., Schlage, W. & Anders, F. (1977). Melanogenesis in genetically determined pigment cell tumours of platyfish and platy-swordtail hybrids: correlation between tyrosinase activity and degree of malignancy. *Z. Krebsforsch.* **90**, 285-299
- Vogelstein, B. & Kinzler, W. (1993). The multistep nature of cancer. *Trends Genet.* **9**, 138-141
- Vos, P., Hogers, R., Bleeker, M., Reijans, M., Van de Lee, T., Hornes, M., Frijters, A., Pot, J., Peleman, J., Kuiper, M. & Zabeau, M. (1995). AFLP: a new technique for DNA fingerprinting. *Nucl. Acids Res.* **23**, 4407-4414
- Wada, H., Naruse, K., Shimada, A. & Shima, A. (1995). A genetic linkage map of a fish, the Japanese medaka *Oryzias latipes*. *Mol. Mar. Biol. Biotechnol.* **4**, 269-274
- Wahls, W.P., Wallace, L.J. & Moore, P.D. (1990a). Hypervariable minisatellite DNA is a hotspot for homologous recombination in human cells. *Cell* **60**, 95-103
- Wahls, W.P., Wallace, L.J. & Moore, P.D. (1990b). The Z-DNA motif d(TG)₃₀ promotes reception of information during gene conversion events while stimulating homologous recombination in human cells in culture. *Mol. Cell. Biol.* **10**, 785-793
- Wakamatsu, Y., Oikawa, A., Obika, M., Hirobe, T., & Ozato, K. (1984). Fish hereditary melanoma cell lines of different degrees of cell differentiation. *Dev. Growth Differ.* **26**, 503-513
- Wakamatsu, Y. (1981). Establishment of a cell line from platyfish-swordtail hybrid melanoma. *Cancer Res.* **41**, 679-680
- Wang, J. & Walsh, K. (1996). Resistance to apoptosis conferred by Cdk inhibitors during myocyte differentiation. *Science* **273**, 359-361
- Watson, C.J., Gaunt, L., Evans, G., Patel, K., Harris, R. & Strachan, T. (1993). A disease-associated germline deletion maps the type 2 neurofibromatosis (NF2) gene between the Ewing sarcoma region and the leukaemia inhibitory factor locus. *Hum. Mol. Genet.* **2**, 701-704
- Weinberg, R.A. (1989). Oncogenes, antioncogenes, and the molecular basis of multistep carcinogenesis. *Cancer Res.* **49**, 3713-3721
- Weis, S. & Schartl, M. (1998). The macromelanophore locus and the melanoma oncogene *Xmrk* are separate genetic entities in the genome of *Xiphophorus*. *Genetics* **149**, 1909-1920
- Welsh, J. & McClelland, M. (1990). Fingerprinting genomes using PCR with arbitrary primers. *Nucl. Acids Res.* **18**, 7213-7218
- Winkler, C., Wittbrodt, J., Lammers, R., Ullrich, A. & Schartl, M. (1994). Ligand-dependent tumor induction in medakafish embryos by a *Xmrk* receptor tyrosine kinase. *Oncogene* **9**, 1517-1525
- Wittbrodt, J., Lammers, R., Malitschek, B., Ullrich, A. & Schartl, M. (1992). The *Xmrk* receptor tyrosine kinase is activated in *Xiphophorus* malignant melanoma. *EMBO J.* **11**, 4239-4246
- Wittbrodt, J., Adam, D., Malitschek, B., Mäueler, W., Raulf, F., Telling, A., Robertson, S.M. & Schartl, M. (1989). Novel putative receptor tyrosine kinase encoded by the melanoma-inducing *Tu* locus in *Xiphophorus*. *Nature* **341**, 415-421

Wong, A.J., Ruppert, J.M., Eggleston, J., Hamilton, S.R., Baylin, S.B. & Vogelstein, B. (1986). Gene amplification of *c-myc* and *N-myc* in small cell carcinoma of the lung. *Science* **233**, 461-464

Woolcock, B.W., Schmidt, B.M., Kallman, K.D. & Vielkind, J.R. (1994). Differences in transcription and promoters of *Xmrk-1* and *Xmrk-2* genes suggest a role for *Xmrk-2* in pigment pattern development in the Platypus, *Xiphophorus maculatus*. *Cell Growth Diff.* **5**, 575-583

Yanisch-Perron, C., Vieira, J. & Messing, J. (1985). Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**, 103-119

Yeager, T., Stadler, W., Belair, C., Puthenveettil, J., Olopade, O. & Reznikoff, C. (1995). Increased p16 levels correlate with pRb alterations in human urothelial cells. *Cancer Res.* **55**, 493-497

Zimmerer, E.J. & Passmore, H.C. (1991). Structural and genetic properties of the Eb recombinational hotspot in the mouse. *Immunogenetics* **33**, 132-140