

TECHNICAL ADVANCE

Open Access



The Wuerzburg procedure: the tensor fasciae latae perforator is a reliable anatomical landmark to clearly identify the Hueter interval when using the minimally-invasive direct anterior approach to the hip joint

Maximilian Rudert¹, Konstantin Horas¹, Maik Hoberg¹, Andre Steinert¹, Dominik Emanuel Holzapfel², Stefan Hübner³ and Boris Michael Holzapfel^{1,4*}

Abstract

Background: The key for successful delivery in minimally-invasive hip replacement lies in the exact knowledge about the surgical anatomy. The minimally-invasive direct anterior approach to the hip joint makes it necessary to clearly identify the tensor fasciae latae muscle in order to enter the Hueter interval without damaging the lateral femoral cutaneous nerve. However, due to the inherently restricted overview in minimally-invasive surgery, this can be difficult even for experienced surgeons.

Methods and Surgical Technique: In this technical note, we demonstrate for the first time how to use the tensor fasciae latae perforator as anatomical landmark to reliably identify the tensor fasciae latae muscle in orthopaedic surgery. Such perforators are used for flaps in plastic surgery as they are constant and can be found at the lateral third of the tensor fasciae latae muscle in a direct line from the anterior superior iliac spine.

Conclusion: As demonstrated in this article, a simple knowledge transfer between surgical disciplines can minimize the complication rate associated with minimally-invasive hip replacement.

Keywords: Direct anterior approach, Hueter interval, Minimally-invasive, Hip replacement, Perforator, Anatomical landmark

Background

Several anatomical landmarks have been proposed for the reliable identification of minimally-invasive surgical approaches in total hip arthroplasty [1–3]. The correct identification of such landmarks is a *conditio sine qua non* to avoid complications related to the restricted overview of the surgical situs [4]. An elaborate description of

the surgical anatomy of the anterior approach to the hip joint was first published by Carl Hueter in 1881 [5]. In 1917 Marius Smith-Peterson was the first to propagate the use of this approach throughout the English-speaking surgical community [6]. Since then multiple other authors developed minimally-invasive techniques and continued to use those in various modifications to treat a *plethora* of different disorders around the hip joint [7–11]. All those methods have in common that they utilise the muscular interval between the sartorius and the tensor fasciae latae muscle, which is known as the Hueter interval. Using this interval poses the risk of damaging the lateral femoral cutaneous nerve as its

* Correspondence: holzapfel@orthopaedic-oncology.net

¹Department of Orthopaedic Surgery, University of Wuerzburg
Koenig-Ludwig Haus, Brettreichstr. 11, 97074 Wuerzburg, Germany

⁴Regenerative Medicine, Institute of Health and Biomedical Innovation,
Queensland University of Technology, 60 Musk Avenue, Kelvin Grove, QLD
4049 Brisbane, Australia

Full list of author information is available at the end of the article




Fig. 1 Axial cross-section through the hip joint and its surrounding soft tissues: The dotted line indicates the surgical approach with lateral incision of the superficial thigh fascia above the belly of the tensor and subsequent blunt dissection to enter the Hueter interval between the tensor fasciae latae muscle (*black asterisk*) and the sartorius muscle (*white asterisk*). This approach allows the protection of the main trunk of the lateral femoral cutaneous nerve (*yellow circle*) leaving it untouched within the fascial tunnel

main trunk usually runs along the medial border of the proximal tensor fasciae latae muscle. At this location, the nerve is covered by the superficial thigh fascia in approximately 90 % of the cases [12]. To minimise the risk for iatrogenic nerve injury, some authors advocated an

elegant technique where they incise the superficial thigh fascia as laterally as possible over the belly of the tensor fasciae latae muscle followed by blunt dissection between the muscle and the superficial fascia thereby entering the Hueter interval [13–15] (Fig. 1). By doing so, the nerve and its branches can stay untouched, embedded in the superficial thigh fascia. However, this technique makes it necessary to clearly identify the tensor fasciae latae muscle and its overlying fascia. Due to the restricted overview in minimally-invasive surgery, this can be challenging even for experienced surgeons [11]. The unambiguous identification of the muscle can furthermore be hindered by a thick subcutaneous fat layer or an inadequate positioning of the skin incision. We therefore propose the use of a constant anatomical landmark that makes it possible to clearly identify the superficial thigh fascia and the tensor muscle.

Surgical technique

Septocutaneous and musculocutaneous perforators emerge from the ascending branch of the lateral circumflex femoral artery and can be found at the lateral border of the tensor fasciae latae muscle where they penetrate the subcutaneous tissue [16] (Fig. 2). Such perforators are reported to be constant and are used in plastic surgery to create pedicles for tensor fasciae latae perforator flaps [17, 18]. Anatomical studies have shown that 100 % of the perforators emerge at the lateral border of the tensor between 6 and 15 cm distal from the anterior superior iliac spine [17]. Due to their consistency in appearance and location, these perforators are ideal anatomical landmarks for the identification of the




Fig. 2 Anatomical dissection of the antero-lateral thigh demonstrating the location of the musculocutaneous tensor fasciae latae perforator (white arrow) at the lateral border of the tensor fasciae latae muscle (TFL M). This perforator emerges from the ascending branch of the lateral circumflex femoral artery (LCFA). The profunda femoris artery, the LCFA and the descending branch of the LCFA are shown at the distal border of the iliopsoas muscle (IP M) after retraction of the sartorius muscle and the rectus femoris muscle (RF M). Note that the femoral nerve and its branches are retracted medially

superficial thigh fascia and the tensor fasciae latae muscle when using the minimally-invasive direct anterior approach for total hip replacement.

The skin incision is placed on an imaginary line between the anterior superior iliac spine and the fibular head. The incision starts two centimetres lateral and distal from the anterior superior iliac spine (Fig. 3a). The superficial subcutaneous fat layer is dissected via monopolar electro-surgical coagulation. The deep subcutaneous fat covering the thigh fasciae (black asterisk) is dissected bluntly using a sterile gauze pad to prevent iatrogenic injury of the lateral femoral cutaneous nerve branches and the tensor fasciae latae perforator (white arrow) is visualized (Fig. 3b). The tensor fasciae latae

muscle covered by the superficial thigh fascia can be clearly identified by the perforator (white arrow) that enters the subcutaneous connective tissue (Fig. 3c). The fascia is incised longitudinally at the lateral third of the tensor muscle directly medial to the perforator and separated from the muscle (white asterisk) to enter the Hueter interval (black arrow) (Fig. 3d).

Discussion

The minimally-invasive direct anterior approach was introduced in our department as the standard approach to the hip joint in 2007 [11]. We and others have demonstrated that this approach results in mid-term and long-term clinical outcomes that are comparable to outcomes




Fig. 3 The key surgical steps involved in the minimally-invasive direct anterior approach to the hip are: the correct placement of the skin incision (a), the identification of the tensor fasciae latae muscle using the perforator as anatomical landmark (b), the incision of the superficial thigh fascia as laterally as possible over the muscle (c) and the blunt dissection to enter the Hueter interval (d)




Fig. 4 Standardized placement of the skin incision in a 63 year old female patient (BMI > 40 kg/m²) undergoing hip replacement via the minimally-invasive direct anterior approach (a). Despite the presence of an enormously thick subcutaneous fat layer, the lateral border of the tensor fasciae latae muscle can be easily identified using the tensor fasciae latae perforator (white arrow) (b)

following standard procedures such as the lateral transgluteal approach. A shorter hospitalization time, reduced postoperative pain levels, a reduced fatty infiltration rate of the gluteus medius muscle and a shorter skin incision have been identified as specific advantages of this surgical approach [9–11, 19–21]. However, one disadvantage of this approach is a relatively high incidence in lesions of the lateral femoral cutaneous nerve. Damaging this purely afferent sensory nerve can result in a simple hypoaesthetic skin area at the lateral thigh, which is usually clinically insignificant, but can also lead to allodynia and eventually a significantly reduced quality of life [4]. To minimize the risk for iatrogenic injury of this nerve, the skin incision and consecutively the incision of the superficial fascia of the tensor fasciae latae muscle should be made as laterally as possible over the belly of the muscle [13–15]. A medial incision is almost inevitably associated with an injury of the main trunk of the nerve. Moreover, in such cases, a later revision and distal extension of the approach poses a very high risk of damaging branches of the femoral nerve [22, 23]. On the other hand, a too far laterally located incision increases the risk of entering the Watson-Jones interval rather than the Hueter interval. As described in our article, a standardized technique for the placement of the skin incision in combination with the use of a constant anatomical landmark such as the tensor fasciae latae perforator to identify the access to the Hueter interval can effectively prevent the above-mentioned complications. Using this technique, even highly adipose patients can be safely treated with total hip replacement via the minimally-invasive direct anterior approach (Fig. 4).

Conclusion

In the presented technical note, we advocate the use of a constant anatomical structure to clearly identify the Hueter interval and to minimize the risk for iatrogenic nerve injury. As demonstrated, a simple transfer of knowledge between surgical disciplines can significantly advance surgical techniques and eventually improve patient outcomes. The tensor fasciae latae muscle perforator that is routinely used in plastic surgery to lift tensor fasciae latae perforator flaps can serve as a reliable anatomical landmark when using the minimally-invasive direct anterior approach for total hip arthroplasty.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MR, MH, AS and BMH developed the presented technique. MR realized the importance of the perforator as an anatomical landmark for the described surgical approach. BMH wrote the paper. KH and DEH drafted the figure outline. BMH and SH were responsible for the concept and design of the manuscript. All authors approved the final version of the manuscript.

Acknowledgments

No approval of the ethics committee of the University of Wuerzburg was necessary for the description of the surgical technique outlined in this article. Written informed consent was obtained from patients whose pictures have been used for this article. This publication was funded by the German Research Foundation (DFG) in the funding program Open Access Publishing of the University of Würzburg.

Author details

¹Department of Orthopaedic Surgery, University of Wuerzburg Koenig-Ludwig Haus, Brettreichstr. 11, 97074 Wuerzburg, Germany.

²Department of Trauma, Orthopaedics, Hand and Reconstructive Surgery, Clinic Harlaching, Sanatoriumsplatz 2, 81545 Munich, Germany. ³Institute of Anatomy and Cell Biology, University of Wuerzburg, Koellikerstr. 6, 97070 Wuerzburg, Germany. ⁴Regenerative Medicine, Institute of Health and

Biomedical Innovation, Queensland University of Technology, 60 Musk Avenue, Kelvin Grove, QLD 4049 Brisbane, Australia.

Received: 16 September 2015 Accepted: 27 January 2016

Published online: 03 February 2016

References

1. Onyemaechi N, Anyanwu E, Obikili E, Ekezie J. Anatomical basis for surgical approaches to the hip. *Ann Med Health Sci Res.* 2014;4(4):487–94.
2. Hoberg M, Rudert M, Tillmann B. Minimally invasive hip arthroplasty - what must be spared? *Orthopade.* 2012;41(5):338–45.
3. Grob K, Manestar M, Ackland T, Filgueira L, Kuster MS. Potential Risk to the Superior Gluteal Nerve During the Anterior Approach to the Hip Joint: An Anatomical Study. *J Bone Joint Surg Am.* 2015;97(17):1426–31.
4. Holzapfel BM, Heinen F, Holzapfel DE, Reiners K, Noth U, Rudert M. Nerve lesions after minimally invasive total hip arthroplasty. *Orthopade.* 2012;41(5):354–64.
5. Hueter C. Fünfte Abteilung: Die Verletzung und Krankheiten des Hüftgelenks. In: *Grundriss der Chirurgie.* 2nd ed. Leipzig: FCW Vogel; 1883. p. 129–200.
6. Smith-Petersen MN. A new supra-articular subperiosteal approach to the hip joint. *J Bone Joint Surg Am.* 1917;2-15:592–5.
7. Judet J, Judet R. The use of an artificial femoral head for arthroplasty of the hip joint. *J Bone Joint Surg Br Vol.* 1950;32-B(2):166–73.
8. Kennon RE, Keggi JM, Wetmore RS, Zatorski LE, Huo MH, Keggi KJ. Total hip arthroplasty through a minimally invasive anterior surgical approach. *J Bone Joint Surg Am.* 2003;85-A Suppl 4:39–48.
9. Matta JM, Shahrdrar C, Ferguson T. Single-incision anterior approach for total hip arthroplasty on an orthopaedic table. *Clin Orthop Relat Res.* 2005;441:115–24.
10. Siguier T, Siguier M, Brumpt B. Mini-incision anterior approach does not increase dislocation rate: a study of 1037 total hip replacements. *Clin Orthop Relat Res.* 2004;426:164–73.
11. Noth U, Nedopil A, Holzapfel BM, Koppmair M, Rolf O, Goebel S, et al. Minimally invasive anterior approach. *Orthopade.* 2012;41(5):390–8.
12. Carai A, Fenu G, Sechi E, Crotti FM, Montella A. Anatomical variability of the lateral femoral cutaneous nerve: findings from a surgical series. *Clin Anat.* 2009;22(3):365–70.
13. Oinuma K, Eingartner C, Saito Y, Shiratsuchi H. Total hip arthroplasty by a minimally invasive, direct anterior approach. *Oper Orthop Traumatol.* 2007;19(3):310–26.
14. Rachbauer F. Minimally invasive total hip arthroplasty via direct anterior approach. *Orthopade.* 2005;34(11):1103–4. 1106–1108, 1110.
15. Rachbauer F, Krismer M. Minimally invasive total hip arthroplasty via direct anterior approach. *Oper Orthop Traumatol.* 2008;20(3):239–51.
16. Ishida LH, Munhoz AM, Montag E, Alves HR, Saito FL, Nakamoto HA, et al. Tensor fasciae latae perforator flap: minimizing donor-site morbidity in the treatment of trochanteric pressure sores. *Plast Reconstr Surg.* 2005;116(5):1346–52.
17. Hubmer MG, Schwaiger N, Windisch G, Feigl G, Koch H, Haas FM, et al. The vascular anatomy of the tensor fasciae latae perforator flap. *Plast Reconstr Surg.* 2009;124(1):181–9.
18. Tuinder S, Baetens T, De Haan MW, Piatkowski de Grzymala A, Booi AD, Van Der Hulst R, et al. Septocutaneous tensor fasciae latae perforator flap for breast reconstruction: radiological considerations and clinical cases. *J Plast Reconstr Aesthet Surg.* 2014;67(9):1248–56.
19. Goebel S, Steinert AF, Schillinger J, Eulert J, Broscheit J, Rudert M, et al. Reduced postoperative pain in total hip arthroplasty after minimal-invasive anterior approach. *Int Orthop.* 2012;36(3):491–8.
20. Ludemann M, Kreutner J, Haddad D, Kenn W, Rudert M, Noth U. MRI-based measurement of muscle damage after minimally invasive hip arthroplasty. *Orthopade.* 2012;41(5):346–53.
21. Reichert JC, Volkmann MR, Koppmair M, Rackwitz L, Ludemann M, Rudert M, et al. Comparative retrospective study of the direct anterior and transgluteal approaches for primary total hip arthroplasty. *Int Orthop.* 2015;39(12):2309–13.
22. Mast NH, Laude F. Revision total hip arthroplasty performed through the Hueter interval. *J Bone Joint Surg Am.* 2011;93 Suppl 2:143–8.
23. Grob K, Monahan R, Gilbey H, Yap F, Filgueira L, Kuster M. Distal extension of the direct anterior approach to the hip poses risk to neurovascular structures: an anatomical study. *J Bone Joint Surg Am.* 2015;97(2):126–32.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

