

CardioB **BEAT**

Economic Analysis of TeleHealth

GEFÖRDERT VOM



Bundesministerium
für Bildung
und Forschung

First outline and baseline data of a randomized, controlled multicenter trial to evaluate the health economic impact of home telemonitoring in chronic heart failure – CardioBBEAT

Hofmann *et al.*

METHODOLOGY

Open Access



First outline and baseline data of a randomized, controlled multicenter trial to evaluate the health economic impact of home telemonitoring in chronic heart failure – CardioBBEAT

Reiner Hofmann¹, Heinz Völler^{2,3}, Klaus Nagels¹, Dominik Bindl^{1*}, Eik Vettorazzi⁴, Ronny Dittmar^{1,5}, Walter Wohlgemuth^{1,6}, Till Neumann⁷, Stefan Störk⁸, Oliver Bruder⁹, Karl Wegscheider⁴, Eckhard Nagel¹, Eckart Fleck¹⁰ and on behalf of the CardioBBEAT Investigators

Abstract

Background: Evidence that home telemonitoring for patients with chronic heart failure (CHF) offers clinical benefit over usual care is controversial as is evidence of a health economic advantage.

Methods: Between January 2010 and June 2013, patients with a confirmed diagnosis of CHF were enrolled and randomly assigned to 2 study groups comprising usual care with and without an interactive bi-directional remote monitoring system (Motiva®). The primary endpoint in CardioBBEAT is the Incremental Cost-Effectiveness Ratio (ICER) established by the groups' difference in total cost and in the combined clinical endpoint "days alive and not in hospital nor inpatient care per potential days in study" within the follow-up of 12 months.

Results: A total of 621 predominantly male patients were enrolled, whereof 302 patients were assigned to the intervention group and 319 to the control group. Ischemic cardiomyopathy was the leading cause of heart failure. Despite randomization, subjects of the control group were more often in NYHA functional class III–IV, and exhibited peripheral edema and renal dysfunction more often. Additionally, the control and intervention groups differed in heart rhythm disorders. No differences existed regarding risk factor profile, comorbidities, echocardiographic parameters, especially left ventricular and diastolic diameter and ejection fraction, as well as functional test results, medication and quality of life. While the observed baseline differences may well be a play of chance, they are of clinical relevance. Therefore, the statistical analysis plan was extended to include adjusted analyses with respect to the baseline imbalances.

Conclusions: CardioBBEAT provides prospective outcome data on both, clinical and health economic impact of home telemonitoring in CHF. The study differs by the use of a high evidence level randomized controlled trial (RCT) design along with actual cost data obtained from health insurance companies. Its results are conducive to informed political and economic decision-making with regard to home telemonitoring solutions as an option for health care. Overall, it contributes to developing advanced health economic evaluation instruments to be deployed within the specific context of the German Health Care System.

Trial registration: ClinicalTrials.gov NCT02293252; date of registration: 10 November 2014

Keywords: Home telemonitoring, Chronic heart failure (CHF), Incremental Cost-Effectiveness Ratio (ICER), Mortality, Telemedicine, Health economics

* Correspondence: dominik.bindl@uni-bayreuth.de

¹Institute for Healthcare Management and Health Sciences, University of Bayreuth, Prieserstraße 2, 95444 Bayreuth, Germany

Full list of author information is available at the end of the article

Background

Chronic heart failure (CHF) is one of the most frequently diagnosed diseases causing disability and death in the Western hemisphere. It is characterized by a prevalence that increases with age [1]. In Germany, heart failure is the most common reason for hospitalization with about 396,000 cases in 2013 [2]. Direct medical costs related to heart failure account for 1–2 % of total health care expenditure [3].

In the majority of cases, home telemonitoring solutions in health care delivery to patients with CHF show advantages over usual care in terms of clinical outcomes. Several meta-analyses reveal that total mortality and number of hospitalizations tend to decrease, while patients' quality of life improves [4–6]. Two subsequently published trials (Telemedical Interventional Management in Heart Failure trial (TIM-HF) [7], Telemonitoring in patients with Heart Failure trial (TELE-HF) [8]) show neutral findings in general. However, the health economic impact has not been clearly demonstrated so far [9]. A meta-analysis by Klersy et al. (2011) states that the difference in costs between remote patient monitoring and usual care ranges from Euro300 to Euro1000, favoring remote patient monitoring because of a lower hospitalization rate. Thus, direct costs for hospitalization were approximated by diagnosis-related group tariffs [10]. A more detailed evaluation of efficiency and economic feasibility could help to determine cost-effectiveness and to avoid misallocation of resources [11].

The CardioBBEAT trial was designed to assess the health economic impact of a dedicated home telemonitoring system for patients with CHF based on actual costs directly obtained from patients' health care providers. The present report provides details on the outline of the study and an analysis of the study population's baseline data.

Methods

Study design

CardioBBEAT represents a randomized, controlled, open, multicenter trial with two prospective study arms. Patients were recruited at ten study sites from five areas of varying economic status in Germany: namely, Berlin, Brandenburg, Bavaria, Hamburg and North Rhine-Westphalia. This diversity allows for investigating the impact of regional differences in medical care, with general medical care predominating in rural districts compared to predominantly specialist care in urban areas. Each site was responsible for recruiting as well as following-up on their patients. Specified information about inclusion and exclusion criteria is displayed in Table 1.

The study has been conducted in accordance with the principles stated in the Declaration of Helsinki, the Good Clinical Practice (International Conference on

Table 1 Summary of inclusion and exclusion criteria at screening

Inclusion criteria	
Confirmed diagnosis of CHF based on ESC guidelines	
Symptoms corresponding to NYHA functional class II–IV	
AHA classification stages C–D	
LVEF \leq 40 %	
Age \geq 18 years	
Patient is discharged after being hospitalized for CHF within the last 12 months	
Patient is able to understand the German language	
Patient has sufficient eyesight to understand and follow the instructions communicated by Motiva®	
Patient is willing and able to use the required hardware and software and to maintain a patient diary	
Patient is residing within geographical reach of one of the ten telemonitoring centers in order to receive additional treatment if required as well as follow-up consultation	
Patient gives informed consent regarding benefits and risks related to the trial, and to sign a participation agreement for the installation of the home telemonitoring system Motiva®	
Exclusion criteria	
Myocardial infarction within the past 4 weeks	
Heart surgery or any coronary intervention within the past 8 weeks	
Cardiogenic shock within the past 4 weeks	
Intended cardiac surgery within the next 6 months or priority status on a waiting list for organ transplantation	
Severe chronic and pulmonary illness with an immediate impact on the main outcome measures	
Renal dysfunction requiring dialysis	
Dementia or other severe cognitive impairment	
Psychiatric disorders prohibiting a participation in the trial	
Patient is discharged to or living in an older persons clinic or a nursing home	
Patient is participating in another clinical trial	

AHA American Heart Association, staging of heart failure, CHF chronic heart failure, ESC European Society of Cardiology, LVEF left ventricular ejection fraction NYHA New York Heart Association, classification of heart failure

Harmonization), and national as well as local regulations. The research protocol was approved by the responsible ethics committees (Table 2), and written informed consent was obtained from all patients prior to any study-related procedures. The study is monitored by an independent external institute to ensure that every participating site abides by the study protocol and to perform external quality control of the data.

Setting

During the trial, all patients receive best medical treatment according to the current guidelines of the European Society of Cardiology (ESC) [12, 13].

Table 2 List of ethical bodies

Ethical body	Reference number
Ethik-Kommission für Forschungsfragen der Universität Bayreuth	O 1305 – HB/ID
Landesärztekammer Brandenburg, Ethik-Kommission	AS 94(a)/2009
Ethikkommission – Ethikausschuss 2 am Campus Virchow-Klinikum Berlin	EA2/084/09
Ethik-Kommission für Forschungsfragen der Universität Bayreuth	O 1305 – HB/ID
Ethik-Kommission für Forschungsfragen der Universität Bayreuth	O 1305 – HB/ID
Medizinische Fakultät der Universität Duisburg-Essen, Ethik-Kommission	10-4536
Ethik-Kommission der Bayerischen Landesärztekammer	7/11014
Universität Würzburg, Ethik-Kommission bei der Medizinischen Fakultät	128/11
Ethik-Kommission der Ärztekammer Hamburg	MC-141/11
Ethik-Kommission für Forschungsfragen der Universität Bayreuth	O 1305 – GB
Ärztekammer Nordrhein, Ethikkommission	2012451

Patients in the control group only receive best medical treatment as stated above, whereas patients in the intervention group additionally receive home telemonitoring-supported care that connects them to the participating care providers by individual guideline-compliant care plans using the telemedicine-system Motiva® (Philips Medical Systems GmbH, Hamburg, Germany).

Motiva® is an interactive bi-directional home telemonitoring system that provides remote monitoring, empowers patients to manage their disease state more effectively and enables physicians to keep in contact with the patient at home on a daily basis. Patients measure their vital signs (blood pressure, heart rate, and weight) every day and Motiva® transfers the data to the relevant telemonitoring center. In doing so, signals of decompensation regarding their heart function can be detected at an early stage and counteractive measures can be taken. In addition, patients receive information via Motiva®, i.e. coaching material, evaluations, reminders, and feedback regarding their health status as well as references to potentially necessary CHF treatment adaptations. If questionnaires reveal any problems, patients receive a phone call from the study site. Additionally, patients receive standardized questionnaires related to symptoms of cardiac decompensation, hypotension or hypertension or abnormal pulse rates. A call from the telemedicine center is made if patients gain more than 2 kg within 3 days, if their systolic blood pressure exceeds 140 mmHg or is lower than 90 mmHg, or their resting heart rate exceeds 80 bpm or is lower than 50 bpm.

A secured broadband connection (Digital Subscriber Line (DSL) or Universal Mobile Telecommunications System (UMTS)) and a set-top box turn the patient's television into their center of personalized care protected by a patient-specific password. Thus, patients can transfer all information about their health status to their attending physician safely.

Motiva® is been provided by Philips Medical Systems GmbH, Hamburg, Germany. The telecommunication infrastructure to transfer patients' data is made available by T-Systems International GmbH, Frankfurt, Germany. Both are provided without any obligations that could influence the study.

Randomization

To assign patients to one of the two study arms, CardioBBEAT used a centralized stacked randomization technique. Patients at home who were managed in cardiologic practices were randomized patient-individually. Patients primarily managed by their general practitioner (GP), on the other hand, were cluster-randomized by their GP's medical practices, to minimize carry-over effects and to keep the organizational effort manageable. The results of the randomization process with regard to the patients were displayed via the study's electronic Case Report Form (eCRF) directly after inclusion into the trial. The study center in charge did inform the patients' attending physicians whether their respective patients were enrolled in the trial and to which study arm they were assigned.

Treatment patterns

After patients are discharged from inpatient care, their GPs or outpatient medical specialists will provide their ambulatory care. These physicians have access to individual patient care plans and are authorized to complete or modify them.

Subjects enrolled in the control group receive best medical treatment according to the current guidelines of the European Society of Cardiology (ESC). Subjects enrolled in the intervention group are additionally supported by the telemedicine system Motiva® installed at the patients' home usually within 2 weeks.

All trial participants maintain a patient diary and are urged to document any health disturbances at least once a week, such as hospitalizations (date of hospitalization, reason for admission, and length of stay), consultations by any physician, and change in medication or dose rate as well as adverse effects. In addition, every patient has to participate in 3 trial-specific examinations (Table 3) at their relevant study site, that take place at the time of enrollment and after 6 as well as 12 months.

Clinical outcome measures

The primary outcome measure to assess the benefit of home telemonitoring is the combined clinical endpoint “days alive and not in hospital nor inpatient care per potential days in study.” For deceased patients, the loss in lifetime is taken into account by setting the denominator to 360 days, for patients lost to follow-up, time to last contact is used. Secondary outcome measures are total mortality, number of inpatient treatments, length of stay in hospital or nursing home, functional state of health and health-related quality of life. These will be determined by the following parameters: days survived in the study, number of hospitalizations for any reason during the study (especially cardiac and heart failure-related reasons), number of days in hospital or nursing home per study month, generic (short form health survey with 36 questions using norm-based scoring (SF-36v2), World Health Organization Five, well-being index (WHO-5)) and disease-specific (Kansas City Cardiomyopathy Questionnaire, (KCCQ)) health-related quality of life as well as medical condition and capacity of each patient.

CardioBBEAT may also be able to differentiate between particular sub-groups (e.g. gender-specific, NYHA-specific, urban/rural, diabetes mellitus) while analyzing the effectiveness of the intervention. The expectation is that several patient groups can be identified which are particularly suited for home telemonitoring with regard to clinical and/or economic outcome.

Cost data

CardioBBEAT aims to reflect the impact of home telemonitoring within an actual health care setting based on originally obtained cost data subdivided into cost of intervention, cost of inpatient and outpatient care, rehabilitation, nursing, and life-saving appliances. To this end, cost data are obtained from patients’ health insurance companies and later on validated using the records of the telemonitoring centers, GPs and medical specialists as well as patients’ diaries. Health insurance data will be obtained similarly for both, patients in the intervention group as well as the control group to avoid ascertainment bias.

Table 3 Trial-specific examinations

Enrollment examination	
1.	Patient briefing
2.	Written informed consent
3.	Verification of inclusion and exclusion criteria
4.	Demography
	Sex
	Date of birth
	Marital status
	Size and weight
5.	Hemodynamic parameters
	Heart rate
	Blood pressure (systolic/diastolic)
6.	Disease-related parameters
	Medical history
	CHF (date of diagnosis, aetiology, inpatient treatments, NYHA classification, AHA stadium)
	Comorbidities
	Medication
	Care plan compilation
	6 MWD
7.	Health-related quality of life assessment
	SF-36v2 (generic quality of life questionnaire)
	WHO-5 (generic quality of life questionnaire)
	KCCQ (disease-specific quality of life questionnaire)
Follow-up and final examination	
1.	Demography
	Weight
2.	Hemodynamic parameters
	Heart rate
	Blood pressure (systolic/diastolic)
3.	Disease-related parameters
	CHF (NYHA classification, AHA stadium)
	Newly occurring comorbidities
	Medication
	Hospitalizations or admissions to a nursing home
	Patient diary monitoring and validation of AE/SAE
	6 MWD
4.	Health-related quality of life assessment
	SF-36v2 (generic quality of life questionnaire)
	WHO-5 (generic quality of life questionnaire)
	KCCQ (disease-specific quality of life questionnaire)

Table 3 Trial-specific examinations (*Continued*)

In case a patient dies during the trial
Date of death
Cause of death

6 MWD 6-minute walking distance, AE adverse event, AHA American Heart Association, staging of heart failure, ESC European Society of Cardiology, KCCQ Kansas City Cardiomyopathy Questionnaire with 23 items for measuring disease-specific domains in CHF, LVEF left ventricular ejection fraction, NYHA New York Heart Association, classification of heart failure, SAE serious adverse event, SF-36v2 short form health survey with 36 questions using norm-based scoring, WHO-5 World Health Organization Five, well-being index

All data are analyzed with appropriate statistical methods to determine the economic effectiveness of home telemonitoring for patients with CHF. Common approaches for the analysis of cost data such as *t* tests, analysis of covariance, bootstrap techniques or permutation tests will be compared regarding their feasibility, the validity of underlying assumptions and their stability and robustness in particular if missing values have to be taken into account.

Especially when analyzing cost data or determining the Incremental Cost-Effectiveness Ratio (ICER), assumptions are made regarding discount rate, utilities, projections or estimation of costs, which are based on uncertain hypotheses. To better understand the outcomes of these analyses, CardioBBEAT uses sensitivity analyses considering best-case and worst-case scenarios to demonstrate in which way the outcomes depend on these assumptions and how they affect their assessment.

Statistical analysis

The primary endpoint ICER, consisting of the group's difference in total cost and the combined clinical endpoint "days alive and not in hospital nor inpatient care per potential days in study", is calculated with confidence intervals obtained by resampling methods. The comparative conventional endpoint "event-free survival" to measure the intervention's effectiveness is evaluated using the Kaplan-Meier analysis and log rank tests. To incorporate possible repeated hospitalizations of a patient, additional analyses will be performed, e.g. comparison of quarterly data and recurrent event analysis.

Secondary outcome measures such as number of stays in hospital per quarter, health-related quality of life or time of survival are analyzed via permutation test, covariance analysis or log rank test as part of the Kaplan-Meier analysis.

Furthermore, the trial uses a cluster-randomization technique and, therefore, correlation effects can evolve due to the collective treatment of patients in a cardiology center, medical practice, or by a single study

nurse. Such effects can result in incorrect *p* values. CardioBBEAT uses frailty and multi-level models to assess if and where such correlations occur. The magnitude of these correlations will be measured and the *p* values will be rectified.

Since at planning stage neither an established statistical method to directly compare costs and ICERs nor sufficient data to estimate the variability of cost estimates was available, the sample size was determined based on literature data. With respect to clinical endpoints, the figures were rather stable and converged to a minimum of 300 patients per group. Since the primary endpoint was expected to be mainly driven by clinical events, it was assumed that the sample size will also be sufficient for the continuously distributed ICER.

Results

The study group comprised 621 patients, predominantly men. Four hundred and seventy-two (76 %) patients were treated by 449 GPs and 149 (24 %) were treated in 119 cardiologic practices. Three hundred and two patients were randomized into the intervention group and 319 patients into the control group. Ischemic cardiomyopathy was the leading cause of heart failure (59 %). Although randomly assigned, subjects of the control group were significantly more often in NYHA functional classes III or IV and exhibited peripheral edema or renal dysfunction, respectively, more frequently (Table 4). Additionally, the control and intervention group differed in heart rhythm disturbances (Table 5). No differences were detected regarding risk factor profile, comorbidities, echocardiographic parameters, especially left ventricular and diastolic diameter and ejection fraction, as well as functional test (6 MWD) results, medication and quality of life (Tables 4, 5, 6 and 7).

In comparison with recently published German trials (TIM-HF [7], Interdisciplinary Network for Heart failure study (INH) [14]) the CardioBBEAT target population was slightly younger and comprised more male patients with fewer in NYHA classes III and IV. Nevertheless, all patients were either categorized in AHA stages C or D and every fifth patient of the given cohort was diagnosed with peripheral edema; this suggests that the study population had a comparable degree of heart failure. Regarding comorbidities and risk factor profile, the study population was very similar, particularly for diabetes or renal dysfunction. Remarkably, it revealed a left bundle branch block in approximately 25 % of the population, a rhythm disorder with a known worse prognosis [15]. Furthermore, detailed information on prognostic relevant therapies was documented. Besides a high proportion of guideline-based pharmacotherapy

Table 4 Baseline characteristics of the CardioBBEAT study participants

Characteristic	All patients <i>n</i> = 621	Usual care <i>n</i> = 319 (51 %)	Monitoring <i>n</i> = 302 (49 %)	<i>p</i> value
Demographic profile				
Age (years)				
mean ± SD	63.0 ± 11.5	63.5 ± 11.4	62.5 ± 11.6	0.303
median (IQR)	65 (55–72)	65 (55–73)	64 (54–72)	
Male sex, <i>n</i> (%)	544 (88)	280 (88)	264 (87)	0.990
Living alone, <i>n</i> (%)	159 (26)	77 (24)	82 (27)	0.442
Education (years) – number of patients (% valid)				
mean ± SD	12 ± 3	12 ± 3	12 ± 3	0.803
median (IQR)	11 (10–13)	11 (10–13)	12 (10–13)	
Causes of heart failure, <i>n</i> (%)				
Ischemic CM	363 (59)	185 (58)	178 (59)	0.797
Non-ischemic CM	258 (42)	134 (42)	124 (41)	
NYHA class, <i>n</i> (%)				
II	430 (69)	209 (66)	221 (73)	0.086
III	186 (30)	108 (34)	78 (26)	
IV	5 (1)	2 (1)	3 (1)	
NYHA class III–IV, <i>n</i> (%)	191 (31)	110 (35)	81 (27)	0.048
Peripheral edema, <i>n</i> (%)	131 (21)	83 (26)	48 (16)	0.003
Comorbidities, <i>n</i> (%)				
Stroke/TIA	41 (7)	21 (7)	20 (7)	1.000
PAD	56 (9)	24 (8)	32 (11)	0.232
COPD	87 (14)	48 (15)	39 (13)	0.516
Sleep apnea	45 (7)	25 (8)	20 (7)	0.668
Renal dysfunction (GFR ≤ 60 ml/min)	148 (24)	87 (27)	61 (20)	0.048
Depression	51 (8)	24 (8)	27 (9)	0.619
Resuscitation	81 (13)	37 (12)	44 (15)	0.327
Risk factor profile				
BMI (kg/m ²) – number of patients (% valid)				
mean ± SD	28 ± 5	28 ± 5	28 ± 5	0.529
median (IQR)	28 (25–31)	28 (25–32)	27 (25–31)	
History of smoking – number/total number (%)	435/620 (70)	221/319 (69)	214/301 (71)	0.684
Diabetes mellitus, <i>n</i> (%)	219 (35)	103 (32)	116 (38)	0.131
Hypertension, <i>n</i> (%)	533 (86)	269 (84)	264 (87)	0.323

BMI body mass index, *CM* cardiomyopathy, *COPD* chronic obstructive pulmonary disease, *GFR* glomerular filtration rate, *IQR* interquartile range, *NYHA* New York Heart Association, classification of heart failure, *PAD* peripheral arterial disease, *TIA* transient ischemic attack

including beta blockers and ACE inhibitors/ARB inhibitors, the study population was treated with mineralocorticoid receptor blockers in 71 % of all cases. A cardioverter-defibrillator or a resynchronization system was implanted in two thirds of our patients (comparable to TIM-HF study [7]) with a considerable prognostic impact on primary and secondary endpoints. Table 8 compares the most important baseline characteristics with TIM-HF [7] and INH [14].

Discussion

Several recent randomized controlled trials (RCTs), including TIM-HF [7, 16] and INH [14], have proven the positive clinical effects of home telemonitoring on several groups of patients diagnosed with CHF. The meta-analyses of Clark et al. [4], Klersy et al. [5] and Inglis et al. [6] also showed its potential to improve several clinical outcomes such as quality of life. However, many results were not statistically significant

Table 5 Baseline characteristics of the CardioBBEAT study participants - Diagnostic

Characteristic	All patients <i>n</i> = 621	Usual care <i>n</i> = 319 (51 %)	Monitoring <i>n</i> = 302 (49 %)	<i>p</i> value
ECG				
Heart rate (1/min) – number of patients (% valid)	616 (99)	315 (99)	301 (100)	
mean ± SD	72 ± 14	72 ± 14	72 ± 13	0.862
median (IQR)	71 (63–81)	71 (63–82)	71 (62–80)	
Heart rhythm – number/total number (%)				0.031
Sinus rhythm	379/619 (61)	185/318 (58)	194/301 (65)	
Atrial fibrillation	78/619 (13)	46/318 (15)	32/301 (11)	
Pacemaker ECG	154/619 (25)	86/318 (27)	68/301 (23)	
Other	8/619 (1)	1/318 (0)	7/301 (2)	
Conduction disorder – number/total number (%)				
LBBB	145/543 (27)	69/272 (25)	76/271 (28)	0.543
RBBB	49/544 (9)	28/273 (10)	21/271 (8)	0.383
QRS duration (ms) – number of patients (% valid)	568 (92)	288 (90)	280 (93)	
mean ± SD	123 ± 33	125 ± 34	121 ± 32	0.151
median (IQR)	110 (100–144)	116 (100–150)	110 (100–140)	
2D echocardiography				
LVEDD (mm) – number of patients (% valid)	584 (94)	297 (93)	287 (95)	
mean (mm) ± SD	62 ± 9	62 ± 9	63 ± 9	0.580
median (IQR)	62 (57–68)	62 (57–68)	62 (57–68)	
LVEF (%) – number of patients (% valid)	619 (100)	317 (100)	302 (100)	
mean ± SD	30 ± 7	31 ± 7	30 ± 8	0.580
median (IQR)	31 (25–37)	31 (25–37)	30 (25–36)	
Mitral insufficiency – number/total number (%)				0.319
none	102/614 (17)	55/315 (18)	47/299 (16)	
mild	370/614 (60)	180/315 (57)	190/299 (64)	
moderate	121/614 (20)	70/315 (22)	51/299 (17)	
severe	21/614 (3)	10/315 (3)	11/299 (4)	
6-minute walk test – number of patients (% valid)	559 (90)	284 (89)	275 (91)	
mean (m) ± SD	375 ± 132	376 ± 132	374 ± 131	0.804
median (IQR)	400 (300–458)	404 (300–455)	400 (300–460)	

2D two-dimensional, ECG electrocardiogram, IQR interquartile range, LBBB left bundle branch block, LVEDD left ventricular end-diastolic dimension, LVEF left ventricular ejection fraction, QRS combination of three of the graphical deflections seen on a typical ECG, RBBB right bundle branch block

mostly due to the fact that effects were too small despite adequately sized studies.

Another meta-analysis by Klersy et al. [10] focused on the economic impact of remote patient monitoring. It showed that management of HF patients by remote monitoring is cost-saving due to a substantial reduction in health care resource utilization, mostly driven by a reduction in the number of HF hospitalizations. However, cost data in this meta-analysis was estimated using 3 diagnosis-related group reimbursements (minimum, median, maximum over countries) and 3 different incidence rates and their lower and upper 95 % CI (confidence interval). These facts reflect the requirement for

additional study-derived and reliable evidence based on originally obtained cost data unlike previously negotiated prices.

In CardioBBEAT, the follow-up care of patients was more diverse than expected: the number of participating practices was higher whereas the number of patients per practice was lower than expected, resulting in an incomplete use of the random blocks implemented in the eCRF and slightly unequal sample sizes between the random groups. However, slight differences in group sizes are of no concern with respect to unbiasedness of results.

At first glance, an imbalance in baseline variables stands out. But, even with perfect randomization the

Table 6 Baseline characteristics of the CardioBBEAT study participants - Therapy

Characteristic	All patients <i>n</i> = 621	Usual care <i>n</i> = 319 (51 %)	Monitoring <i>n</i> = 302 (49 %)	<i>p</i> value
Medication				
ACE inhibitor/ARB, <i>n</i> (%)	577 (93)	297 (93)	280 (93)	0.974
Beta blocker – number/total number (%)	591/620 (95)	303/319 (95)	288/301 (96)	0.826
MR antagonist, <i>n</i> (%)	439 (71)	232 (73)	207 (69)	0.291
Diuretics, <i>n</i> (%)	506 (82)	255 (80)	251 (83)	0.360
Glycosides, <i>n</i> (%)	95 (15)	50 (16)	45 (15)	0.876
Amiodarone, <i>n</i> (%)	80 (13)	48 (15)	32 (11)	0.125
Anticoagulation, <i>n</i> (%)				
Vitamin K antagonist	231 (37)	122 (38)	109 (36)	0.637
Other	42 (7)	24 (8)	18 (6)	0.538
Devices, <i>n</i> (%)				
Pacemaker	101 (16)	61 (19)	40 (13)	0.061
ICD				
with monitoring	63 (10)	31 (10)	32 (11)	0.280
without monitoring	242 (39)	134 (42)	108 (36)	
CRT-D	89 (14)	40 (13)	49 (16)	0.232

ACE angiotensin converting enzyme, ARB angiotensin receptor blocker, CRT-D cardiac resynchronization therapy combined with defibrillation ICD implantable cardioverter defibrillator, MR mineralocorticoid receptor

Table 7 Baseline characteristics of the CardioBBEAT study participants – Quality of life

Characteristic	All patients <i>n</i> = 621	Usual care <i>n</i> = 319 (51 %)	Monitoring <i>n</i> = 302 (49 %)	<i>p</i> value
SF-36v2				
Physical comp. sum – number of patients (% valid)				
mean ± SD	581 (94)	299 (94)	282 (93)	
median (IQR)	39 ± 10	39 ± 10	38 ± 10	0.458
Mental comp. sum – number of patients (% valid)				
mean ± SD	581 (94)	299 (94)	282 (93)	
median (IQR)	45 ± 13	45 ± 12	44 ± 13	0.239
Physical functioning – number of patients (% valid)				
mean ± SD	591 (95)	304 (95)	287 (95)	
median (IQR)	51 ± 27	52 ± 27	50 ± 27	0.445
WHO-5				
Score – number of patients (% valid)	586 (94)	301 (94)	285 (94)	
mean ± SD	55 ± 25	55 ± 25	54 ± 25	0.588
median (IQR)	56 (36–76)	56 (36–76)	56 (32–76)	
KCCQ				
Overall sum – number of patients (% valid)				
mean ± SD	591 (95)	305 (96)	286 (95)	
median (IQR)	59 ± 24	59 ± 24	60 ± 23	0.574
Clinical sum – number of patients (% valid)				
mean ± SD	591 (95)	305 (96)	286 (95)	
median (IQR)	63 ± 25	63 ± 26	64 ± 24	0.409
	67 (45–85)	67 (44–85)	69 (49–84)	

Comp component, IQR interquartile range, KCCQ Kansas City cardiomyopathy questionnaire with 23 items for measuring disease-specific domains in CHF, SF-36v2 short form health survey with 36 questions using norm-based scoring, sum summary, WHO-5 World Health Organization Five, well-being index

Table 8 Compared baseline characteristics of the CardioBBEAT, TIM-HF and INH study participants

Variable	CardioBBEAT		TIM-HF		INH	
	Usual care (n = 319)	Monitoring (n = 302)	Usual care (n = 356)	Monitoring (n = 354)	Usual care (n = 363)	Monitoring (n = 352)
Demographic profile						
Age (years)	63.5 ± 11.4	62.5 ± 11.6	66.9 ± 10.5	66.9 ± 10.8	69.4 ± 11.5	67.7 ± 12.8
Male sex (%)	88	87	82	81	71	71
Living alone (%)	24	27	22	21	35	30
Clinical profile						
NYHA class (%)						
I	0	0	0	0	2	3
II	65	73	51	50	62	54
III	34	26	49	50	31	40
IV	1	1	0	0	5	3
Risk factor profile						
BMI (kg/m ²)	28 ± 5	28 ± 5	28 ± 5	28 ± 5	n.a.	n.a.
Diabetes mellitus (%)	32	38	39	40	36	36
Hypertension (%)	84	87	66	68	77	72
Diagnostic						
Heart rate (1/min)	72 ± 14	72 ± 13	71 ± 13	71 ± 13	80 ± 18	80 ± 20
LVEF (%)	31 ± 7	30 ± 8	27 ± 6	27 ± 6	30 ± 8	30 ± 8
Medication (%)						
ACE inhibitor/ARB	93	93	97	94	87	89
Beta blocker	95	96	93	92	79	81
Diuretics	80	83	94	94	86	90

ACE angiotensin converting enzyme, ARB angiotensin receptor blocker, BMI body mass index, LVEF left ventricular ejection fraction, n.a. not available, NYHA New York Heart Association, classification of heart failure

expected number of statistically significant differences between baseline variables is 5 % or 2.45 of the 49 baseline comparisons in Tables 4, 5, 6 and 7, on average. In case of independence of the baseline variables, the observed number of significances will thus follow a binomial distribution with $p = 0.05$. In CardioBBEAT, 4 out of 49 baseline comparisons (8.2 %) were significant. In case of independence, 4 or more significant comparisons would occur in 23 % of the cases even in perfect randomization. The observed baseline differences could thus well be a play of chance. However, NYHA functional class, peripheral edema, heart rhythm, and renal dysfunction were clinically highly relevant variables that might bias the conclusions even if evoked by chance. Therefore, the statistical analysis plan was extended to include adjusted analyses with respect to the baseline imbalances.

Conclusions

CardioBBEAT is a RCT that adds a comprehensive cost assessment to the clinical component of the study

including actual costs generated by patients, health services and health products. The corresponding data have been obtained directly from patients' health insurances including statutory sickness funds and private insurances. This will provide more reliable information about the cost-effectiveness of home telemonitoring in CHF patients based on the actual health care setting. CardioBBEAT may also be able to differentiate between particular sub-groups (gender-specific, NYHA-specific, urban/rural, diabetes mellitus) while analyzing the effectiveness of the intervention. The expectation is that important patient groups, which are better suited for the input of telemedicine with regard to the clinical and/or economic outcome, can be identified. The study results, reflecting a guideline-compliant, highly accurate treatment of the whole CardioBBEAT study population shown above, will significantly contribute to the existing data basis on home telemonitoring in CHF. Therefore, it adds to informed political and economic decision-making within the specific context of the German Health Care System.

Individual gratitude

Study center home telemonitoring teams

German Heart Institute Berlin

Fleck Eckart, Prof. Dr.

Roser Mattias, Dr.

Gültekin Nicole

Klinik am See Rüdersdorf

Völler Heinz, Prof. Dr.

Hartwig-Zaidan Ines

University Hospital Essen

Neumann Till, Prof. Dr.

Comprehensive Heart Failure Center Würzburg

Störk Stefan, Prof. Dr.

Schmidt Maximilian, Dr.

Schupfner Elisabeth

Contilia Heart and Vascular Center Essen

Bruder Oliver, PD Dr.

Seifert Vanessa, Dr.

Reuter Vanessa

Munich Municipal Hospital Bogenhausen

Leber Alexander, PD Dr.

Landwehr Peter, Dr.

Vivantes Medical Center Berlin Neukölln

Darius Harald, Prof. Dr.

Maselli Astrid

Jewish Hospital Berlin

Graf Kristof, Prof. Dr.

Fischer Lidia

University Heart Center Hamburg-Eppendorf

Patten-Hamel Monica, PD Dr.

de Boer Imkje

Hospitals Essen Süd

Koslowski Bernd, Dr.

Technology partners

Philips Medical Systems Boeblingen GmbH

Brüge Armin, MBA, Dipl.-Ing.

Richter Wolfgang, Dr.

T-Systems International GmbH

Bruns Uta

Hauptmann Imke

Study and project management

Institute of Health Care Management and Health Sciences

Furundzija Vesna, Dr.

Ahmed-Taner Funda, Dr.

Post Monica

Belozarov Sergej, Dr.

Stolze Kirsten

Krings Peter, Dr.

Angermann Christiane, Prof. Dr.

Menhofer Dominik, Dr.

Blank Elisabeth

Waidelich Lioba, Dr.

Steffen Melanie

Antoni Diethmar, Dr.

Tomelden June

Meincke Carsten, Dr.

Bolay Miriam, Dr.

Molz Simon, Dr.

Hermes Monika

Kemper Nicole

Bui Nhat Kha, Certified Engineer, Dipl. Can. Theol.

Sarantos Melanie

Cech Martin, Dipl.-Ing.

Götze Stephan , PD Dr.

Weinkopf Katharina

Michely Beate, Dr.

Salzwedel Annett

Reichert Clemens, Dr.

Hartner Gabriele

Grosch Bernhard, Dr.

Keinhorst Jens

Kloos Patrick, Dr.

Kirchner Aenn

Pförtner Mona, Dr.

Sinning Christoph, Dr.

Kupper-Schmidt Claudia

Reintges Nicole

Goldbach Udo, Dipl.-Ing.

Westerteicher, Christoph, Dipl.-Ing.

Foth Gerd, Dipl.-Ing.

(Continued)

Nagels Klaus, Prof. Dr. Dr.	Nagel Eckhard, Prof. Dr. mult.	Wohlgemuth Walter, Prof. Dr. Dr.
Dittmar Ronny, Dr.	Hofmann Reiner	Bindl Dominik
Department of Medical Biometry and Epidemiology		
Wegscheider Karl, Prof. Dr.	Balzer Klaus	Treszl Andras, Dr.
Vettorazzi Eik		
Monitoring		
Clinical Trial Center North		
Freese Ralf, Dr.	Papavlassopoulos Martin, Dr.	Henkes Liliane, Dr.
Borregaard Saskia, Dr.		
Development of cost data set		
AOK Nordost		
Kornek Stefanie	Schönfelder Moritz	
DAK-Gesundheit		
Wobbe Stefanie		
TK		
Kettlitz Mandy	Petereit Frank	

Abbreviations

ACE: angiotensin converting enzyme; AE: adverse event; AHA: American Heart Association, staging of heart failure; ARB: angiotensin receptor blocker; BMI: body mass index; CHF: chronic heart failure; CI: confidence interval; CM: cardiomyopathy; COPD: chronic obstructive pulmonary disease; CRT-D: cardiac resynchronization therapy combined with defibrillation; DSL: Digital Subscriber Line (broadband connection); ECG: electrocardiogram; eCRF: electronic Case Report Form; ESC: European Society of Cardiology; GFR: glomerular filtration rate; GP: general practitioner; HF: heart failure; ICD: implantable cardioverter defibrillator; ICER: Incremental Cost-Effectiveness Ratio; INH: Interdisciplinary Network for Heart failure study; KCCQ: Kansas City Cardiomyopathy Questionnaire with 23 items for measuring disease-specific domains in CHF; LBBB: left bundle branch block; LVEDD: left ventricular end-diastolic dimension; LVEF: left ventricular ejection fraction; MR: mineralocorticoid receptor; MWD: minute walking distance; n.a.: not available; NYHA: New York Heart Association, classification of heart failure; PAD: peripheral arterial disease; RBBB: right bundle branch block; RCT: randomized controlled trial; SAE: serious adverse event; SF-36v2: short form health survey with 36 questions using norm-based scoring; TELE-HF: Telemonitoring in patients with Heart Failure trial; TIA: transient ischemic attack; TIM-HF: Telemedical Interventional Management in Heart Failure trial; UMTS: Universal Mobile Telecommunications System (mobile cellular system); WHO-5: World Health Organization Five, well-being index.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

RH made substantial contributions to the development, implementation and management of the study, drafted the manuscript and will perform the economic analysis and interpretation of data. HV made substantial contributions to conception and design of the study, is principal investigator and head of clinical trial, member of steering committee, revised the manuscript critically and will perform the clinical interpretation of data. KN made substantial contributions to the development and coordination of the study, is head of health economics evaluation, member of steering committee, sponsor, and revised the manuscript critically. DB made substantial contributions to the development and management of the study, drafted the manuscript, is corresponding author and will perform the economic analysis and interpretation of data. EV made substantial contributions to conception and design of the study, drafted the manuscript and will perform the biometry and statistical analysis and interpretation of data. RD made substantial contributions to conception, design, implementation and management of the study and revised the manuscript critically. WW made substantial contributions to

conception, design and implementation of the study and revised the manuscript critically. TN made substantial contributions to acquisition of data, is principal investigator and revised the manuscript critically. SS made substantial contributions to acquisition of data, is principal investigator and revised the manuscript critically. OB made substantial contributions to acquisition of data, is head of health economics evaluation, member of steering committee, sponsor, and revised the manuscript critically. EF made substantial contributions to conception and design of the study, is principal investigator and head of clinical trial, member of steering committee, revised the manuscript critically and will perform the clinical interpretation of data. All authors read and approved the final manuscript.

Funding

The CardioBBEAT trial is supported by the Federal Ministry of Education and Research in cooperation with (and managed by) the German Aerospace Center (grant number 01KX0805). The home telemonitoring system Motiva® is provided by Philips Medical Systems GmbH, the electronic infrastructure by T-Systems International GmbH.

Author details

¹Institute for Healthcare Management and Health Sciences, University of Bayreuth, Prieserstraße 2, 95444 Bayreuth, Germany. ²Rehabilitation Center for Internal Medicine, Klinik am See, Seebad 84, 15562 Rüdersdorf, Germany. ³Center of Rehabilitation Research, University of Potsdam, Am Neuen Palais 10, 14469 Potsdam, Germany. ⁴Department of Medical Biometry and Epidemiology, University Medical Center Hamburg-Eppendorf, Martinistraße 52, 20246 Hamburg, Germany. ⁵Professional Board of German Surgeons, Luisenstraße 58/59, 10117 Berlin, Germany. ⁶Radiology, University Medical Center of Regensburg, Franz-Josef-Strauß-Allee 11, 93053 Regensburg, Germany. ⁷Clinic for Cardiology, University Hospital Essen, Hufelandstraße 55, 45147 Essen, Germany. ⁸Comprehensive Heart Failure Center Würzburg and Department of Internal Medicine I, University of Würzburg, Straubmühlweg 2a, 97078 Würzburg, Germany. ⁹Contilia Heart and Vascular Center, Department of Cardiology and Angiology, Elisabeth Hospital Essen, Klara-Kopp-Weg 1, 45138 Essen, Germany. ¹⁰Department of Medicine/ Cardiology, German Heart Institute Berlin, Augustenburger Platz 1, 13353 Berlin, Germany.

Received: 26 November 2014 Accepted: 24 July 2015

Published online: 11 August 2015

References

- Cleland JGF, Swedberg K, Follath F, Komajda M, Cohen-Solal A, Aguilar JC, et al. The EuroHeart Failure survey programme – a survey on the quality of care among patients with heart failure in Europe: Part 1: patient characteristics and diagnosis. *Eur Heart J*. 2003;24:442–63.
- Federal Statistical Office – Destatis. Gesundheit: Fallpauschalenbezogene Krankenhausstatistik. https://www.destatis.de/DE/Publikationen/Thematisch/Gesundheit/Krankenhaeuser/FallpauschalenKrankenhaus2120640137004.pdf?__blob=publicationFile. 2013. Accessed 24 Sep 2014.
- Federal Statistical Office – Destatis. Gesundheitsberichterstattung des Bundes: Krankheitskosten in Mio. €. <http://www.gbe-bund.de>. 2014. Accessed 24 Sep 2014.
- Clark RA, Inglis SC, McAlister FA, Cleland JGF, Stewart S. Telemonitoring or structured telephone support programmes for patients with chronic heart failure: systematic review and meta-analysis. *BMJ*. 2007;334:942.
- Klersy C, de Silvestri A, Gabutti G, Regoli F, Auricchio A. A meta-analysis of remote monitoring of heart failure patients. *J Am Coll Cardiol*. 2009;54:1683–94.
- Inglis SC, Clark RA, McAlister FA, Stewart S, Cleland JGF. Which components of heart failure programmes are effective? A systematic review and meta-analysis of the outcomes of structured telephone support or telemonitoring as the primary component of chronic heart failure management in 8323 patients: Abridged Cochrane Review. *Eur J Heart Fail*. 2011;13:1028–40.
- Koehler F, Winkler S, Schieber M, Sechtem U, Stangl K, Böhm M, et al. Impact of remote telemedical management on mortality and hospitalizations in ambulatory patients with chronic heart failure: the Telemedical Interventional Monitoring in Heart Failure Study. *Circulation*. 2011;123:1873–80.
- Chaudry SI, Mattera JA, Curtis JP, Spertus JA, Herrin J, Zhenqiu L, et al. Telemonitoring in patients with heart failure. *N Engl J Med*. 2010;363:2301–9.
- Heinen-Kammerer T, Wiosna W, Nelles S, Rychlik R. Monitoring von Herzfunktionen mit Telemetrie. Cologne, Germany: DAHTA@DIMDI. 2006;30.
- Klersy C, de Silvestri A, Gabutti G, Raisaro A, Curti M, Regoli F, et al. Economic impact of remote patient monitoring: an integrated economic model derived from a meta-analysis of randomized controlled trials in heart failure. *Eur J Heart Fail*. 2011;13:450–9.
- von der Schulenburg JM G, Greiner W, Jost F, Klusen N, Kubin M, Leidl R, et al. German recommendations on health economic evaluation: third and updated version of the Hanover Consensus. *Value Health*. 2008;11:539–44.
- Dickstein K, Cohen-Solal A, Filippatos G, McMurray JJV, Ponikowski P, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *Eur Heart J*. 2008;29:2388–442.
- McMurray JJV, Adamopoulos S, Anker SD, Auricchio A, Böhm M, Dickstein K, et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2012. The Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2012 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association (HFA) of the ESC. *Eur Heart J*. 2012;33:1787–847.
- Angermann CE, Störk S, Gelbrich G, Faller H, Jahns R, Frantz S, et al. Mode of action and effects of standardized collaborative disease management on mortality and morbidity in patients with systolic heart failure: the Interdisciplinary Network for Heart Failure (INH) study. *Circ Heart Fail*. 2012;5:25–35.
- Dissmann R, Kamke W, Reibis R, Herbstle J, Wegscheider K, Völler H. Prognostic impact of left bundle-branch block in the early stable phase after acute myocardial infarction. *Int J Cardiol*. 2008;130:438–43.
- Koehler F, Winkler S, Schieber M, Sechtem U, Stangl K, Böhm M, et al. Telemedicine in heart failure: pre-specified and exploratory subgroup analyses from the TIM-HF trial. *Int J Cardiol*. 2012;161:143–50.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

