
H. REFERENCES

- Akopyants, N.S., Clifton, S.W., Kersulyte, D., Crabtree, J.E., Youree, B.E., Reece, C.A., Bukanov, N.O., Drazek, E.S., Roe, B.A., and Berg, D.E. (1998). Analyses of the *cag* pathogenicity island of *Helicobacter pylori*. *Mol Microbiol* **28**:37-53.
- Alm, R.A., Guerry, P., and Trust, T.J. (1993). The *Campylobacter* σ^{54} *flaB* flagellin promoter is subject to environmental regulation. *J Bacteriol* **175**:4448-4455.
- Amsler, C.D., and Matsumura, P. (1995). Chemotactic signal transduction in *Escherichia coli* and *Salmonella typhimurium*. J. A. Hoch and T. J. Silhavy ed. ASM press, Washington, D.C., pp 89-103.
- Appelmelk, B.J., Negrini, R., Moran, A., and Kuipers, E. (1997). Molecular mimicry between *Helicobacter pylori* and the host. *Trends Microbiol* **5**:70-73.
- Arora, S.K., Ritchings, B.W., Almira, E.A., Lory, S., and Ramphal, R. (1997). A transcriptional activator, FleQ, regulates mucin adhesion and flagellar gene expression in *Pseudomonas aeruginosa* in a cascade manner. *J Bacteriol* **179**:5574-5581.
- Ascencio, F., Fransson, L.A., and Wadström, T. (1993). Affinity of the gastric pathogen *Helicobacter pylori* for the N-sulphated glycosaminoglycan heparan sulphate. *J Med Microbiol* **38**:240-244.
- Aspinall, G.O., Monteiro, M.A., Pang, H., Walsh, E.J., and Moran, A.P. (1996). Lipopolysaccharide of the *Helicobacter pylori* type strain NCTC11637 (ATCC43504): structure of the O antigen chain and core oligosaccharide regions. *Biochemistry* **35**:2489-2497.
- Atherton, J.C., Cao, P., Peek Jr., R.M., Tummuru, M.K.R., Blaser, M.J., and Cover, T.L. (1995). Mosaicism in vacuolating cytotoxin alleles of *Helicobacter pylori*: association of specific *vacA* types with cytotoxin production and peptic ulceration. *J Biol Chem* **270**:17771-17777.
- Atlung, T., and Ingmer, H. (1997). H-NS: a modulator of environmentally regulated gene expression. *Mol Microbiol* **24**:7-17.
- Attey, A., Belyaeva, T., Savery, N., Hogget, J., Fujita, N., Ishihama, A., and Busby, S. (1994). Interactions between the cyclic AMP receptor protein and the alpha subunit of RNA polymerase at the *Escherichia coli* galactose operon *P1* promoter. *Nucl Acids Res* **22**:4375-4380.
- Austin, J.W., Doig, P., Stewart, M., and Trust, T.J. (1992). Structural comparison of urease and a GroEL analog from *Helicobacter pylori*. *J Bacteriol* **174**:7470-7473.
- Bajaj, V., Lucas, R.L., Hwang, C., and Lee, C.A. (1996). Co-ordinate regulation of *Salmonella typhimurium* invasion genes by environmental and regulatory factors is mediated by control of *hilA* expression. *Mol Microbiol* **22**:703-714.

- Barer, M.R., Elliott, T.S.J., Berkeley, D., Thomas, J.E., and Eastham, E.J. (1990). Intracellular vacuolization caused by the urease of *Helicobacter pylori*. *J Infect Dis* **161**:1302-1304.
- Barton, S.G., Winrow, V.R., Rampton, D.S., Crabtree, J.E., Beales, I.L., and Calam, J. (1998). Circulating antibodies to the 60-kD heat shock protein (hsp) family in patients with *Helicobacter pylori* infection. *Clin Exp Immunol* **112**:490-494.
- Bashyam, M.D., Kaushal, D., Dasgupta, S.K., and Tyagi, A.K. (1996). A study of the mycobacterial transcriptional apparatus: Identification of novel features in promoter elements. *J Bacteriol* **178**:4847-4853.
- Bauerfeind, P., Garner, R.M., and Mobley, H.L.T. (1996). Allelic exchange mutagenesis of *nixA* in *Helicobacter pylori* results in reduced nickel transport and urease activity. *Infect Immun* **64**:2877-2880.
- Beier, D., Spohn, G., Rappuoli, R., and Scarlato, V. (1997). Identification and characterization of an operon of *Helicobacter pylori* that is involved in motility and stress adaptation. *J Bacteriol* **179**:4676-4683.
- Beier, D., Spohn, G., Rappuoli, R., and Scarlato, V. (1998). Functional analysis of the *Helicobacter pylori* principal sigma subunit of RNA polymerase reveals that the spacer region is important for efficient transcription. *Mol Microbiol* **30**:121-134.
- Benaissa, M., Babin, P., Quellard, N., Pezennec, L., Cenatiempo, Y., and Fauchere, J.L. (1996). Changes in *Helicobacter pylori* ultrastructure and antigens during conversion from the bacillary to the coccoid form. *Infect Immun* **64**:2331-2335.
- de Bernard, M., Papini, E., de Filippis, V., Gottardi, E., Telford, J., Manetti, R., Fontana, A., Rappuoli, R., and Montecucco, C. (1995). Low pH activates the vacuolating toxin of *Helicobacter pylori*, which becomes acid and pepsin resistant. *J Biol Chem* **270**:23937-23940.
- Birnboim, H.C. and Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* **7**:1513-1523.
- Blaser, M.J. (1987). Gastric *Campylobacter*-like organisms, gastritis and peptic ulcer disease. *Gastroenterology* **93**:371-382.
- Blaser, M.J. (1993). *Helicobacter pylori*: microbiology of a slow bacterial infection. *Trends Microbiol* **1**:255-260.
- Bode, G., Malfertheiner, P., Lehnhardt, G., Nilius, M., and Dischuneit, H. (1993a). Ultrastructural localization of urease of *Helicobacter pylori*. *Med Microbiol Immunol* **182**:233-242.
- Bode, G., Mauch, F., and Malfertheiner, P. (1993b). The coccoid forms of *Helicobacter pylori*. Criteria for their viability. *Epidemiol Infect* **111**:483-490.
- Borén, T., Falk, P., Roth, K.A., Larson, G., and Normark, S. (1993). Attachment of *Helicobacter pylori* to human gastric epithelium mediated by blood group antigens. *Science* **262**:1892-1895.
- Brahms, G., Brahms, S., and Magasanik, B. (1995). A sequence-induced superhelical DNA segment serves as transcriptional enhancer. *J Mol Biol* **246**:35-42.
- Bucca, G., Farina, G., Puglia, A.M., and Smith, C.P. (1995). The *dnaK* operon of *Streptomyces coelicolor* encodes a novel heat shock protein which binds to the promoter region of the operon. *Mol Microbiol* **17**:663-674.

- Cao, P., McClain, M.S., Forsyth, M.H., and Cover, T.L. (1998). Extracellular release of antigenic proteins by *Helicobacter pylori*. *Infect Immun* **66**:2984-2986.
- Censini, S., Lange, C., Xiang, Z., Crabtree, J.E., Ghiara, P., Borodovsky, M., Rappuoli, R., and Covacci, A. (1996). *cag*, a pathogenicity island of *Helicobacter pylori*, encodes type I-specific and disease-associated virulence factors. *Proc Natl Acad Sci USA* **93**:14648-14653.
- Chan, W.Y., Hui, P.K., Leung, K.M., Chow, J., Kwok, F., and Ng, C.S. (1994). Coccoid forms of *Helicobacter pylori* in the human stomach. *Am J Clin Pathol* **102**:503-507.
- Christie, P.J. (1997). *Agrobacterium tumefaciens* T-complex transport apparatus: a paradigm for a new family of multifunctional transporters in eubacteria. *J Bacteriol* **179**:3085-3094.
- Claeys, D., Faller, G., Appelmelk, B.J., Negrini, R., and Kirchner, T. (1998). The gastric H⁺,K⁺-ATPase is a major autoantigen in chronic *Helicobacter pylori* gastritis with body mucosa atrophy. *Gastroenterology* **115**:340-347.
- Cosloy, S.D. and Oishi, M. (1973). Genetic transformation in *Escherichia coli* K12. *Proc Natl Acad Sci USA* **70**:84-87.
- Costa, K., Bacher, G., Allmaier, G., Dominguez-Bello, M.G., Engstrand, L., Falk, P., de Pedro, M.A., and García-del Portillo, F. (1999). The morphological transition of *Helicobacter pylori* cells from spiral to coccoid is preceded by a substantial modification of the cell wall. *J Bacteriol* **181**:3710-3715.
- Covacci, A., Censini, S., Bugnoli, M., Petracca, R., Burroni, D., Macchia, G., Massone, A., Papini, E., Xiang, Z., Figura, N., and Rappuoli, R. (1993). Molecular characterization of the 128-kDa immunodominant antigen of *Helicobacter pylori* associated with cytotoxicity and duodenal ulcer. *Proc Natl Acad Sci USA* **90**:5791-5795.
- Covacci, A., Falkow, S., Berg, D.E., and Rappuoli, R. (1997). Did the inheritance of a pathogenicity island modify the virulence of *Helicobacter pylori*? *Trends Microbiol* **5**:205-208.
- Cover, T.L., and Blaser, M.J. (1992). Purification and characterization of the vacuolating toxin from *Helicobacter pylori*. *J Biol Chem* **267**:10570-10575.
- Cover, T.L., Tummuru, M.K., Cao, P., Thompson, S.A., and Blaser, M.J. (1994). Divergence of genetic sequences for the vacuolating cytotoxin among *Helicobacter pylori* strains. *J Biol Chem* **269**:10566-10573.
- Crabtree, J.E., Figura, N., Taylor, J.D., Bugnoli, M., Armellini, D., and Tompkins, D.S. (1992). Expression of 120 kilodalton protein and cytotoxicity in *Helicobacter pylori*. *J Clin Pathol* **45**:733-734.
- Cussac, V., Ferrero, R.L., and Labigne, A. (1992). Expression of *Helicobacter pylori* urease genes in *Escherichia coli* grown under nitrogen-limiting conditions. *J Bacteriol* **174**:2466-2473.
- Dagert, M. and Ehrlich, S.D. (1979). Prolonged incubation in calcium chloride improves the competence of *Escherichia coli* cells. *Gene* **6**:23-28.
- Deckert, G., Warren, P.W., Gaasterland, T., Young, W.G., Lenox, A.L., Graham, D.E., Overbeek, R., Snead, M.A., Keller, M., Aujay, M., Huber, R., Feldman, R.A., Short, J.M., Olson, G.J., and Swanson, R.V. (1998). The complete genome of the hyperthermophilic bacterium *Aquifex aeolicus*. *Nature* **392**:353-358.

- Dick, J.D. (1990). *Helicobacter (Campylobacter) pylori*: a twist on an old disease. *Annu Rev Microbiol* **108**:70-90.
- DiRita, V.J., and Mekalanos, J.J. (1991). Periplasmic interaction between two membrane regulatory proteins, ToxR and ToxS, results in signal transduction and transcriptional activation. *Cell* **64**:29-37.
- DiRita, V.J. (1995). Regulation of virulence in *Vibrio cholerae* by the ToxR system. In: Rappuoli *et al.*, (eds.): Signal transduction and bacterial virulence. R.G. Landes Company.
- Dunn, B.E., Campbell, G.P., Perez-Perez, G.I., and Blaser, M.J. (1990). Purification and characterization of urease from *Helicobacter pylori*. *J Biol Chem* **265**:9464-9469.
- Dunn, B.E., Roop II, R.M., Sung, C.-C., Sharma, S.A., Perez-Perez, G.I., and Blaser, M.J. (1992). Identification and purification of a cpn60 heat shock protein homologue from *Helicobacter pylori*. *Infect Immun* **60**:1946-1951.
- Dunn, B.E., Vakil, N.B., Schneider, B.G., Miller, M.M., Zitzer, J.B., Peutz, T., and Phadnis, S.H. (1997). Localization of *Helicobacter pylori* urease and heat shock protein in human gastric biopsies. *Infect Immun* **65**:1181-1188.
- Eaton, K.A., and Krakowka, S. (1994). Effect of gastric pH on urease-dependent colonization of gnotobiotic piglets by *Helicobacter pylori*. *Infect Immun* **62**:3604-3607.
- Eaton, K.A., Catrenich, C.E., Makin, K.M., and Krakowka, S. (1995). Virulence of coccoid and bacillary forms of *Helicobacter pylori* in gnotobiotic piglets. *J Infect Dis* **171**:459-462.
- Eaton, K.A., Suerbaum, S., Josenhans, C., and Krakowka, S. (1996). Colonization of gnotobiotic piglets by *Helicobacter pylori* deficient in two flagellin genes. *Infect Immun* **64**:2445-2448.
- Eisenbach, M. (1996). Control of chemotaxis. *Mol Microbiol* **20**:903-910.
- van der Ende, A., Rauws, E.A., Feller, M., Mulder, C.J., Tytgat, G.N., and Dankert, J. (1996). Heterogeneous *Helicobacter pylori* isolates from members of a family with a history of peptic ulcer disease. *Gastroenterology* **111**:638-647.
- Evans, D.J., Evans, D.G., Engstrand, L., and Graham, D. (1992). Urease-associated heat shock protein of *Helicobacter pylori*. *Infect Immun* **60**:2125-2127.
- Evans, D.G., Karjalainen, T.K., Evans Jr., D.J., Graham, D.Y., and Lee, C.-H. (1993). Cloning, nucleotide sequence, and expression of a gene encoding an adhesin subunit protein of *Helicobacter pylori*. *J Bacteriol* **175**:674-683.
- Figura, N., Guglielmetti, P., Rossolini, A., Barberi, A., Cusi, G., Musmanno, R., Russi, M., and Quaranta, S. (1989). Cytotoxin production by *Campylobacter pylori* strains isolated from patients with peptic ulcers and from patients with chronic gastritis only. *J Clin Microbiol* **27**:225-226.
- Garner, J.A., Jetton, T., and Cover, T.L. (1995). Binding and internalization of *Helicobacter pylori* cytotoxin by epithelial cells. *Gastroenterology* **108** (Suppl):A99.
- Ge, Z., Hiratsuka, K., and Taylor, D.E. (1995). Nucleotide sequence and mutational analysis indicate that two *Helicobacter pylori* genes encode a P-type ATPase and a cation-binding protein associated with copper transport. *Mol Microbiol* **15**:97-106.

- Ge, Y., Old, I.G., Saint Girons, I., and Charon, N.W. (1997). Molecular characterization of a large *Borrelia burgdorferi* motility operon which is initiated by a consensus σ^{70} promoter. *J Bacteriol* **179**:2289-2299.
- Ghiara, P., Marchetti, M., Blaser, M.J., Tummuru, M.K.R., Cover, T.L., Segal, E.D., Tompkins, L.S., and Rappuoli, R. (1995). Role of the *Helicobacter pylori* virulence factors vacuolating cytotoxin, CagA, and urease in a mouse model of disease. *Infect Immun* **63**:4154-4160.
- Gilbert, J.V., Ramakrishna, J., Sunderman Jr., F.W., Wright, A., and Plaut, A.G. (1995). Protein HpN: cloning and characterization of a histidine-rich metal-binding polypeptide in *Helicobacter pylori* and *Helicobacter mustelae*. *Infect Immun* **63**:2682-2688.
- Glocker, E., Lange, C., Covacci, A., Bereswill, S., Kist, M., and Pahl, H.L. (1998). Proteins encoded by the *cag* pathogenicity island of *Helicobacter pylori* are required for NF-kappaB activation. *Infect Immun* **66**:2346-2348.
- Grandvalet, C., Servant, P., and Mazodier, P. (1997). Disruption of *hspR*, the repressor gene of the *dnaK* operon in *Streptomyces albus* G. *Mol Microbiol* **23**:77-84.
- Grandvalet, C., de Crécy-Lagard, V., and Mazodier, P. (1999). The ClpB ATPase of *Streptomyces albus* G belongs to the HspR heat shock regulon. *Mol Microbiol* **31**:521-532.
- Gross, C.A. (1996). Function and regulation of the heat shock proteins. In *Escherichia coli and Salmonella typhimurium: Cellular and Molecular Biology*. Neidhardt, F.C., Ingraham, J.L., Low, K.B., Magasanik, B., Schaechter, M., and Umbarger, H.E. (eds). Washington, D.C.: American Society for Microbiology, pp. 1382-1399.
- Gross, R., and Rappuoli, R. (1989). Pertussis toxin promoter sequences involved in modulation. *J Bacteriol* **171**:4026-4030.
- Haas, R., Meyer, T.F., and van Putten, J.P.M. (1993). Aflagellated mutants of *Helicobacter pylori* generated by genetic transformation of naturally competent strains using shuttle mutagenesis. *Mol Microbiol* **8**:753-760.
- Hacker, J., Blum-Oehler, G., Mühldorfer, I., and Tschäpe, H. (1997). Pathogenicity islands of virulent bacteria: structure, function and impact on microbial evolution. *Mol Microbiol* **23**:1089-1097.
- Hanahan, D. (1983). Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* **166**:557-580.
- Harkey, C.W., Everiss, K.D., and Peterson, K.M. (1994). The *Vibrio cholerae* toxin-coregulated-pilus gene *tcpI* encodes a homologue of methyl-accepting protein. *Infect Immun* **62**:2669-2678.
- Hendricks, J.K., and Mobley, H.L.T. (1997). *Helicobacter pylori* ABC transporter: effect of allelic exchange mutagenesis on urease activity. *J Bacteriol* **179**:5892-5902.
- Higgins, C.F., Dorman, C.J., Stirling, D.A., Waddell, L., Booth, I.R., May, G., and Bremer, E. (1988). A physiological role for DNA supercoiling in the osmotic regulation of gene expression in *S. typhimurium* and *E. coli*. *Cell* **52**:569-584.

- Homma, M., Kutsukake, K., Hasebe, M., Iino, T., and Macnab, R.M. (1990a). FlgB, FlgC, FlgF and FlgG. A family of structurally related proteins in the flagellar basal body of *Salmonella typhimurium*. *J Mol Biol* **211**:465-477.
- Homma, M., DeRosier, D.J., and Macnab, R.M. (1990b). Flagellar hook and hook-associated proteins of *Salmonella typhimurium* and their relationship to other axial components of the flagellum. *J Mol Biol* **213**:819-832.
- Huesca, M., Borgia, S., Hoffman, P., and Lingwood, C. (1996). Acidic pH changes receptor binding specificity of *Helicobacter pylori*: a binary adhesion model in which surface heat shock (stress) proteins mediate sulfatide recognition in gastric colonization. *Infect Immun* **64**:2643-2648.
- Huesca, M., Goodwin, A., Bhangwansingh, A., Hoffman, P., and Lingwood, C.A. (1998). Characterization of an acidic-pH-inducible stress protein (hsp70), a putative sulfatide binding adhesin, from *Helicobacter pylori*. *Infect Immun* **66**:4061-4067.
- Hulton, C.S.J., Seirafi, A., Hinton, J.C.D., Sidebotham, J.M., Waddell, L., Pavitt, G.D., Owen-Hughes, T., Spassky, A., Buc, H., and Higgins, C.F. (1990). Histone-like protein H1 (H-NS), DNA supercoiling and gene expression in bacteria. *Cell* **63**:631-642.
- Ilver, D., Arnqvist, A., Ögren, J., Frick, I.-M., Kersulyte, D., Incecik, E.T., Berg, D.E., Covacci, A., Engstrand, L., and Borén, T. (1998). *Helicobacter pylori* adhesin binding fucosylated histo-blood group antigens revealed by retagging. *Science* **279**:373-377.
- Jenks, P.J., Foynes, S., Ward, S.J., Constantinidou, C., Penn, C.W., and Wren, B.W. (1997). A flagellar-specific ATPase (FliI) is necessary for flagellar export in *Helicobacter pylori*. *FEMS Microbiol Lett* **152**:205-211.
- Josenhans, C., Labigne, A., and Suerbaum, S. (1995). Comparative ultrastructural and functional studies of *Helicobacter pylori* and *Helicobacter mustelae* flagellin mutants: both flagellin subunits, FlaA and FlaB, are necessary for full motility in *Helicobacter* species. *J Bacteriol* **177**:3010-3020.
- Kansau, I., Guillain, F., Thiberge, J.-M., and Labigne, A. (1996). Nickel binding and immunological properties of the C-terminal domain of the *Helicobacter pylori* GroES homologue (HspA). *Mol Microbiol* **22**:1013-1023.
- Katsukake, K., and Doi, H. (1994). Nucleotide sequence of the *flgD* gene of *Salmonella typhimurium* which is essential for flagellar hook formation. *Biochim Biophys Acta* **1218**:443-446.
- Kostrzynska, M., Betts, J.D., Austin, J.W., and Trust, T.J. (1991). Identification, characterization, and spatial localization of two flagellin species in *Helicobacter pylori* flagella. *J Bacteriol* **173**:937-946.
- Kumar, A., Malloch, R.A., Fujita, N., Smillie, D.A., Ishihama, A., and Hayward, R.S. (1993). The minus 35 - recognition region of *Escherichia coli* sigma 70 is inessential for initiation of transcription at an “extended minus 10” promoter. *J Mol Biol* **232**:406-418.
- Kustu, S., North, K., and Weiss, D. (1991). Prokaryotic transcriptional enhancers and enhancer-binding proteins. *Trends Biochem Sci* **16**:397-402.

- Labigne-Roussel, A., Courcoux, P., and Tompkins, L. (1988). Gene disruption and replacement as feasible approach for mutagenesis of *Campylobacter jejuni*. *J Bacteriol* **170**:1704-1708.
- Labigne, A., Cussac, V., and Courcoux, P. (1991). Shuttle cloning and nucleotide sequences of *Helicobacter pylori* genes responsible for urease activity. *J Bacteriol* **173**:1920-1931.
- Laemmli, U.K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**:680-685.
- Landini, P., and Volkert, M.R. (1995). RNA polymerase α subunit binding site in positively controlled promoters: a new model for RNA polymerase-promoter interaction and transcriptional activation in the *Escherichia coli ada* and *aidB* genes. *EMBO J* **14**:4329-4335.
- Leunk, R.D., Johnson, P.T., David, B.C., Kraft, W.G., and Morgan, D.R. (1988). Cytotoxic activity in broth-culture filtrates of *Campylobacter pylori*. *J Med Microbiol* **26**:93-99.
- Leying, H., Suerbaum, S., Geis, G., and Haas, R. (1992). Cloning and genetic characterization of a *Helicobacter pylori* flagellin gene. *Mol Microbiol* **6**:2863-2874.
- Lindquist, S. and Craig, E.A. (1988). The heat shock proteins. *Annu Rev Genet* **22**:631-677.
- Lingwood, C.A., Huesca, M., and Kuksis, A. (1992). The glycerolipid receptor for *Helicobacter pylori* (and exoenzyme S) is phosphatidylethanolamine. *Infect Immun* **60**:2470-2474.
- Lupetti, P., Heuser, J.E., Manetti, R., Massari, P., Lanzavecchia, S., Bellon, P.L., Dallai, R., Rappuoli, R., and Telford, J.L. (1996). Oligomeric and subunit structure of the *Helicobacter pylori* vacuolating cytotoxin. *J Cell Biol* **133**:801-807.
- Macchia, G., Massone, A., Burroni, D., Covacci, A., Censini, S., and Rappuoli, R. (1993). The Hsp60 protein of *Helicobacter pylori*: structure and immune response in patients with gastroduodenal diseases. *Mol Microbiol* **9**:645-652.
- Macnab R.B. (1995). Flagellar switch. J. A. Hoch and T. J. Silhavy ed. ASM Press, Washington, D.C., pp 181-199.
- Maxam, A.M., and Gilbert, W. (1980). Sequencing end-labeled DNA with base-specific chemical cleavages. *Methods Enzymol* **65**:499-560.
- Megraud, F. (1995). Transmission of *Helicobacter pylori*: faecal-oral versus oral-oral route. *Aliment Pharmacol Ther* **9**(Suppl. 2):85-91.
- Melchers, K., Weitzenegger, T., Buhmann, A., Steinhilber, W., Sachs, G., and Schafer, K.P. (1996). Cloning and membrane topology of a P-type ATPase from *Helicobacter pylori*. *J Biol Chem* **271**:446-457.
- Meyer, T.S., and Lamberts, B.L. (1965). Use of coomassie brilliant blue R250 for the electrophoresis of microgram quantities of parotid saliva proteins on acrylamide-gel strips. *Biochim Biophys Acta* **107**:144-145.
- Miller, J.H. (1972). Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, N. Y.
- Miller, V.L., Taylor, R.K., and Mekalanos, J.J. (1987). Cholera toxin transcriptional activator ToxR is a transmembrane DNA binding protein. *Cell* **48**:271-279.

- Mobley, H.L.T., Garner, R.E., and Bauerfeind, P. (1995). *Helicobacter pylori* nickel transport gene *nixA*; synthesis of catalytically active urease independent of growth conditions. *Mol Microbiol* **16**:97-109.
- Morales, V.M., Backman, A., and Bagdasarian, M. (1991). A series of wide-host-range low-copy-number vectors that allow direct screening for recombinants. *Gene* **97**:39-47.
- Munzenmaier, A., Lange, C., Glocker, E., Covacci, A., Moran, A., Bereswill, S., Baeuerle, P.A., Kist, M., and Pahl, H.L. (1997). A secreted/shed product of *Helicobacter pylori* activates transcription factor nuclear factor-kappa B. *J Immunol* **159**:6140-6147.
- Muotiala, A., Helander, I.M., Pyhala, L., Kosunen, T.U., and Moran, A.P. (1992). Low biological activity of *Helicobacter pylori* lipopolysaccharide. *Infect Immun* **60**:1714-1716.
- Neidhardt, F.C., and VanBogelen, R.A. (1987). The heat shock response. In *Escherichia coli and Salmonella typhimurium: Cellular and Molecular Biology*. Neidhardt, F.C., Ingraham, J.L., Low, K.B., Magasanik, B., Schaechter, M., and Umbarger, H.E. (eds). Washington, D.C.: American Society for Microbiology, pp. 1334-1345.
- Nomura, A., Stemmermann, G.N., Chyou, P.-H., Kato, I., Perez-Perez, G.I., and Blaser, M.J. (1991). *Helicobacter pylori* infection and gastric carcinoma among Japanese Americans in Hawaii. *N Engl J Med* **325**:1132-1136.
- Odenbreit, S., Wieland, B., and Haas, R. (1996). Cloning and genetic characterization of *Helicobacter pylori* catalase and construction of a catalase-deficient mutant strain. *J Bacteriol* **178**:6960-6967.
- Odenbreit, S., Till, M., Hofreuter, D., Faller, G., and Haas, R. (1999). Genetic and functional characterization of the *alpAB* gene locus essential for adhesion of *Helicobacter pylori* to human gastric tissue. *Mol Microbiol* **31**:1537-1548.
- O'Toole, P.W., Kostrzynska, M., and Trust, T.J. (1994). Non-motile mutants of *Helicobacter pylori* and *Helicobacter mustelae* defective in flagellar hook production. *Mol Microbiol* **14**:691-703.
- O'Toole, P.W., Janzon, L., Doig, P., Huang, J., Kostrzynska, M., and Trust, T. (1995). The putative neuraminylactose-binding hemagglutinin HpaA of *Helicobacter pylori* CCUG 17874 is a lipoprotein. *J Bacteriol* **177**:6049-6057.
- Papini, E., de Bernard, M., Milia, E., Bugnoli, M., Zerial, M., Rappuoli, R., and Montecucco, C. (1994). Cellular vacuoles induced by *Helicobacter pylori* originate from late endosomal compartments. *Proc Natl Acad Sci USA* **91**:9720-9724.
- Parsonnet, J., Friedman, G.D., Vandersteen, D.P., Chang, Y., Vogelman, J.H., Orentreich, N., and Sibley, R.K. (1991). *Helicobacter pylori* infection and risk of gastric carcinoma. *N Engl J Med* **325**:1127-1131.
- Phadnis, S.H., Parlow, M.H., Levy, M., Ilver, D., Caulkins, J.B., Connors, J.B., and Dunn, B.E. (1996). Surface localization of *Helicobacter pylori* urease and heat shock protein homolog requires bacterial autolysis. *Infect Immun* **64**:905-912.
- Porwollik, S., Noonan, B., and O'Toole, P.W. (1999). Molecular characterization of a flagellar export locus of *Helicobacter pylori*. *Infect Immun* **67**:2060-2070.

- Rosenshine, I., Ruschkowski, S., Stein, M., Reinscheid, D.J., Mills, S.D., and Finlay, B.B. (1996). A pathogenic bacterium triggers epithelial signals to form a functional bacterial receptor that mediates actin pseudopod formation. *EMBO J* **15**:2613-2624.
- Ross, W., Gosink, K.K., Salomon, J., Igarashi, K., Zou, C., Ishihama, A., Severinov, K., and Gourse, R.L. (1993). A third recognition element in bacterial promoters: DNA binding by the α subunit of RNA polymerase. *Science* **262**:1407-1413.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory press, Cold Spring Harbor, N. Y.
- Sanger, F., Nicklen, S., and Coulson, A.R. (1977). DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA* **74**:5463-5467.
- Sarkari, J., Pandit, N., Moxon, E.R., and Achtman, M. (1994). Variable expression of the Opc outer membrane protein in *Neisseria meningitidis* is caused by size variation of a promoter containing poly-cytidine. *Mol Microbiol* **13**:207-217.
- Scarlato, V., Prugnola, A., Aricò, B., and Rappuoli, R. (1990). Positive transcriptional feedback at the *bvg* locus controls expression of virulence factors in *Bordetella pertussis*. *Proc Natl Acad Sci* **87**:6753-6757.
- Scarlato, V., Aricò, B., Goyard, S., Ricci, S., Manetti, R., Prugnola, A., Manetti, R., Polverino-De-Laureto, P., Ullmann, A., and Rappuoli, R. (1995). A novel chromatin-forming histone H1 homologue is encoded by a dispensable and growth-regulated gene in *Bordetella pertussis*. *Mol Microbiol* **15**:871-881.
- Schmitt, W., and Haas, R. (1994). Genetic analysis of the *Helicobacter pylori* vacuolating cytotoxin: structural similarities with the IgA protease type of exported protein. *Mol Microbiol* **12**:307-319.
- Schmitz, A., Josenhans, C., and Suerbaum, S. (1997). Cloning and characterization of the *Helicobacter pylori* *flbA* gene, which codes for a membrane protein involved in coordinated expression of flagellar genes. *J Bacteriol* **179**:987-997.
- Schulz, A., and Schumann, W. (1996). *hrcA*, the first gene of the *Bacillus subtilis dnaK* operon encodes a negative regulator of class I heat shock genes. *J Bacteriol* **178**:1088-1093.
- Segal, G., and Ron, E.Z. (1993). Heat shock transcription of the *groESL* operon of *Agrobacterium tumefaciens* may involve a hairpin-loop structure. *J Bacteriol* **175**:3083-3088.
- Segal, E.D., Lange, C., Covacci, A., Tompkins, L.S., and Falkow, S. (1997). Induction of host signal transduction pathways by *Helicobacter pylori*. *Proc Natl Acad Sci USA* **94**:7595-7599.
- Smith, A.W., Chahal, B., and French, G.L. (1994). The human gastric pathogen *Helicobacter pylori* has a gene encoding an enzyme first classified as a mucinase in *Vibrio cholerae*. *Mol Microbiol* **13**:153-160.
- Smoot, D.T., Mobley, H.L.T., Chippendale, G.R., Lewison, J.F., and Resau, J.H. (1990). *Helicobacter pylori* urease activity is toxic to human gastric epithelial cells. *Infect Immun* **58**:1992-1994.

- Southern, E.M. (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* **98**:503-517.
- Spiegelhalder, C., Gerstenecker, B., Kersten, A., Schiltz, E., and Kist, M. (1993). Purification of *Helicobacter pylori* superoxide dismutase and cloning and sequencing of the gene. *Infect Immun* **61**:5315-5325.
- Studier, F.W., Rosenberg, A.H., Dunn, J.J., and Dubendorf, J.W. (1990). Use of T7 RNA polymerase to direct expression of cloned genes. *Methods Enzymol* **185**:60-89.
- Suerbaum, S., Josenhans, C., and Labigne, A. (1993). Cloning and genetic characterization of the *Helicobacter pylori* and *Helicobacter mustelae* flagellin genes and construction of *H. pylori* *flaA*- and *flaB*-negative mutants by electroporation-mediated allelic exchange. *J Bacteriol* **175**:3278-3288.
- Suerbaum, S., Thibierge, J.-M., Kansau, I., Ferrero, R.L., and Labigne, A. (1994). *Helicobacter pylori* *hspA-hspB* heat shock gene cluster: nucleotide sequence, expression, putative function and immunogenicity. *Mol Microbiol* **14**:959-974.
- Suerbaum, S. (1995). The complex flagella of gastric *Helicobacter* species. *Trends Microbiol* **3**:168-170.
- Suerbaum, S., Brauer-Steppkes, T., Labigne, A., Cameron, B., and Drilica, K. (1998). Topoisomerase I of *Helicobacter pylori*: juxtaposition with a flagellin gene and functional requirement of a zinc finger motif. *Gene* **210**:151-161.
- Tabor, S. and Richardson, C.C. (1987). DNA sequence analysis with a modified bacteriophage T7 DNA polymerase. *Proc Natl Acad Sci USA* **84**:4767-4771.
- Tan, M., Wong, B., and Engel, J.N. (1996). Transcriptional organization and regulation of the *dnaK* and *groE* operons of *Chlamydia trachomatis*. *J Bacteriol* **178**:6983-6990.
- Tang, H., Severinov, K., Goldfarb, A., and Ebright, R.H. (1995). Rapid RNA polymerase genetics: One-day, no-column preparation of reconstituted recombinant *Escherichia coli* RNA polymerase. *Proc Natl Acad Sci USA* **92**:4902-4906.
- Tang, H., Kim, Y., Severinov, K., Goldfarb, A., and Ebright, R.H. (1996). *Escherichia coli* RNA polymerase holoenzyme: rapid reconstitution from recombinant α , β , β' and σ subunits. *Methods Enzymol* **273**:130-134.
- Taylor, D.E., and Blaser, M.J. (1991). The epidemiology of *Helicobacter pylori* infection. *Epidemiol Rev* **13**:42-59.
- Tee, W., Lambert, J.R., and Dwyer, B. (1995). Cytotoxin production by *Helicobacter pylori* from patients with upper gastrointestinal tract diseases. *J Clin Microbiol* **33**:1203-1205.
- Telford, J.L., Ghiara, P., Dell'Orco, M., Comanducci, M., Burroni, D., Bugnoli, M., Tecce, M. F., Censini, S., Covacci, A., Xiang, Z., Papini, E., Montecucco, C., Parente, L., and Rappuoli, R. (1994). Gene structure of the *Helicobacter pylori* cytotoxin and evidence of its role in gastric disease. *J Exp Med* **179**:1653-1658.
- Thomas, J.E., Gibson, G.R., Darboe, M.K., Dale, A., and Weaver, L.T. (1992). Isolation of *Helicobacter pylori* from human faeces. *Lancet* **340**:1194-1195.

- Tomb, J.F., White, O., Kerlavage, A.R., Clayton, R.A., Sutton, G.G., Fleischmann, R.D., Ketchum, K.A., Klenk, H.P., Gill, S., Dougherty, B.A., Nelson, K., Quackenbush, J., Zhou, L., Kirkness, E.F., Peterson, S., Loftus, B., Richardson, D., Dodson, R., Khalak, H.G., Glodek, A., McKenney, K., Fitzgerald, L.M., Lee, N., Adams, M.D., Hickey, E.K., Berg, D.E., Gocayne, J.D., Utterback, T.R., Peterson, J.D., Kelley, J.M., Cotton, M.D., Weidman, J.M., Fujii, C., Bowman, C., Watthey, L., Wallin, E., Hayes, W.S., Borodovsky, M., Karp, P.D., Smith, H.O., Fraser, C.M. and Venter, J.C. (1997). The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* **388**:539-547.
- Trust, T.J., Doig, P., Emody, L., Kienle, Z., Wadström, T., and O'Toole, P. (1991). High-affinity binding of the basement membrane proteins collagen type-IV and laminin to the gastric pathogen *Helicobacter pylori*. *Infect Immun* **59**:4398-4404.
- Tsuda, M., Karita, M., Morshed, M.G., Okita, K., and Nakazawa, T. (1994). A urease-negative mutant of *Helicobacter pylori* constructed by allelic exchange mutagenesis lacks the ability to colonize the nude mouse stomach. *Infect Immun* **62**:3586-3589.
- Tummuru, M.K.R., Cover, T.L., and Blaser, M.J. (1993). Cloning and expression of a high-molecular-mass major antigen of *Helicobacter pylori*: evidence of linkage to cytotoxin production. *Infect Immun* **61**:1799-1809.
- Tummuru, M.K.R., Cover, T.L., and Blaser, M.J. (1994). Mutation of the cytotoxin-associated *cagA* gene does not affect the vacuolating cytotoxin activity of *Helicobacter pylori*. *Infect Immun* **62**:2609-2613.
- Tummuru, M.K.R., Sharma, S.A., and Blaser, M.J. (1995). *Helicobacter pylori* *picB*, homologue of the *Bordetella pertussis* secretion protein, is required for induction of IL-8 in gastric epithelial cells. *Mol Microbiol* **18**:867-876.
- Ueguchi, C., Kakeda, M., Yamada, H. and Mizuno, T. (1994). An analogue of the DnaJ molecular chaperone in *Escherichia coli*. *Proc Natl Acad Sci* **91**:1054-1058.
- Vanet, A., and Labigne, A. (1998). Evidence for specific secretion rather than autolysis in the release of some *Helicobacter pylori* proteins. *Infect Immun* **66**:1023-1027.
- Wang, Y., and Taylor, D.E. (1990). Chloramphenicol resistance in *Campylobacter coli*: nucleotide sequence, expression, and cloning vector construction. *Gene* **94**:23-28.
- Warren, J.R., and Marshall, B. (1983). Unidentified curved bacilli on gastric epithelium in active chronic gastritis. *Lancet* **1**:1273-1275.
- Weel, J.F., van der Hulst, R.W., Gerrits, Y., Roorda, P., Feller, M., Dankert, J., Tytgat, G.N., and van der Ende, A. (1996). The interrelationship between cytotoxin-associated gene A, vacuolating cytotoxin, and *Helicobacter pylori*-related diseases. *J Infect Dis* **173**:1171-1175.
- Weiss, A.A., Johnson, F.D., and Burns, D.L. (1993). Molecular characterization of an operon required for pertussis toxin secretion. *Proc Natl Acad Sci USA* **90**:2970-2974.
- Werner, M.H., and Burley, S.K. (1997). Architectural transcription factors: proteins that remodel DNA. *Cell* **88**:733-736.

- Wetzstein, M., Völker, U., Dedio, J., Löbau, S., Zuber, U., Schiesswohl, M., Herget, C., Hecker, M., and Schumann, W. (1992). Cloning, sequencing, and molecular analysis of the *dnaK* locus from *Bacillus subtilis*. *J Bacteriol* **174**:3300-3311.
- Wirth, H.P., Yang, M., Karita, M., and Blaser, M.J. (1996). Expression of the human cell surface glycoconjugates Lewis x and Lewis y by *Helicobacter pylori* isolates is related to *cagA* status. *Infect Immun* **64**:4598-4605.
- Wu, J., and Newton, A. (1997). Regulation of the *Caulobacter* flagellar gene hierarchy: not just for motility. *Mol Microbiol* **24**:233-239.
- Xiang, Z., Censini, S., Bayeli, P.F., Telford, J.L., Figura, N., Rappuoli, R., and Covacci, A. (1995). Analysis of expression of CagA and VacA virulence factors in 43 strains of *Helicobacter pylori* reveals that clinical isolates can be divided into two major types and that CagA is not necessary for expression of the vacuolating cytotoxin. *Infect Immun* **63**:94-98.
- Yokota, K., Hirai, Y., Haque, M., Hayashi, S., Isogai, H., Sugiyama, T., Nagamachi, E., Tsukada, Y., Fujii, N., and Oguma, K. (1994). Heat shock protein produced by *Helicobacter pylori*. *Microbiol Immunol* **38**:403-405.
- Zou, C., Fujita, N., Igarashi, K., and Ishihama, A. (1992). Mapping of the cAMP receptor protein contact site on the α subunit of *Escherichia coli* RNA polymerase. *Mol Microbiol* **6**:2599-2605.
- Zu, T., Manetti, R., Rappuoli, R., and Scarlato, V. (1996). Differential binding of BvgA to two classes of virulence genes of *Bordetella pertussis* directs promoter selectivity by RNA polymerase. *Mol Microbiol* **21**:557-565.
- Zuber, U., and Schumann, W. (1994). CIRCE, a novel heat shock element involved in regulation of heat shock operon *dnaK* of *Bacillus subtilis*. *J Bacteriol* **176**:1359-1363.