



[⁶⁸Ga]-Pentixafor PET/CT for CXCR4-Mediated Imaging of Vestibular Schwannomas

Maria Breun^{1*}, Camelia M. Monoranu^{2,3}, Almuth F. Kessler¹, Cordula Matthies¹, Mario Löhr¹, Carsten Hagemann¹, Andreas Schirbel⁴, Steven P. Rowe⁵, Martin G. Pomper⁵, Andreas K. Buck⁴, Hans-Jürgen Wester⁶, Ralf-Ingo Ernestus¹ and Constantin Lapa^{4*}

¹ Department of Neurosurgery, University Hospital Würzburg, Würzburg, Germany, ² Department of Neuropathology, University of Würzburg, Institute of Pathology, Würzburg, Germany, ³ Comprehensive Cancer Center (CCC) Mainfranken, Würzburg, Germany, ⁴ Department of Nuclear Medicine, University Hospital Würzburg, Würzburg, Germany, ⁵ Division of Nuclear Medicine and Molecular Imaging, The Russell H. Morgan Department of Radiology and Radiological Science, Johns Hopkins University School of Medicine, Baltimore, MD, United States, ⁶ Pharmaceutical Radiochemistry, Technische Universität München, Munich, Germany

OPEN ACCESS

Edited by:

Georgios S. Limouris,
National and Kapodistrian University
of Athens, Greece

Reviewed by:

Yuji Nakamoto,
Kyoto University, Japan
Luigi Aloj,
University of Cambridge,
United Kingdom

*Correspondence:

Maria Breun
breun_m@ukw.de
Constantin Lapa
lapa_c@ukw.de

Specialty section:

This article was submitted to
Cancer Imaging and Image-directed
Interventions,
a section of the journal
Frontiers in Oncology

Received: 01 April 2019

Accepted: 28 May 2019

Published: 12 June 2019

Citation:

Breun M, Monoranu CM, Kessler AF, Matthies C, Löhr M, Hagemann C, Schirbel A, Rowe SP, Pomper MG, Buck AK, Wester H-J, Ernestus R-I and Lapa C (2019) [⁶⁸Ga]-Pentixafor PET/CT for CXCR4-Mediated Imaging of Vestibular Schwannomas. *Front. Oncol.* 9:503. doi: 10.3389/fonc.2019.00503

We have recently demonstrated CXCR4 overexpression in vestibular schwannomas (VS). This study investigated the feasibility of CXCR4-directed positron emission tomography/computed tomography (PET/CT) imaging of VS using the radiolabeled chemokine ligand [⁶⁸Ga]Pentixafor.

Methods: 4 patients with 6 primarily diagnosed or pre-treated/observed VS were enrolled. All subjects underwent [⁶⁸Ga]Pentixafor PET/CT prior to surgical resection. Images were analyzed visually and semi-quantitatively for CXCR4 expression including calculation of tumor-to-background ratios (TBR). Immunohistochemistry served as standard of reference in three patients.

Results: [⁶⁸Ga]Pentixafor PET/CT was visually positive in all cases. SUV_{mean} and SUV_{max} were 3.0 ± 0.3 and 3.8 ± 0.4 and TBR_{mean} and TBR_{max} were 4.0 ± 1.4 and 5.0 ± 1.7, respectively. Histological analysis confirmed CXCR4 expression in tumors.

Conclusion: Non-invasive imaging of CXCR4 expression using [⁶⁸Ga]Pentixafor PET/CT of VS is feasible and could prove useful for *in vivo* assessment of CXCR4 expression.

Keywords: vestibular schwannoma, CXCR4, PET/CT, molecular imaging, Pentixafor

INTRODUCTION

Vestibular schwannomas (VS) are benign nerve sheath tumors that arise from Schwann cells of the vestibulocochlear nerve (1, 2). VS regularly cause hypoacusis, dizziness, and tinnitus. These tumors usually arise sporadically, however in ~5% of the cases they are associated with a rare (1:33,000) genetic disorder, neurofibromatosis type 2 (NF2). In NF2, various types of tumors including schwannomas, meningiomas, and ependymomas develop due to loss of the NF2 gene, which encodes for Merlin, a tumor suppressor protein (3, 4).

VS are the hallmark tumors of this disease. In NF2, they usually appear bilaterally, and compared to sporadic schwannomas, they grow faster and are much more adherent to the cranial nerves and

the brainstem (5–7). Accordingly, NF2-associated vestibular schwannomas are the more aggressive tumor entity. Surgery is the standard treatment in sporadic schwannoma, but not a long-lasting solution for NF2-related tumors since the disease is often associated with persistent cranial nerve deficits and high recurrence rates. Thus, efficacious systemic or non-invasive therapies would be of value for these patients.

Chemokines are important regulators of the tumor environment, which, in addition to Merlin loss in Schwann cells, is essential for tumor development in VS. C-X-C motif chemokine receptor 4 (CXCR4), a 40-kDa G protein-coupled receptor of the chemokine receptor subfamily, was initially found to regulate leukocyte trafficking (8–11). It plays an important role in the process of homing and recruitment of progenitor and immune cells, and it is integral to the development of the nervous, hematopoietic, and cardiovascular systems during embryogenesis (9, 10, 12). However, it is also involved in diverse pathological processes, including infection, autoimmune disease, and cancer (9, 13). CXCR4 overexpression has been described in more than 30 different tumor entities including breast, prostate, lung, and colon cancer, as well as in neuroblastoma and peripheral nerve sheath tumors (10, 14). Significant overexpression of CXCR4 in both sporadic as well as neurofibromatosis-associated VS was recently demonstrated (15, 16). Therefore, CXCR4 could serve as a new target for systemic therapy with specific inhibitors (e.g., AMD3100) (17, 18). CXCR4 inhibitors have already been approved for leukemia therapy and are under investigation in trials for several solid tumors (19–21).

The radiolabeled CXCR4-targeted ligand [^{68}Ga]Pentixafor was recently developed for PET imaging (22). CXCR4 expression has been demonstrated in multiple types of cancer, including adrenocortical carcinoma (23), SCLC, glioblastoma, and hematologic malignancies (24–27). This manuscript is the first report of non-invasive detection of CXCR4 expression in patients with sporadic and NF2-associated VS.

MATERIALS AND METHODS

Patients

From June to December 2017, a total of four patients with either newly diagnosed ($n = 1$) or pre-treated and observed VS ($n = 3$) underwent imaging with [^{68}Ga]-Pentixafor-PET/CT. Routine diagnosis before surgery included MRI and was available in all patients. Six tumors were VS and one was a facial nerve schwannoma. Patient characteristics regarding tumor extension and clinical impairment are given in more detail in **Table 1**. [^{68}Ga]-Pentixafor was administered on a compassionate use base in compliance with §37 of the Declaration of Helsinki and the German Medicinal Products Act, AMG §13 2b, and in accordance with the responsible regulatory body (Regierung von Oberfranken). All patients gave written, informed consent prior to imaging. Due to the retrospective nature of this study, the local institutional review board (University Hospital Würzburg, Würzburg, Germany) waived the requirement for additional approval.

Imaging and Image Analysis

All PET scans were performed on a dedicated (PET/CT) scanner (Siemens Biograph mCT 64; Siemens Medical Solutions, Erlangen, Germany). [^{68}Ga]Pentixafor PET was performed on the day prior to surgery, 60 min after i.v. injection of 88 to 163 MBq (mean: 135 ± 28 MBq) Low-dose CT scans of the brain for attenuation correction were acquired (35 mAs, 120 keV, a 512×512 matrix, 5 mm slice thickness, increment of 30 mm/s, rotation time of 0.5 s, and pitch of 0.8). All PET images were iteratively reconstructed (3 iterations, 24 subsets with resolution recovery; Gaussian filtering; 2 mm; matrix: 400×400) using corrections for attenuation, dead-time, random events, and scatter. Acquisition and data reconstruction were performed using dedicated manufacturer software (syngo MI.PET/CT; Siemens Healthineers, Erlangen, Germany).

Images were first inspected visually by a reader with expertise in the interpretation of [^{68}Ga]Pentixafor PET (C.L.). Then the axial PET image slice displaying the maximum tumor uptake was selected. Tumor regions of interest (ROIs) were defined in 2 ways. First, a standardized 10-mm circular region was placed over the area with the maximum activity. This first ROI was used to derive maximum (SUV_{max}) and mean standardized uptake values (SUV_{mean}). A normal reference brain region was defined by drawing a ROI (diameter of 25 mm) involving the contralateral cerebral hemisphere at the level of the centrum semiovale to derive tumor-to-background ratios. Additionally, another ROI (3D isocontour) was placed in the superior sagittal sinus (at the tumor level) to derive an estimate of blood pool activity (for respective tumor-to-blood pool ratios). The radiotracer concentration in the ROIs was normalized to the injected dose per kilogram of patient's body weight to derive the SUVs.

Immunohistochemistry

All tumors were histologically assessed and graded on formalin fixed and paraffin embedded tissue sections by an experienced neuropathologist (CMM) according to the 2016 criteria of the World Health Organization (2). Schwann cell origin of the tumor cells was confirmed by the positive reaction with S100 antiserum (1:200, Dako, Hamburg, Germany). To determine the proliferative activity of tumor cells, the Ki-67 labeling index was calculated after immunostaining (monoclonal, clone Ki-67, 1:50, Dako, Hamburg, Germany) by determining the number of positive nuclei among 100 tumor cells per high power field (HPF) ($\times 400$) in a total of 10 HPF per sample.

The VS sections ($3 \mu\text{m}$) were cut from formalin-fixed paraffin-embedded tissue blocks and stained with anti-CXCR4 antibody (Zytomed 503-18440, Berlin, Germany) using a 1:50 dilution in dilution buffer (DCS, Jena, Germany). CXCR4 protein expression was visualized using a poly-link secondary antibody and a peroxidase kit (Dako; DCS Innovative Diagnostic Systems, Jena, Germany). Positive signals resulted in brown staining, and counterstaining was performed with hematoxylin. All immunohistochemically stained slides were analyzed using light microscopy (Leica, Wetzlar, Germany). Negative control experiments were performed by staining

TABLE 1 | Tumor characteristics and imaging results.

No	Tumor location	Tumor extension	Antoni type	Ki 67 (%)	CXCR4 IRS	SUV _{max}	TBR _{max}	TBIR _{max}
1	VS left	T3A 16 × 8 mm	A/B	1	9	3.37	3.59	1.32
2	VS right	T4B 30 × 35 mm	A	1–2	2–6	4.13	4.17	1.59
3	VS left	T4A 23 × 23 mm				3.78	6.87	1.71
	VS right	T3B 16x12 mm		None		4.05	7.36	1.83
4	VS left	T4 26 × 32 mm	A/B	10–15	6	3.26	3.62	1.62
	VS right	T3A				3.99	4.43	1.99

IRS, immunoreactive score; SUV, standardized uptake value; TBIR, tumor to blood pool ratio; TBR, tumor to background ratio; VS, vestibular schwannoma.

low-grade astrocytomas, and positive control experiments were performed by staining placenta sections with primary and secondary antibodies.

The analysis of the stained sections was done semi-quantitatively by light microscopy according to the immunoreactive score (IRS) described by Remmele and Stegner (28). The percentage of CXCR4 positive cells was scored as follows: 0 (no positive cells), 1 (<10% positive cells), 2 (10–50% positive cells), 3 (>50–80% positive cells), 4 (>80% positive cells). Additionally, the intensity of staining was graded: 0 (no color reaction), 1 (mild reaction), 2 (moderate reaction), 3 (intense reaction). Multiplication of both scores for a given sample yielded the IRS classification: 0–1 (negative), 2–3 (mild), 4–8 (moderate), 9–12 (strongly) positive.

Statistical Analysis

Statistical analyses were performed using Graph Pad Prism 6 software (GraphPad Software, La Jolla, CA, USA). For descriptive statistics, quantitative values were expressed as mean ± standard deviation or median and range as appropriate. Comparisons of related metric measurements were performed using the two-tailed *t*-test. A *p* < 0.05 was considered to indicate statistical significance.

RESULTS

Clinical Data

Two patients (patients #3 and #4) had neurofibromatosis type 2 with bilateral VS which had been previously treated with (radio-)surgery. In the remainder, VS were unilateral with a subject with newly diagnosed (patients #2) and a single patient with known but untreated tumors (patient #1).

Two of six tumors had been previously treated with radiosurgery and one with surgery. The remaining three tumors had no prior treatment. Three tumors were progressive as described by consecutive MR imaging with a growth rate higher than the average of 2 mm per year (29, 30), one tumor was newly diagnosed and directly treated after diagnosis. Two tumors were stable regarding their extension after radiosurgery, but caused progressive hearing impairment necessitating further treatment. Patients' characteristics are displayed in **Table 1** and **Supplemental Table 1**.

Imaging Results and Analysis

All VS demonstrated enhanced [⁶⁸Ga]Pentixafor uptake (7/7, 100.0%). All lesions were visually clearly delineated from normal brain tissue and adjacent structures. Of note, the NF2 patients displayed radiotracer accumulation in all VS (**Figure 1**), irrespective of prior treatment.

SUV_{mean} and SUV_{max} were 3.0 ± 0.3 and 3.8 ± 0.4, respectively. With background SUV_{mean} of 0.8 ± 0.2, TBR_{mean} was 4.0 ± 1.4 and TBR_{max} was 5.0 ± 1.7. Blood pool activity ranged between 2.0 and 2.6 (median, 2.4; mean, 2.3 ± 0.3) and was significantly lower than VS SUV (*p* < 0.05). Individual imaging results can be found in **Table 1** and **Supplemental Table 2**.

Immunohistochemistry Analysis


Immunohistological evaluation of CXCR4 expression for comparison with imaging results was available for three patients. In all samples, CXCR4 was detectable at the cell membrane and in the cytoplasm.

Patient #1 (maximum tumor extension: 16 × 8 mm) demonstrated intense (Score: 3) membranous CXCR4 expression in 52% of cells (Score: 3) which results in an IRS of 9. Patient #2 (maximum tumor extension: 30 × 35 mm) showed areas with mild CXCR4 expression in 37.5% of schwannoma cells and areas with moderate CXCR4 expression in 55.5% of tumor cells. Accordingly, IRS of 2 for the low-expression and of 6 for the moderate-expression areas were calculated. Patient #4 (maximum tumor extension: 26 × 32 mm) presented moderate (Score: 2) cell surface CXCR4 expression in 57% of tumor cells (Score: 3) resulting in an IRS of 6 (**Figure 1**).

In this limited cohort, there was no correlation of immunohistochemistry to the Ki67 proliferation index or [⁶⁸Ga]Pentixafor PET uptake.

DISCUSSION

This is the first report of *in vivo* imaging of CXCR4 expression in humans with VS. A recent report from our group evaluating samples of these tumors had demonstrated higher chemokine receptor expression in VS as compared to healthy vestibular nerves, with higher CXCR4 expression levels trending to correlate with greater functional impairment (16). In concordance with *in vitro* data, receptor expression on the



cell surface was visualized by [⁶⁸Ga]-Pentixafor PET/CT in our cohort in all cases, even in VS as small as 8 mm.

Considering the high tumor recurrence rate and frequently debilitating functional outcomes of patients after VS resection, especially in NF2 cases, a new therapeutic approach would be of tremendous value. Given that all tumor lesions in our cohort demonstrated CXCR4-positivity, CXCR4 could be a promising target for chemokine receptor-directed therapies. We have found that treatment with CXCR4 antagonists reduces schwannoma growth in cell culture experiments (unpublished data). Given the commercial availability of specific chemokine receptor inhibitors such as AMD3100, systemic blockage of CXCR4 might be a promising approach to (NF2-associated) VS treatment.

An important pre-requisite for receptor-targeted therapy is robust expression of the target and the possibility of *in vivo* imaging to select patients who are most appropriate for the treatment. PET imaging with the radiolabeled CXCR4 ligand [⁶⁸Ga]Pentixafor has already proven its value for the non-invasive visualization of receptor expression in a number of various tumor entities (25, 31). In this pilot cohort, [⁶⁸Ga]Pentixafor PET/CT was able to detect all schwannomas with sufficient tumor-to-background and tumor-to-blood pool ratios and matched with membranous CXCR4 expression as assessed by immunohistochemistry. Thus, CXCR4-directed PET/CT might serve as a non-invasive, *in vivo* read-out for identification of potential candidates for targeted therapy. Future studies might also investigate the relationship between CXCR4 and somatostatin receptors, which have also been demonstrated to be expressed in peripheral nerve sheath tumors and might therefore represent another suitable option (32).

This pilot study has several limitations. First, only a limited number of patients could be included in the study, thus precluding any robust conclusions from this cohort. Second, histological data for comparison with imaging findings were available in only three cases and no correlation of [⁶⁸Ga]Pentixafor uptake with histological receptor expression

could reasonably be calculated because of the limited number of cases. Autoradiography was not performed.

In the future, larger studies should be carried out to fully explore the binding of [⁶⁸Ga]Pentixafor to membranous CXCR4 and to assess whether CXCR4-directed therapy is a viable option for patients with VS.

CONCLUSION

Our pilot data demonstrate the feasibility of non-invasive imaging of CXCR4 expression in VS. [⁶⁸Ga]Pentixafor PET/CT could prove to be a useful tool for *in vivo* assessment of CXCR4 expression, especially in NF2-mutated patients. Further research to elucidate the biologic implications and potential role of [⁶⁸Ga]Pentixafor PET in selecting patients for CXCR4-directed therapy is warranted.

DATA AVAILABILITY

All datasets generated for this study are included in the manuscript and/or the **Supplementary Files**.

ETHICS STATEMENT

[⁶⁸Ga]-Pentixafor was administered on a compassionate use base in compliance with §37 of the Declaration of Helsinki and the German Medicinal Products Act, AMG §13 2b, and in accordance with the responsible regulatory body (Regierung von Oberfranken). All patients gave written, informed consent prior to imaging. Due to the retrospective nature of this study, the local institutional review board waived the requirement for additional approval.

AUTHOR CONTRIBUTIONS

MB and CL participated in the design of the study. MB and CL performed the experiments and performed the data analysis and interpretation. ML, CM, and CMM provided the samples. MB

and CL coordinated the work and drafted the manuscript, with the help of and critical revision by CM, R-IE, SR, MP, ML, AB, and CH. AFK and AS helped in performing the experiments. H-JW gave technical and pharmaceutical input. All authors read and approved the final manuscript.

FUNDING

This publication was funded by the Interdisciplinary Center of Clinical Research (IZKF) and the University of Würzburg in the funding program Open Access Publishing.

REFERENCES

- Stemmer-Rachamimov AO, Louis DN, Nielsen GP, Antonescu CR, Borowsky AD, Bronson RT, et al. Comparative pathology of nerve sheath tumors in mouse models and humans. *Cancer Res.* (2004) 64:3718–24. doi: 10.1158/0008-5472.CAN-03-4079
- Louis DN, Perry A, Reifenberger G, von Deimling A, Figarella-Branger D, Cavenee WK, et al. The 2016 world health organization classification of tumors of the central nervous system: a summary. *Acta Neuropathol.* (2016) 131:803–20. doi: 10.1007/s00401-016-1545-1
- Hanemann CO. Magic but treatable? Tumours due to loss of merlin. *Brain.* (2008) 131:606–15. doi: 10.1093/brain/awm249
- Lim SH, Ardern-Holmes S, McCowage G, de Souza P. Systemic therapy in neurofibromatosis type 2. *Cancer Treat Rev.* (2014) 40:857–61. doi: 10.1016/j.ctrv.2014.05.004
- Asthagiri AR, Parry DM, Butman JA, Kim HJ, Tsilou ET, Zhuang Z, et al. Neurofibromatosis type 2. *Lancet.* (2009) 373:1974–86. doi: 10.1016/S0140-6736(09)60259-2
- Schulz A, Zoch A, Morrison H. A neuronal function of the tumor suppressor protein merlin. *Acta Neuropathol Commun.* (2014) 2:82. doi: 10.1186/PREACCEPT-181288455213 0941
- Hexter A, Jones A, Joe H, Heap L, Smith MJ, Wallace AJ, et al. Clinical and molecular predictors of mortality in neurofibromatosis 2: a UK national analysis of 1192 patients. *J Med Genet.* (2015) 52:699–705. doi: 10.1136/jmedgenet-2015-103290
- Domanska UM, Kruizinga RC, Nagengast WB, Timmer-Bosscha H, Huls G, de Vries EG, et al. A review on CXCR4/CXCL12 axis in oncology: no place to hide. *Eur J Cancer.* (2013) 49:219–30. doi: 10.1016/j.ejca.2012.05.005
- Jacobson O, Weiss ID. CXCR4 chemokine receptor overview: biology, pathology and applications in imaging and therapy. *Theranostics.* (2013) 3:1–2. doi: 10.7150/thno.5760
- Pozzobon T, Goldoni G, Viola A, Molon B. CXCR4 signaling in health and disease. *Immunol Lett.* (2016) 177:6–15. doi: 10.1016/j.imlet.2016.06.006
- Schulz A, Buttner R, Hagel C, Baader SL, Kluwe L, Salamon J, et al. The importance of nerve microenvironment for schwannoma development. *Acta Neuropathol.* (2016) 132:289–307. doi: 10.1007/s00401-016-1583-8
- Zlotnik A, Burkhardt AM, Homey B. Homeostatic chemokine receptors and organ-specific metastasis. *Nat Rev Immunol.* (2011) 11:597–606. doi: 10.1038/nri3049
- Bar-Shavit R, Maoz M, Kancharla A, Nag JK, Agranovich D, Grisar-Granovsky S, et al. G Protein-coupled receptors in cancer. *Int J Mol Sci.* (2016) 17:E1320. doi: 10.3390/ijms17081320
- Zhao H, Guo L, Zhao H, Zhao J, Weng H, Zhao B. CXCR4 over-expression and survival in cancer: a system review and meta-analysis. *Oncotarget.* (2015) 6:5022–40. doi: 10.18632/oncotarget.3217
- Held-Feindt J, Rehmke B, Mentlein R, Hattermann K, Knerlich F, Hugo HH, et al. Overexpression of CXCL16 and its receptor CXCR6/Bonzo promotes growth of human schwannomas. *Glia.* (2008) 56:764–74. doi: 10.1002/glia.20651
- Breun M, Schwerdtfeger A, Martellotta DD, Kessler AF, Perez JM, Monoranu CM, et al. CXCR4: a new player in vestibular schwannoma pathogenesis. *Oncotarget.* (2018) 9:9940–50. doi: 10.18632/oncotarget.24119
- De Clercq E. The bicyclam AMD3100 story. *Nat Rev Drug Discov.* (2003) 2:581–7. doi: 10.1038/nrd1134
- Debnath B, Xu S, Grande F, Garofalo A, Neamati N. Small molecule inhibitors of CXCR4. *Theranostics.* (2013) 3:47–75. doi: 10.7150/thno.5376
- Liu T, Li X, You S, Bhuyan SS, Dong L. Effectiveness of AMD3100 in treatment of leukemia and solid tumors: from original discovery to use in current clinical practice. *Exp Hematol Oncol.* (2015) 5:19. doi: 10.1186/s40164-016-0050-5
- Chittasupho C, Anuchapreeda S, Sarisuta N. CXCR4 targeted dendrimer for anti-cancer drug delivery and breast cancer cell migration inhibition. *Eur J Pharm Biopharm.* (2017) 119:310–21. doi: 10.1016/j.ejpb.2017.07.003
- Gravina GL, Mancini A, Colapietro A, Vitale F, Vetuschci A, Pompili S, et al. The novel CXCR4 antagonist, PRX177561, reduces tumor cell proliferation and accelerates cancer stem cell differentiation in glioblastoma preclinical models. *Tumour Biol.* (2017) 39:1010428317695528. doi: 10.1177/1010428317695528
- Demmer O, Gourni E, Schumacher U, Kessler H, Wester HJ. PET imaging of CXCR4 receptors in cancer by a new optimized ligand. *ChemMedChem.* (2011) 6:1789–91. doi: 10.1002/cmdc.201100320
- Bluemel C, Hahner S, Heinze B, Fassnacht M, Kroiss M, Bley TA, et al. Investigating the chemokine receptor 4 as potential theranostic target in adrenocortical cancer patients. *Clin Nucl Med.* (2017) 42:e29–e34. doi: 10.1097/RLU.0000000000001435
- Philipp-Abbrederis K, Herrmann K, Knop S, Schottelius M, Eiber M, Luckerath K, et al. *In vivo* molecular imaging of chemokine receptor CXCR4 expression in patients with advanced multiple myeloma. *EMBO Mol Med.* (2015) 7:477–87. doi: 10.15252/emmm.201404698
- Lapa C, Luckerath K, Kleinlein I, Monoranu CM, Linsenmann T, Kessler AF, et al. (68)Ga-Pentixafor-PET/CT for imaging of chemokine receptor 4 expression in glioblastoma. *Theranostics.* (2016) 6:428–34. doi: 10.7150/thno.13986
- Lapa C, Schreder M, Schirbel A, Samnick S, Kortum KM, Herrmann K, et al. [(68)Ga]Pentixafor-PET/CT for imaging of chemokine receptor CXCR4 expression in multiple myeloma - Comparison to [(18)F]FDG and laboratory values. *Theranostics.* (2017) 7:205–12. doi: 10.7150/thno.16576
- Werner RA, Weich A, Higuchi T, Schmid JS, Schirbel A, Lassmann M, et al. Imaging of chemokine receptor 4 expression in neuroendocrine tumors - a triple tracer comparative approach. *Theranostics.* (2017) 7:1489–98. doi: 10.7150/thno.18754
- Kaemmerer D, Peter L, Lupp A, Schulz S, Sanger J, Baum RP, et al. Comparing of IRS and Her2 as immunohistochemical scoring schemes in gastroenteropancreatic neuroendocrine tumors. *Int J Clin Exp Pathol.* (2012) 5:187–94.

ACKNOWLEDGMENTS

We are very grateful to Siglinde Kühnel and Elisabeth Karl for excellent technical assistance and to the Interdisciplinary Center of Clinical Research (IZKF) for funding.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fonc.2019.00503/full#supplementary-material>

29. Irving RM, Harada T, Moffat DA, Hardy DG, Whittaker JL, Xuereb JH, et al. Somatic neurofibromatosis type 2 gene mutations and growth characteristics in vestibular schwannoma. *Am J Otol.* (1997) 18:754–60.
30. Sughrue ME, Kane AJ, Kaur R, Barry JJ, Rutkowski MJ, Pitts LH, et al. A prospective study of hearing preservation in untreated vestibular schwannomas. *J Neurosurg.* (2011) 114:381–5. doi: 10.3171/2010.4.JNS091962
31. Lapa C, Luckerath K, Rudelius M, Schmid JS, Schoene A, Schirbel A, et al. [68Ga]Pentixafor-PET/CT for imaging of chemokine receptor 4 expression in small cell lung cancer—initial experience. *Oncotarget.* (2016) 7:9288–95. doi: 10.18632/oncotarget.7063
32. Mawrin C, Schulz S, Hellwig-Patyk A, Kirches E, Roessner A, Lendeckel U, et al. Expression and function of somatostatin receptors in peripheral nerve sheath tumors. *J Neuropathol Exp Neurol.* (2005) 64:1080–8. doi: 10.1097/01.jnen.0000190065.36182.25

Conflict of Interest Statement: H-JW is founder and shareholder of Scintomics.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Breun, Monoranu, Kessler, Matthies, Löhr, Hagemann, Schirbel, Rowe, Pomper, Buck, Wester, Ernestus and Lapa. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.