

7. Literaturverzeichnis

- Ahrendt, H.; Hunziker, J.C.; Weber, K. (1978): Age and degree of metamorphism and time of Nappe emplacement along the Southern Margin of the Damara Orogen/ Namibia (SW-Africa).- *Geol. Rdsch.*, **67**, 2, 719-742.
- Ahrendt, H.; Behr, H.-J.; Clauer, N.; Hunziker, J.C.; Porada, H.; Weber, K. (1983a): The northern branch: Depositional development and timing of the structural and metamorphic evolution within the framework of the Damara Orogen.- In „Intracontinental fold belts“, H. Martin & F.W. Ender, eds., Springer-Verlag, Berlin-Heidelberg, 723-743.
- Ahrendt, H.; Behr, H.J.; Clauer, N.; Porada, H.; Weber, K. (1983b): K/Ar age determinations of the northern Damara branch and their implications for the structural and metamorphic evolution of the Damara Orogen, South West Africa/ Namibia.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 299-306.
- Albee, A.L. (1972): Metamorphism of pelitic rocks: reaction relations of chloritoid und staurolite.- *Bull. Geol. Soc. Am.*, **83**, 3249-3268.
- Attoh, K. (1998): High-pressure granulite facies metamorphism in the Pan-African Dahomeyide Orogen, West Africa.- *J. Geol.*, **106**, 235-246.
- Babinski, M.; Chemale, F.; Hartmann, L.A.; Van Schmus, W.R.; da Silva, L.C. (1996): Juvenile accretion at 750 - 700 Ma in southern Brazil.- *Geology*, **24**, 5, 439-442.
- Badenhorst, F.P. (1988): The lithostratigraphy of the Chuos mixtite in part of the southern Central Zone of the Damara Orogen, South West Afrika.- *Communs Geol. Surv. S.W. Africa/ Namibia*, **4**, 103-110.
- Barnes, S.J.; Sawyer, E.W. (1980): An alternative model for the Damara mobile belt: Ocean crust subduktion and continental convergence.- *Precamb. Res.*, **13**, 297-336.
- Berman, R.G. (1988): Internally-consistant thermodynamic data for minerals in the system Na₂O-K₂O-CaO-MgO-FeO-Fe₂O₃-Al₂O₃-SiO₂-TiO₂-H₂O-CO₂.- *J. Petrol.*, **29**, 445-522.
- Berman, R.G. (1990): Mixing properties of Ca-Mg-Fe-Mn garnets.- *Am. Mineral.*, **75**, 328-344.
- Berman, R.G. (1991): Thermobarometry using multi-equilibrium calculations: a new technique, with petrological applications.- *Can. Mineral.*, **29**, 833-855.
- Bhatia, M.R. (1983): Plate tectonics and geochemical composition of sandstones.- *J. Geol.*, **91**, 611-627.
- Bhatia, M. R.; Crook, K.A.W. (1986): Trace element characteristics of greywackes and tectonic discrimination of sedimentary basins.- *Contrib. Mineral. Petrol.*, **92**, 181-193.
- Bhattacharya, A.; Mazumar, A.C.; Sen, S.K. (1988): Fe-Mg mixing in cordierite; constraints from natural data and implications for cordierite-garnet geothermometry in granulites.- *Am. Mineral.*, **73**, 338-344.
- Bhattacharya, A.; Mohanty, L.; Maji, A.; Sen, S.K.; Raith, M. (1992): Non-ideal mixing in the phlogopite-annite binary: constraints from experimental data on Mg-Fe partitioning and a reformulation of the biotite-garnet geothermometer.- *Contrib. Mineral. Petrol.*, **111**, 87-93.
- Blümel, P.; Schreyer, W. (1977): Phase relations in pelitic and psammitic gneisses of the sillimanite-potash feldspar and cordierite-potash feldspar zones in the Moldanubicum of the Lam-Bodenmais area, Bavaria.- *J. Petrol.*, **18**, 431-459.
- Bohlen, S.R.; Liotta, J.J. (1986): Barometer for garnet amphibolites and garnet granulites.- *J. Petrol.*, **27**, 1025-1034.

- Bohlen, S.R.; Montana, A.; Kerrik, D.M. (1991): Precise determination of the equilibria kyanite = sillimanite and kyanite = andalusite and a revisited triple point for Al_2SiO_5 polymorphs.- Am. Mineral., **76**, 677-680. ($530 \pm 20^\circ\text{C}$ / $4.2 \pm 0.3\text{kbar}$)
- Bucher-Nurminen, K. (1987): A recalibration of the chlorite-biotite-muscovite geobarometer.- Contrib. Mineral. Petrol., **96**, 519-522.
- Bühn, B.; Häußinger, H.; Kramm, U.; Kukla, C.; Kukla, P.A.; Stanistreet, I.G. (1994): Tectono-metamorphic patterns developed during Pan-African continental collision in the Damara Inland Belt, Namibia.- Chem. Erde, **54**, 329-354.
- Caby, R. (1994): Precambrian coesit from northern Mali: first record and implications for plate tectonics in the trans-Saharan segment of the Pan-African belt.- Eur. J. Mineral., **6**, 235-244.
- Carrington, D.P.; Harley, S.L. (1995): Partial melting and phase relations in high-grade metapelites: an experimental petrogenetic grid in the KFMASH system.- Contrib. Mineral. Petrol., **120**, 270-291.
- Castaing, C.; Triboulet, C.; Feybesse; Chèvremont (1993): Tectonometamorphic evolution of Ghana, Togo and Benin in the light of the Pan-African/ Brasiliano orogeny.- Tectonophysics, **218**, 323-342.
- Chatterjee, N.D.; Flux, S. (1986): Thermodynamic mixing properties of muscovite-paragonite crystalline solutions at high temperatures and pressures, and their geological applications.- J. Petrology., **3**, 677-693.
- Chatterjee, N.D.; Froese, E. (1975): A thermodynamic study of the pseudobinary join muscovite-paragonite in the system KAlSi_3O_8 - $\text{NaAlSi}_3\text{O}_8$ - Al_2O_3 - SiO_2 - H_2O .- Am. Mineral., **60**, 985-993.
- Chemale, F.; Hartmann, L.A.; da Silva, L.C. (1995): Stratigraphy and Tectonism of Brasiliano Cycle in Southern Brazil.- Communs Geol. Surv. Namibia, **10**, 151-166.
- Coward, M.P. (1983): The tectonic history of the Damara belt.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 409-421.
- Dalziel, I.W.D. (1991): Gondwana-Laurentia supercontinent from 1.0 to 0.55 Ga; evidence, reconstruction, implications.- GSA, **23**, 2, 17.
- Dasgupta, S.; Sengupta, P.; Guha, D.; Fukuoka, M. (1991): A refined garnet-biotite Fe-Mg exchange geothermometer and its application in amphibolites and granulites.- Contrib. Mineral. Petrol., **109**, 130-137.
- Deer, W.A.; Howie, R.A.; Zussman, J. (1991): An introduction to the rock-forming minerals.- Longman Scientific & Technical, Essex, England, 17.Auflage.
- Dingeldey, D.P. (1996): Tectonic setting of metabasites from the Pan-African Kaoko Belt, NW-Namibia.- 6. Symposium TSK, Salzburg, erw. Kurzf., 91-94.
- Dingeldey, D.P. (1997): Tectono-metamorphic evolution of the Pan-African Kaoko belt, NW-Namibia.- Dissertation, Universität Würzburg, Würzburg, 247 p.
- Dingeldey, D.P.; Dürr, S.B. (1994): Transpressionsmodell für den nördlichen Küstenast des pan-afrikanischen Damara-Orogens (Namibia).- Göttinger Arb. Geol. Paläont., **Sb 1**, 5. Symposium TSK, 234-235.
- Dingeldey, D.P.; Dürr, S.B.; Charlesworth, E.G.; Franz, L.; Okrusch, M.; Stanistreet, I.G. (1994): A geotraverse through the northern coastal branch of the Damara orogen west of Sesfontein, Namibia.- J. African Earth Sci., **19**, 4, 315-329.

- Dingeldey, D.P.; Okrusch, M.; Franz, L. (1995): Tectonometamorphic history of the Damara orogen (northern coastal branch): Metamorphic history.- Ext. Abstr. Centenniel Geocongress Vol. II, 1117-1120.
- Dingeldey, D.P.; Franz, L.; Dürr, S.B. (1996): Metamorphic evolution of the Pan-African Kaoko Belt, NW-Namibia: Evidence for lithospheric delamination.- 6. Symposium TSK, Salzburg, erw. Kurzf., 95-98.
- Dombrowski, A. (1998): Fluid evolution in scapolite-bearing shear zones of the Pan-African Kuiseb Formation in the northern Khomas Trough, Damara Orogen, central Namibia.- Dissertation, Universität Würzburg, Würzburg, 236 p.
- Downing, K.N.; Coward, M.P. (1981): The Okahandja Lineament and its significance for Damaran tectonics in Namibia.- Geol. Rdsch., **70**, 3, 972-1000.
- Dougan, T.W. (1974): Cordierite gneisses and associated lithologies of the Guri area Northwest Guayana shield, Venezuela.- Contrib. Mineral. Petrol., **46**, 169-188.
- Droop, G.T.R.; Harte, B. (1995): The effect of Mn on the phase relations of medium-grade pelites: Constraints from natural assemblages on petrogenetic grid topology.- J. Petrol., **36**, 1549-1578.
- Dürr, S.B.; Dingeldey, D.P. (1995): Tectonometamorphic history of the Damara orogen (northern coastal branch): Structural and kinematic evolution.- Ext. Abstr. Centenniel Geocongress Vol. II, 1126-1129.
- Dürr, S.B.; Dingeldey, D.P. (1996a): Gondwana-Amalgamation, gesehen vom panafrikanischen Kaoko-Gürtel (Namibia).- 6. Symposium TSK, Salzburg, erw. Kurzf., 103-106.
- Dürr, S.B.; Dingeldey, D.P. (1996b): The Kaoko belt (Namibia): Part of a late Neoproterozoic continental-scale strike-slip system.- Geology, **24**, 6, 503-506.
- Dürr, S.B.; Dingeldey, D.P.; Okrusch, M.; Franz, L. (1996): Transpression in the Panafrican Kaoko Belt of northern Namibia: structural and petrological implications.- Zbl. Geol. Paläont. I, **3/4**, 495-505.
- Dutrow, B.L.; Holdaway, M.J.; Hinton, R.W. (1986): Lithium in staurolite and its petrologic significance.- Contrib. Mineral. Petrol., **94**, 496-506.
- Ebert, H.D.; Chemale, F.; Babinski, M.; Artur, A.C.; Van Schmus, W.R. (1996): Tectonic setting and U/Pb zircon dating of the plutonic Socorro Complex in the transpressiv Rio Paraiba do Sul Shear Belt, SE Brazil.- Tectonics, **15**, 2, 688-699.
- Eugster, H.P.; Albee, A.L.; Bence, A.E.; Thompson, J.B.; Waldbaum, D.R. (1972): The two-phase region and excess mixing properties of paragonite-muscovite crystalline solutions.- J. Petrol., **13**, 147-179.
- Ferry, J.M.; Spear, F.S. (1978): Experimental calibration of the partitioning of Fe and Mg between biotite and garnet.- Contrib. Mineral. Petrol., **66**, 113-117.
- Finnemore, S.H. (1978): The geochemistry and origin of the Matchless amphibolite belt, Windhoek district, South West Africa.- In „Mineralization in metamorphic terranes“, W.J. Verwoerd, ed., Spec. Publ. Geol. Soc. S. Afr., **4**, 433-477.
- Franz, L. (1991): Geologische, geochemische und petrologische Untersuchungen an präalpidischen Metamorphiten der Inseln Kreta und des Dodekanes (Griechenland).- Dissertation, Universität Würzburg, Würzburg, 388 S.
- Franz, L.; Romer, R.L.; Dingeldey, D.P. (1999): Diachronous Pan-African granulite-facies metamorphism (650 Ma and 550 Ma) in the Kaoko belt, NW Namibia.- Eur. J. Mineral., **11**, 167-180.

- Frimmel, H.E. (1995): Metamorphic evolution of the Gariep Belt.- South Afric. J. Geol., **98**, 2, 176 - 190.
- Frimmel, H.E.; Frank, W. (1998): Neoproterozoic tectono-thermal evolution of the Gariep Belt and its basement, Namibia and South Africa.- Precamb. Res., **90**, 1-28.
- Frimmel, H.E.; Hartnady, C.J.H. (1992): Blue amphiboles and their significance for the metamorphic history of the Pan-African Gariep belt, Namibia.- J. Metamorph. Geol., **10**, 651-669.
- Frimmel, H.E.; Hartnady, C.J.H.; Koller, F. (1996a): Geochemistry and tectonic setting of magmatic units in the Pan-African Gariep belt, Namibia.- Chem. Geol., **130**, 101-121.
- Frimmel, H.E.; Klötzli, U.S.; Siegfried, P.R. (1996b): New Pb-Pb single zircon age constraints on the timing of Neoproterozoic glaciation and continental break-up in Namibia.- J. Geol., **104**, 459 - 469.
- Fuhrman, M.L.; Lindsley, D.H. (1988): Ternary-feldspar modeling and thermometry.- Am. Mineral., **73**, 201-215.
- Garcia-Casco, A.; Torres-Roldan, R.L. (1996): Disequilibrium induced by fast decompression in St-Bt-Grt-Ky-Sil-And metapelites from the Betic Belt (Southern Spain).- J. Petrol, **37**, 1207-1239.
- Gevers, T.W. (1931): An ancient tillite in South West Africa.- Trans. Geol. Soc. S. Afr., **34**, 1-17.
- Ghent, E.D. (1976): Plagioclase-garnet-Al₂SiO₅-quartz; a potential geobarometer-geothermometer.- Am. Mineral., **61**, 710-714.
- Ghent, E.D.; Stout, M.Z. (1981a): Geobarometry and geothermometry of plagioclase-biotite-garnet-muscovite assemblages.- Contrib. Mineral. Petrol., **76**, 92-97.
- Ghent, E.D.; Stout, M.Z. (1981b): Geothermometry and geobarometry of clinopyroxene-garnet-amphibolites, sillimanite zone, Mica Creek, British Columbia.- EOS, Trans. Am. Geophys. Union, **62**, 45, 1060.
- Ghent, E.D.; Stout, M.Z. (1984): TiO₂ activity in metamorphosed pelitic and basic rocks: principles and applications to metamorphism in southeastern Canadian Cordillera.- Contrib. Mineral Petrol., **86**, 248-255.
- Ghent, E.D.; Stout, M.Z.; Black, P.M.; Brothers, R.N. (1987): Chloritoid-bearing rocks associated with blueschists and eclogites, northern Caledonia.- J. Metamorph. Geol., **5**, 239-254.
- Ghildiyal, H. (1999): Structural geology and geodynamic evolution of the Panafrican Kaoko belt, NW-Namibia.- Dissertation, Universität Würzburg, Würzburg, 150 p.
- Gottschalk, M. (1997): Internally consistant thermodynamic data for rock-forming minerals in the system SiO₂-TiO₂-Al₂O₃-Fe₂O₃-CaO-MgO-FeO-K₂O-Na₂O-H₂O-CO₂.- Eur. J. Mineral., **9**, 175-223.
- Green, T.H.; Hellmann, P.L. (1982): Fe-Mg partitioning between coexisting garnet and phengite at high pressure, and comments on a garnet-phengite geothermometer.- Lithos, **15**, 253-266.
- Gresse, P.G. (1994): Strain partitioning in the southern Gariep Arc as reflected by sheath folds and stretching directions.- South Afr. J. Geol., **97**, 1, 52-61.
- Griffen, D.T.; Ribbe, P.H. (1973): The crystal chemistry of staurolite.- Am. J. Sci., **273 A**, 479-495.
- Gruner, B. (1999): P-T evolution of metapelites in the Pan-African Kaoko belt, NW Namibia.- in: „Aspects of Pan-African tectonics“, DeWall, H.; Greiling, R.O. (eds.), Schriften des Forschungszentrums Jülich, **32**, 29-34. (ISBN 3-89336-253-3)
- Gruner, B.B.; Will, T.M.; Okrusch, M. (1999): Pan-African metapelites of the Kaoko belt in Namibia: P-T history.- Conference EUG 10, March 28 - April 01 1999, Strasbourg, Terra Abstracts, **11**, 105.

- Guj, P. (1970): The Damara mobile belt in the south-western Kaokoveld, South West Africa.- Bull. Precamb. Res. Unit, Univ. Cape Town, **10**, 168p.
- Hartmann, O.; Hoffer, E.; Haak, U. (1983): Regional metamorphism in the Damara Orogen: Interaction of crustal motion and heat transfer.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 233-241.
- Hartnady, C.J. (1978): The stratigraphy and structure of the Naukluft nappe complex.- Precamb. Res. Unit, Annual Rep., **14/15**, Univ. Cape Town, 163-170.
- Hartnady, C.J.; Joubert, P.; Stowe, C. (1985): Proterozoic crustal evolution in southwestern Africa.- Episodes, **8**, 4, 236-244.
- Häußinger, H. (1990): Geochemische und petrologische Untersuchungen an metasedimentären Nebengesteinen und assoziierten Metabasiten der Sulfiderz-Vorkommen von Gorob, Damara Orogen, Namibia.- Dissertation, Universität Würzburg, Würzburg, 258S.
- Hawkesworth, C.J.; Marlow, A.G. (1983): Isotope evolution of the Damara orogenic belt:- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 397-407.
- Hawkesworth, C.J.; Gledhill, A.R.; Roddick, J.C.; Miller, R.McG.; Kröner, A. (1983): Rb-Sr and $^{40}\text{Ar}/^{39}\text{Ar}$ studies bearing on models for the thermal evolution of the Damara belt, Namibia.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 323-338.
- Hawkesworth, C.J.; Menzies, A.L.; van Calsteren, D. (1986): Chemical and tectonic evolution of the Damara Belt, Namibia.- In „Collision tectonics“, M.P. Coward; C. Ries-Alison, eds., Spec. Publ. Geol. Soc. of London, London, 305-319.
- Hedberg, R.M. (1979): Stratigraphy of the Ovamboland Basin, South West Afrika.- Bull. Precamb. Res. Unit, **24**, Univ. Cape Town, 325pp.
- Henry, G.; Stanistreet, I.G.; Maiden, K.J. (1986): Preliminary results of a sedimentological study of the Chuos Formation in the Central Zone of the Damara Orogen: evidence for mass flow processes and glacial activity.- Communs Geol. Surv. S.W.Afr./ Namibia, **2**, 75-92.
- Henry, G.; Osborne, M.A.; Schmerold, R.K. (1992/1993): Proposed lithostratigraphic subdivision of the Ugab Subgroup (Damara Sequence) in Kaokoland, Namibia.- Communs Geol. Surv. Namibia, **8**, 143-145.
- Hensen, B.J.; Harley, S.L. (1990): Graphical analysis of P-T-X relations in granulite facies metapelites.- In: „High-temperature metamorphism and crustal anatexis“, J.R. Ashworth & M. Brown, eds., Unwin Hyman, London, 19-56.
- Hey, M.T. (1954): A new review of chlorites.- Mineral. Mag., **30**, 278-292.
- Hodges, K.V.; Spear, F.S. (1982): Geothermometry, geobarometry and the Al_2SiO_5 triple point at Mt. Moosilanke, New Hampshire.- Am. Mineral., **67**, 118-1134.
- Hoernes, S.; Hoffer, E. (1979): Equilibrium relations of prograde metamorphic mineral assemblages. A stable isotope study of rocks of the Damara Orogen, from Namibia.- Contrib. Mineral. Petrol., **68**, 377-389.
- Hoernes, S.; Hoffer, E. (1985): Stable isotope evidence for fluid-present and fluid-absent metamorphism in metapelites from the Damara Orogen, Namibia.- Contrib. Mineral. Petrol., **90**, 322-330.
- Hoffer, E. (1975): Die zeitliche Beziehung zwischen Tektogenese und Metamorphose: ein Beitrag zur Rekonstruktion der thermischen Geschichte des Damara-Orogen.- Bericht Sonderforschungsbereich 48, Universität Göttingen (unveröff.), 170-183.

- Hoffer, E. (1977): Petrologische Untersuchungen zur Regionalmetamorphose Al-reicher Metapelite im südlichen Damara-Orogen (Südwest-Afrika).- Habilitationsschrift, Universität Göttingen, 150 S.
- Hoffer, E. (1978): On the „late“ formation of paragonite and its breakdown in pelitic rocks of the southern Damara Orogen (Namibia).- *Contrib. Mineral. Petrol.*, **67**, 204-214.
- Hoffer, E. (1983): Compositional variations of minerals in metapelites involved in low to medium grade isograds reactions in the southern Damara Orogen, Namibia/ SW-Africa.- In „Intra-continental fold belts“, H. Martin & F.W. Ender, eds., Springer-Verlag, Berlin- Heidelberg, 745-766.
- Hoffer, E.; Puhan, D. (1975): Isoreaktionsgerade in den Metapeliten des Damara-Orogen.- Bericht Sonderforschungsbereich 48, Universität Göttingen (unveröff.), 111-151.
- Hoffmann, P.F.; Hawkins, D.P.; Isachsen, C.E.; Bowring, S.A. (1996): Precise U-Pb zircon ages for early Damaran magmatism in the Summas Mountains and Welwitschia Inlier, northern Damara Belt, Namibia.- *Communs Geol. Surv. Namibia*, **11**, 47-52.
- Hoffmann, K.-H. (1983): Lithostratigraphy and facies of the Swakop Group of the southern Damara Belt, SWA/Namibia.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., **11**, 43-63.
- Hoffmann, K.-H. (1987): Application of the tectonostratigraphic terrane analysis in the Damara Province of central and northern Namibia.- Abstr. Volume, Alex L. Du Toit Golden Jubilee Conference on Tectonostratigraphic Terrane Analysis, Geol. Soc. and Royal Soc. of S Afr., Cape Town, 25-27.
- Hoffmann, K.-H. (1989): New aspects of lithostratigraphic subdivision and correlation on the late Proterozoic to early Cambrian rocks of the southern Damara belt und their correlation with the central and northern Damara belt and the Gariep belt.- *Communs Geol. Surv. Namibia*, **5**, 59-67.
- Hoffmann, K.-H. (1990): Sedimentary depositional history of the Damara Belt related to continental breakup, passive to active margin transition and foreland basin development.- Ext. Abstr. 23rd Earth Sci. Congr., Geol. Soc. S. Afr., Cape Town, 250-253.
- Hoffmann; K.H.; Prave, A.R. (1996): A preliminary note on a revised subdivision and regional correlation of the Otavi Group based on glaciogenic diamictites and associated cap dolostones.- *Communs Geol. Surv. Namibia*, **11**, 77-82.
- Hoinkes, G. (1986): Effect of grossular-content in garnet on the partitioning of Fe and Mg between garnet and biotite.- *Contrib. Mineral. Petrol.*, **92**, 393-399.
- Hoisch, T.D. (1989): A muscovite-biotite geothermometer.- *Am. Mineral.*, **74**, 565-572.
- Hoisch, T.D. (1990): Empirical calibration of six geobarometers for the mineral assemblage quartz + muscovite + biotite + plagioclase + garnet.- *Contrib. Mineral. Petrol.*, **104**, 225-234.
- Holdaway, M.J.; Lee, S.M. (1977): Fe-Mg cordierite stability in high-grade pelitic rocks based on experimental, theoretical, and natural observations.- *Contrib. Mineral. Petrol.*, **63**, 175-198.
- Holdaway, M.J.; Dutrow, B.L.; Borthwick, J.; Shore, P.; Harmon, R.S.; Hinton, R.W. (1986a): H content of staurolite as determined by hydrogen extraction line and iron microprobe.- *Am. Mineral.*, **71**, 1135-1141.
- Holdaway, M.J.; Dutrow, B.L.; Shore, P. (1986b): A model of crystal chemistry of staurolite.- *Am. Mineral.*, **71**, 1142-1159.
- Holdaway, M.J.; Dutrow, B.L.; Hinton, R.W. (1988): Devonian and Carboniferous metamorphism in west-central Maine; the muscovite-almandine geobarometer and the staurolite problem revisited.- *Am. Mineral.*, **73**, 20-47.

- Holdaway, M.J.; Mukhopadhyay, B.; Dyar, M.D.; Dutrow, B.L.; Rumble, D.III; Grambling, J.A. (1991): A new perspective on staurolite crystal chemistry: use of stoichiometric and chemical end-members for a mole fraction model.- Am. Mineral., **76**, 1910-1919.
- Holdaway, M.J.; Mukhopadhyay, B.; Dutrow, B.L. (1995): Thermodynamic properties of stoichiometric staurolite $H_2Fe_4Al_{18}Si_8O_{48}$ and $H_6Fe_2Al_{18}Si_8O_{48}$.- Am. Mineral., **80**, 520-533.
- Holland, T.J.B.; Powell, R. (1985): An internally consistant thermodynamic data set with uncertainties and correlations : 2. Data and results.- J. Metamorph. Geol., **3**, 343-370.
- Holland, T.J.B.; Powell, R. (1990): An enlarged and updated internally consistant thermodynamic data set with uncertainties and correlations : the system $Na_2O - K_2O - CaO - MgO - MnO - FeO - Fe_2O_3 - Al_2O_3 - SiO_2 - TiO_2 - C - H_2 - O_2$.- J. Metamorph. Geol., **8**, 89-124.
- Holland, T.; Powell, R. (1992a): Plagioclase feldspars: Activity-composition relations based upon Darken's quadratic formalism and Landau theory.- Am. Mineral., **77**, 53-61.
- Holland, T.J.B.; Powell, R. (1992b): Updated thermodynamic data set from 1990.
- Holland, T.J.B.; Powell, R. (1998): An internally consistant thermodynamic data set for phases of petrological interest.- J. Metamorph. Geol., **16**, 309-343.
- Hollister, L.S. (1969): Metastabile paragenetic sequence of andalusite, kyanite, and sillimanite, Kwoiek area, British Columbia.- Amer. J. Sci., **267**, 352-370.
- Hollister, L.S.; Crawford, M.L. (1981): Fluid inclusions: Applications to petrology.- Mineral. Ass. of Canada, Calgary, Short Course Handbook, **6**, 304 p.
- Imeokparia, E.G.; Emofurieta, W.O. (1991): Protoliths and petrogenesis of Precambrian gneisses from the Igbedi area, SW Nigeria.- Chem. Erde; **51**, 1, 39-54.
- Indares, A.; Martignole, J. (1985a): Biotite-garnet geothermometry in the granulite facies; the influence of Ti and Al in biotite.- Am. Mineral., **70**, 272-278.
- Indares, A.; Martignole, J. (1985b): Biotite-garnet geothermometry in granulite-facies rocks; evaluation of equilibrium criteria.- Can. Mineral., **23**, 187-193.
- Jasper, M.J.U.; Stanistreet, I.G.; Charlesworth, E.G. (1992/93): Preliminary results of a study of the structural and sedimentological evolution of the late Proterozoic / early Palaeozoic Gariep Belt, southern Namibia.- Communs Geol. Surv. Namibia, **8**, 99-118.
- Jung; S.; Mezger, K. (1999): Age of crustal melting and leucogranite formation: An U-Pb monazite and Sm-Nd study of magmatic and anatexitic garnets from the central Pan-African Damara Orogen, Namibia.- Conference EUG 10, March 28 - April 01 1999, Strasbourg, Terra Abstracts, **11**, 106.
- Jung, S.; Metzger, K.; Masberg, P.; Hoffer, E.; Hoernes, S. (1998a): Petrology of an intrusion-related high-grade migmatite: implications for partial melting of metasedimentary rocks and leucosome-forming processes.- J. Metamorph. Geol., **16**, 425-445.
- Jung, S.; Mezger, K.; Hoernes, S. (1998b): Petrology and geochemistry of syn- to post-collisional metaluminous A-type granites - a major and trace element and Nd-Sr-Pb-O-isotope study from the Proterozoic Damara Belt, Namibia.- Lithos, **45**, 147-175.
- Jung, S.; Hoffer, E.; Masberg, P.; Hoernes, S. (1995): Geochemistry of granitic in-situ low-melt fractions - an example from the central Damara Orogen.- Communs Geol. Surv. Namibia, **10**, 21-31.
- Jung, S.; Hoernes, S.; Masberg, P.; Hoffer, E. (1999): The petrogenesis of some migmatites and granites (central Damara orogen, Namibia): Evidence for disequilibrium melting, wall-rock contamination and crystall fractionation.- J. Petrol., **40**, 8, 1241-1269.

- Kalt, A.; Altherr, R.; Ludwig, T. (1998): Contact metamorphism in pelitic rocks on the island Kos (Greece, Eastern Aegean Sea): a test for the Na-in-cordierite thermometer.- *J. Petrol.*, **39**, 4, 663-688.
- Kasch, K.W. (1979): Metamorphic history of the southern Damara belt around Omitara, SWA/Namibia.- *Precamb. Res. Unit, Annual Rep.*, **16**, Univ. Cape Town, 90-100.
- Kasch, K.W. (1980): The Damara metamorphic traverse.- *Precamb. Res. Unit, Annual Rep.*, **17**, Univ. Cape Town, 118-129.
- Kasch, K.W. (1981): The structural geology, metamorphic petrology and tectonothermal evolution of the southern Damara Belt around Omitara, SWA/Namibia.- Ph.D. thesis, Univ. Cape Town, 333p.
- Kasch, K.W. (1983a): Regional P-T variations in the Damara Orogen with particular reference to early high-pressure metamorphism along the southern margin.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., **11**, 243-253.
- Kasch, K.W. (1983b): Tectonothermal evolution of the southern Damara Orogen.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., **11**, 255-265.
- Kasch, K.W. (1983c): The structural geology, metamorphic petrology and tectonothermal evolution of the southern Damara Belt around Omitara, SWA/Namibia.- *Bull. Precamb. Res. Unit, Univ. Cape Town*, **27**, 333p.
- Kepezhinskas, K.B.; Khlestov, V.V. (1977): The petrogenetic grid and subfacies for middle-temperature metapelites.- *J. Petrol.*, **18**, 114-143.
- Kerrick, D.M. (1972): Experimental determination of muscovit + quartz stability with $P_{H2O} < P_{total}$.- *Am. J. Sci.*, **272**, 946-958.
- Kleemann, U.; Reinhardt, J. (1994): Garnet-biotite thermometry revisited: The effect of Al^{VI} and Ti in biotite.- *Eur. J. Mineral.*, **6**, 925-941.
- Klemd, R.; Okrusch, M. (1990): Phase relationships and fluid inclusions characteristics of the metamorphosed, stratiform sulfide deposit Matchless, Namibia.- *Geol. Rdsch.*, **79**, 433-449.
- Klemd, R.; Maiden, K.J.; Okrusch, M.; Richter, P. (1989): Geochemistry of metamorphosed massive sulphide deposit Matchless, South West Africa/Namibia: wallrock alteration during submarine ore-forming processes.- *Econ. Geol.*, **84**, 603-617.
- Klemd, R.; Bröcker, M.; Schramm, J. (1995): Characterisation of amphibolite-facies fluides of Variscan eclogites from the Orlica-Snieznik dome (Sudetes, SW Poland).- *Chem. Geol.*, **119**, 101-113.
- Koch-Müller, M. (1996): Fe-Mg exchange experiments between staurolite and garnet.- Sixth international symposium on experimental mineralogy, petrology and geochemistry, Bayreuth 10. - 13. April 1996, Terra abstracts, **8**, suppl. 1, 115.
- Koch-Müller, M. (1997): Experimentally determined Fe-Mg exchange between synthetic staurolite and garnet in the system MgO-FeO-Al₂O₃-SiO₂-H₂O.- *Lithos*, **41**, 185-212.
- Koziol, A.M., Bohlen, S.R. (1992): Solution properties of almandine-pyrope garnet as determined by phase equilibrium experiments.- *Am. Mineral.*, **77**, 765-773.
- Koziol, A.M.; Newton, R.C. (1988): Redetermination of the anorthite breakdown reaction and improvement of the plagioclase-garnet-Al₂SiO₅-quartz geobarometer.- *Am. Mineral.*, **73**, 216-223.
- Kriegsman, L.M.; Hensen, B.J. (1999): Restite-melt back reactions: implications for P-T paths and inferred tectonic settings.- Conference EUG 10, March 28 - April 01 1999, Strasbourg, Terra Abstracts, **11**, 715.

- Kroll, H.; Evangelakis, C.; Voll, G. (1993): Two-feldspar geothermometry: a review and revision for slowly cooled rocks.- *Contrib. Mineral. Petrol.*, **114**, 510-518.
- Kröner, A. (1980): Pan-African crustal evolution.- *Episodes*, **2**, 3-8.
- Kröner, A. (1982): Rb-Sr geochronology and tectonic evolution of the Pan-African Damara belt of Namibia, southwestern Africa.- *Amer. J. Sci.*, **282**, 1471-1507.
- Kröner, A.; Correia, H. (1980): Continuation of the Pan-African Damara belt into Angola: A proposed correlation of the Chela Group in southern Angola with the Nosib Group in northern Namibia/SWA.- *Trans. Geol. Soc. S. Afr.*, **83**, 5-16.
- Kröner; A.; Hawkesworth, C. (1977): Late Pan-African emplacement ages for Rössing alaskitic granite (Damara belt) and Rooi Lepel bostonite (Gariep belt) in Namibia and their significance for the timing of metamorphic events.- 20th a. Rep. Res. inst. afr. Geol., Univ. Leeds, 14-17.
- Kröner, A.; Halpern, M.; Jacob, R.E. (1978): Rb-Sr geochronology in favour of polymetamorphism in the Pan African Damara belt of Namibia (South West Africa).- *Geol. Rdsch.*, **67**, 2, 688-705.
- Kukla, C. (1993): Strontium isotope heterogenities in amphibolite facies, banded metasediments - A case study from the late Proterozoic Kuiseb Formation in the southern Damara Orogen, Central Namibia.- *Mem. Geol. Surv. Namibia*, **15**, 138p.
- Kukla, C.; Kramm, U.; Kukla, P.A.; Okrusch, M. (1991): U-Pb monazite data relating to metamorphism and granite intrusion in the northwestern Khomas Trough, Damara Orogen, central Namibia.- *Communs Geol. Surv. Namibia*, **7**, 49-54.
- Kukla, P.A. (1992): Tectonics and sedimentation of a late Proterozoic Damaran convergent continental margin, Khomas Hochland, central Namibia.- *Mem. Geol. Surv. Namibia*, **12**, 95p.
- Kukla, P.A.; Stanistreet, I.G. (1991): Record of the Damaran Khomas Hochland accretionary prism in central Namibia: Refutation of an „ensialic“ origin of a Late Proterozoic orogenic belt.- *Geology*, **19**, 473-476.
- Kukla, P.A.; Opitz, C.; Stanistreet, I.G.; Charlesworth, E.G. (1988): New aspects of the sedimentology and structure of the Kuiseb Formation in the western Khomas Trough, Damara Orogen, SWA/Namibia.- *Communs Geol. Surv. Namibia*, **4**, 33-42.
- Kukla, P.A.; Kukla, C.; Stanistreet, I.G.; Okrusch, M. (1990): Unusual preservation of sedimentary structures in sillimanite-bearing metaturbidites of the Damara Orogen, Namibia.- *J. Geol.*, **98**, 91-99.
- Laird, C.A.; Albee, A.L. (1981a): High-pressure metamorphism in mafic schist from Northern Vermont.- *Amer. J. Sci.*, **281**, 97-126.
- Laird, C.A.; Albee, A.L. (1981b): Pressure, temperature, and time indicators in mafic schist: Their application to reconstructing the polymetamorphic history of Vermont.- *Amer. J. Sci.*, **281**, 127-175.
- Le Breton, N.; Tompson, A.A. (1988): Fluid-absent (dehydration) melting of biotite in metapelites in the early stages of crustal anatexis.- *Contrib. Mineral. Petrol.*, **99**, 226-237.
- Lehtonen, M.I.; Manninen, T.E.T.; Schreiber, U.M. (1996): Lithostratigraphy of the area between the Swakop, Khan and lower Omaruru rivers, Namib Desert.- *Communs Geol. Surv. Namibia*, **11**, 65-75.
- Le Maitre, R.W.; Bateman, P.; Dudek, A.; Keller, J.; Lameyre Le Bas, M.J.; Sabine, P.A.; Schmid, R.; Sorensen, H.; Streckeisen, A.; Woolley, A.R.; Zanettin, B. (1989): A classification of igneous rocks and glossary of terms.- Blackwell, Oxford, 193p.

- Lindsley, D.H. (1973): Delimination of the hematite-ilmenite miscibility gap.- *Geol. Soc. Am. Bull.*, **84**, 657-662.
- Machado, N.; Valladares, C.; Heilbron, M.; Valeriano, C. (1996): U-Pb geochronology of the central Ribeira belt (Brazil) and implications for the evolution of the Brazilian Orogeny.- *Precamb. Res.*, **79**, 347-361.
- Mahar, E.M.; Baker, J.M.; Powell, R.; Holland, T.J.B.; Howell, N. (1997): The effect of Mn on stability in metapelites.- *J. Metamorph. Geol.*, **15**, 223-238.
- Marlow, A.G. (1983): Geology and Rb-Sr geochronology of mineralized and radioaktive granites and alaskites, Namibia.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 289-298.
- Martin, H. (1965): The Precambrian geology of South West Africa and Namaqualand.- Precam. Res. Unit, Univ. Cape Town, 159p.
- Martin, H. (1983): Overview of the geosynclinal, structural, and metamorphic development of the intracontinental branch of the Damara Orogen.- In „Intracontinental fold belts“, H. Martin & F.W. Ender, eds., Springer-Verlag, Berlin-Heidelberg, 473-502..
- Martin, H.; Porada, H. (1977a): The intracratonic branch of the Damara Orogen in South West Africa: I. Discussion of geodynamic models.- *Precamb. Res.*, **5**, 311-338.
- Martin, H.; Porada, H. (1977b): The intracratonic branch of the Damara Orogen in South West Africa: II. Discussion of relationships with the Pan-African mobile system.- *Precamb. Res.*, **5**, 339-357.
- Masberg, P. (1996): Multiple migmatization in the HT/LP Central Damara Orogen, Namibia.- *N. Jb. Mineral. Abh.*, **170**, 3, 257-289.
- Masberg, H.P.; Hoffer, E.; Hoernes, S. (1992): Microfabrics indicating granulite-facies metamorphism in the low-pressure central Damara-Orogen, Namibia.- *Precamb. Res.*, **55**, 243-257.
- Massone, H.-J. (1991): High-pressure, low temperature metamorphism of pelitic and other protoliths based on experiments in the system K₂O-MgO-Al₂O₃-SiO₂-H₂O.- Habilitations-schrift, Ruhr-Universität, Bochum, 172 p.
- Massone, H.-J.; Schreyer, W. (1987): Phengite geobarometry based on the limiting assemblage with K-feldspar, phlogopite, and quartz.- *Contrib. Mineral. Petrol.*, **96**, 212-224.
- Matthes, S. (1993): Mineralogie.- Springer-Verlag, Berlin-Heidelberg, 4. Auflage, 461 S.
- Ménot, R.-P.; Seddoh, K.F. (1985): The eclogites of the Lato hills, south Togo, west Africa: relics from the early tectonometamorphic evolution of the Pan-African orogeny.- *Chem. Geol.*, **50**, 313-330.
- Miller, R.McG. (1979): The Okahandja Lineament, a fundamental tectonic boundary in the Damara Orogen of South West Africa/ Namibia.- *Trans. Geol. Soc. S. Afr.*, **82**, 349-361.
- Miller, R.McG. (ed.) (1983a): Evolution of the Damara Orogen of South West Africa/Namibia.- Spec. Publ. **No. 11**, Geol. Soc. South Africa, 515p.
- Miller, R.McG. (1983b): The Pan-African Damara Orogen of South West Africa/Namibia.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 431-515.
- Miller, R.McG. (1991): The framework of the Damara Orogen, Namibia.- In „Precambrian Sedimentary basins of Southern Africa“, P.G. Eriksson, ed., Terra Abstracts, **3**, 22-23.
- Miller, R.McG.; Burger, A.J. (1983a): U-Pb zircon ages of the early Damaran Naauwpoort Formation.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 267-272.

- Miller, R.McG.; Burger, A.J. (1983b): U-Pb zircon ages of members of the Salem granitic suite along the northern edge of the Central Damara Belt .- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 273-280.
- Miller, R.McG.; Grote, W. (1988): Geological map of the Damara Orogen South West Africa/Namibia, 1 : 500 000.- Geol. Survey of Namibia, Windhoek.
- Miller R. McG; Schalk, K.E.L. (1980): Geological Map of Namibia, 1 : 1 000 000, Sheet 1 - 4, Reprinted 1990, Geol. Survey of Namibia, Windhoek.
- Miller, R.McG.; Freyer, E.E.; Häblich, I.W. (1983): A turbidite succession equivalent to the entire Swakop Group.- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 65-71.
- Mirwald, P. (1986): Ist Cordierit ein Geothermometer? Fortschr. Mineral., **64**, Bh.1, 119.
- Miyashiro, A. (1973): Metamorphism and metamorphic belts.- Allen & Unwin LTD, London, 492p.
- Neves, S.P.; Vauchez, a.; Ferand, G. (2000): Tectono-thermal evolution, magma emplacement, and shear zone development in the Caruaru area (Borborema Province, NE Brazil).- Precamb. Res., **99**, 1-32.
- Newton, R.C.; Haselton, H.T. (1981): Thermodynamics of the garnet - plagioclase - Al_2SiO_5 - quartz geobarometer.- In „Thermodynamics of minerals and melts“, Advances in Phys. Geochem., R.C. Newton, ed., Springer-Verlag, New York, **1**, 131-147.
- Nichols, G.T.; Berry, R.F.; Green, D.H. (1992): Internally consistant gahnitic spinel-cordierite-garnet equilibria in the FMASHZn system: geothermometry and applications.- Contrib. Mineral. Petrol., **111**, 362-377.
- Ohmoto, H.; Kerrick, D. (1977): Devolatilization equilibria in graphitic systems.- Am. J. Sci., **277**, 1013-1044.
- Oliver, G.J.H. (1994): Mid-crustal detachment and domes in the central zone of the Damaran Orogen, Namibia.- J. Afr. Earth Sci., **19**, 4, 331-344.
- Pattinson, D.R.M. (1992): Stability of andalusite and sillimanite and the Al_2SiO_5 triple point: Constraints from the Ballachulish Aureole, Scotland.- J. Geol., **100**, 423-446. ($550 \pm 35^\circ\text{C}$ / $4.5 \pm 0.5\text{kbar}$)
- Perchuk, L.L. (1969): The staurolite - garnet thermometer.- Akademij Nauk USSR, Doklady Earth Science Sections, 189-191.
- Perchuk, L.L.; Fedkin, V.V. (1974): Staurolite-garnet equilibria as geothermometers.- In „Energetics of geological processes“, S.K. Saxena & S. Bhattacharij, eds., 1977, Springer Verlag, New York, 318-320.
- Perchuk, L.L.; Lavrent'eva, I.V. (1983): Experimental investigation of exchange equilibria in the system cordierite - garnet - biotite.- In „Kinetics and equilibrium in mineral reactions“, S.K. Saxena, ed., Springer-Verlag, New York, 199-239.
- Porada, H. (1979): The Damara-Ribeira orogen of the Pan-African-Brasiliano cycle in Namibia (South West Africa) and Brazil as interpreted in terms of continental collision.- Tectono-physics, **57**, 237-265.
- Porada, H. (1985): Stratigraphy and facies in the upper Proterozoic Damara Orogen, Namibia, based on a geodynamic model.- Precamb. Res., **29**, 235-264.
- Powell, R.; Holland, T.J.B. (1988): An internally consistent dataset with uncertainties and correlations: 3. Applications to geobarometry, worked examples and a computer programm (Thermocalc v2.3).- J. Metamorph. Geol., **6**, 173-204.

- Powell, R.; Holland, T. (1990): Calculated mineral equilibria in the pelite system, KFMASH (K_2O - FeO - MgO - Al_2O_3 - SiO_2 - H_2O).- Am. Mineral., **75**, 367-380.
- Pownceby, M.I.; Wall, V.J.; O'Neill, H.S.C. (1987a): Fe-Mn partitioning between garnet and ilmenite; experimental calibration and applications.- Contrib. Mineral. Petrol., **97**, 116-126.
- Pownceby, M.I.; Wall, V.J.; O'Neill, H.S.C. (1987b): Fe-Mn partitioning between garnet and ilmenite; experimental calibration and application.- Contrib. Mineral. Petrol., **97**, 539.
- Prave, A.R. (1996): Tale of three cratons: Tectonostratigraphic anatomy of the Damara orogen in northwestern Namibia and the assembly of Gondwana.- Geology, **28**, 12, 1115-1118.
- Puhan, D. (1977): Metamorphic temperature determined by means of the dolomite-calcite solvus geothermometer - examples from the central Damara Orogen (SW Africa).- Contrib. Mineral. Petrol., **58**, 23-28.
- Puhan, D. (1983): Metamorphism of siliceous dolomites of the central and southern part of the Damara Orogen.- In „Intracontinental fold belts“, H. Martin & F.W. Ender, eds., Springer-Verlag, Berlin-Heidelberg, 767-784.
- Reid, D.L.; Ransome, I.G.D.; Onstott, T.C.; Adams, C.J. (1991): Time of emplacement and metamorphism of Late Precambrian mafic dykes associated with the Pan-African Gariep orogeny, Southern Africa: implications for the age of the Nama Group.- J. Afr. Earth Sci., **13**, 531-541.
- Ribbe, P.H. (1983): Chemistry, structure and nomenclature of feldspars.- Mineral. Soc. Am., Rev. in Mineral., **2**, 1-19.
- Roedder, E. (1984): Fluid inclusions.- Mineral. Soc. Am., Rev. in Mineral., **12**, 644 p.
- Rollinson, H. (1993): Using geochemical data: evaluation, presentation, interpretation.- Longman Scientific & Technical, New York, U.S.A., 325 p.
- Roser, B.P.; Korsch, R.J. (1986): Determination of tectonic setting of sandstone-mudstone suites using SiO_2 content and K_2O/Na_2O ratio.- J. Geol., **94**, 635-650.
- Roser, B.P.; Korsch, R.J. (1988): Provenance signatures of sandstone-mudstone suites determined using discriminant function analysis of major-element data.- Chem. Geol., **67**, 119-139.
- Rumble, D. III (1973): Fe-Ti oxide minerals from regionally metamorphosed quartzites of eastern New Hampshire.- Contrib. Mineral. Petrol., **42**, 181-195.
- Rumble, D. III (1976): Oxide minerals in metamorphic rocks.- In „Oxide minerals“, Rumle, D. III, ed., Mineral. Soc. Am. Short Course Note **3**, R1-R24.
- SACS - South African Committee for Stratigraphy (1980): The Damara Sequence.- In „The Stratigraphy of South Africa, Part 1, Lithostratigraphy of the Republic of South Africa, South West Africa/ Namibia, and the Republics of Bophutatswana, Transkei and Venda“, L.Kent & Co., eds., Handbook **8**, Geol. Surv. S. Afr., 415-438.
- Saunders, A.D.; Tarney, J.; Weaver, S.D. (1980): Transverse geochemical variations across the Antarctic Peninsula; implications for the genesis of calc-alkaline magmas.- Earth Planet Sci. Lett., **46**, 3, 344-360.
- Sawyer, E.W. (1981): Damaran structural and metamorphic geology in an area southeast of Walvisbay, SWA/ Namiba.- Mem. Geol. Surv. SW Afrika/ Namiba, **7**, 1-94.
- Saxena, S.K. (ed.) (1973): Thermodynamics of rock-forming crystalline solutions, minerals, rocks and inorganic materials.- Springer-Verlag, Berlin- Heidelberg, 140p.
- Saxena, S.K. (ed.) (1983): Kinetics and equilibrium in mineral reactions.- Springer-Verlag, New York, 273p.

- Schenk, V.; Herms, P.; LeBreton, N.; Craven, J.; Harley, S. (1992): Cordierit als Fluidindikator in metamorphen Gesteinen.- Eur. J. Mineral., **4**, Beih. 1, 238.
- Schliestedt, M. (1980): Phasengleichgewichte in Hochdruckgesteinen von Sifnos, Griechenland.- Dissertation, Universität Braunschweig, Braunschweig, 142 S.
- Schmädicke, E. (1992): Die Eklogite des Erzgebirges.- Dissertation, Universität Würzburg, Würzburg, 338 S.
- Schmidt, A.; Wedepol, K.H. (1983): Chemical composition and genetic relations of the Matchless Amphibolite (Damara Orogenic Belt).- In „Evolution of the Damara Orogen“, R.McG. Miller, ed., Spec. Publ. Geol. Soc. S. Afr., 11, 139-145.
- Schmitt, R.S.; Trow, R.A.J.; Van Schmus, W.R. (1998): U/Pb ages of Transamazonian (2.0 Ga) and Cambrian (0.5 Ga) orogenic events in the Ribeira Belt, SE Brazil; constraints on the history of Gondwana.- Geological Society of America (GSA), 1998 annual meeting, abstracts with programs, **30**, 7, 97.
- Schreyer, W. (1985): Experimental studies on cation substitutions and fluid incorporation in cordierite.- Bull. Soc. Fr. Minéral., **108**, 273-291.
- Schwandt, C.S.; Cygan, R.T.; Westrich, H.R. (1995): Mg self-diffusion in pyrope garnet.- Am. Mineral., **80**, 483-490.
- Schwandt, C.S.; Cygan, R.T.; Westrich, H.R. (1996a): Ca self-diffusion in grossular garnet.- Am. Mineral., **81**, 448-451.
- Schwandt, C.S.; Papike, J.J.; Shearer, C.K. (1996b): Trace element zoning in garnet of the Black Hills, South Dakota.- Am. Mineral., **81**, 1195-1207.
- Seck, H.A. (1971): Der Einfluß des Drucks auf die Zusammensetzung koexistierender Alkali-feldspäte und Plagioklase im System NaAlSi₃O₈ - KAlSi₃O₈ - CaAl₂Si₂O₈ - H₂O.- Contrib. Mineral. Petrol., **31**, 67-86.
- Seth, B. (1999): Crustal evolution of the Kaoko belt, NW Namibia: Geochemical and geochronological study of Archaean to Mesoproterozoic basement gneisses and Pan-African migmatites and granitoids.- Dissertation, Universität Würzburg, Würzburg, 120 p.
- Seth, B.; Kröner, A. (1999): Pan-African multiphase magmatism and implications for metamorphic events in the Kaoko belt of NW Namibia.- Conference EUG 10, March 28 - April 01 1999, Strasbourg, Terra Abstracts, **11**, 105-106.
- Seth, B.; Kröner, A.; Dürr, S.B.; Dingeldey, D.P. (1996): Speculations on the origin of a late Archaean crustal fragment in the Kaoko belt, Namibia, and geodynamic implications.- 1996 Fall Meeting AGU, Suppl. to EOS, Transactions AGU, **77**, No. **46**, F820-F821.
- Seth, B.; Kröner, A.; Dürr, S.B.; Dingeldey, D.P.; Okrusch, M. (1997): Archaean terrane identification and late Pan-African metamorphism in the Kaoko belt, Namibia: significance for Gondwana assembly.- Conference EUG 9, March 23-27 1997, Strasbourg, Terra Nova, **9**, 168.
- Seth, B.; Kröner, A.; Mezger, K.; Nemchin, A.A.; Pidgeon, R.T.; Okrusch, M. (1998): Archaean to Neoproterozoic magmatic events in the Kaoko belt of NW Namibia and their geodynamic significance.- Precamb. Res., **92**, 4, 341-363.
- Shepherd, T.J.; Rankin, A.H.; Alderton, D.H.M. (1985): A practical guide to fluid inclusion studies.- Blackie, Glasgow - London, 239 p.
- Smoler, M. (1987): Petrographische, geochemische und phasenpetrologische Untersuchungen an Metasedimenten des NW-Spessarts/ Bayern.- Dissertation, Universität Würzburg, Würzburg, 256 S.

- Spear, F.S. (1988): The Gibbs method and Duhem's theorem: The quantitative relationships among P, T, chemical potential, phase composition and reaction progress in igneous and metamorphic systems.- *Contrib. Mineral. Petrol.*, **99**, 249-256.
- Spear, F.S. (1993): Metamorphic phase equilibria and pressure-temperature-time paths.- *Mineral. Soc. America*, Washington DC, 799S.
- Spear, F.S.; Cheney, J.T. (1989): A petrogenetic grid for pelitic schists in the system SiO₂-Al₂O₃-FeO-MgO-K₂O-H₂O.- *Contrib. Mineral. Petrol.*, **101**, 149-164.
- Spear, F.S.; Menard, T. (1989): Program GIBBS: a generalized Gibbs methode algorithm.- *Am. Mineral.*, **74**, 942-943.
- Spear, F.S.; Peacock, S.M. (eds.) (1989): Metamorphic pressure-temperature-time paths.- Short Course in Geology, Am. Geophys. Union, Washington DC, 1-55.
- Spear, F.S.; Selverstone, J. (1983): Quantitative P-T paths from zoned minerals. Theory and tectonic applications.- *Contrib. Mineral. Petrol.*, **83**, 348-357.
- Spear, F.S.; Peacock, S.M.; Kohn, M.J.; Florence, F.P.; Menart, T. (1991a): Computer programs for petrological P-T-t path calculations.- *Ibid*, **76**, 2009-2012.
- Spear, F.S.; Kohn, M.J.; Florence, F.P.; Menard, T. (1991b): A model for garnet and plagioclase growth in pelitic schists: implications for thermobarometry and P-T path determinations.- *J. Metamorph. Geol.*, **8**, 683-696.
- Spear, F.S.; Kohn, M.; Menart, T.; Florence, F. (1997): Program Gibbs v.4.7, May 1997 (Computer program and manual).- Dept. of Geology, Rensselaer Polytechnic Institute, Troy, New York 12180, U.S.A. (Spear home page).
- Spear, F.S.; Kohn, J.; Cheney, J.T. (1999): P-T paths from anatetic pelites.- *Contrib. Mineral. Petrol.*, **134**, 17-32.
- Stanistreet, I.G.; Kukla, P.A.; Henry, G. (1991): Sedimentary basinal responses to a late Precambrian Wilson Cycle: the Damara Orogen and Nama Foreland, Namibia.- *J. Afr. Earth Sci.*, **13**, 1, 141-156.
- Steven, N.M.; Armstrong, R.A.; Moore, J.M. (1992/93): New Rb-Sr data from the Central Zone of the Damara Orogen, Namibia.- *Communs Geol. Surv. Namibia*, **4**, 71-81.
- Symmes, G.H.; Ferry, J.M. (1991): Evidence from mineral assemblages for infiltration of pelitic schists by aqueous fluids during metamorphism.- *Contrib. Mineral. Petrol.*, **108**, 419-438.
- Symmes, G.H.; Ferry, J.M. (1992): The effect of whole-rock MnO content on the stability of garnet in pelitic schists during metamorphism.- *J. Metamorph. Geol.*, **10**, 221-237.
- Tack, L.; Boven, A.; Bowden, P.; Oliver, G. (1995): The significance of ⁴⁰Ar/³⁹Ar ages on a mylonitic cover gneiss from the contact shear zone of the Ida dome, central Damaran belt, western Namibia.- *Mus. roy. Afr. centr., Tervuren (Belg.) Dépt. Géol. Min., Rapp. ann.*; 1993 & 1994, 159-172. (*Royal Museum of Central Africa, Tervuren, Belg.*)
- Tegtmeyer, A.; Kröner, A. (1985): U-Pb zircon ages for granitoid gneisses in northern Namibia and their significance for Proterozoic evolution of southern Africa.- *Precamb. Res.*, **28**, 311-326.
- Thélin, P. (1983): Les gneiss œillés de la nappes du Grand Saint-Bernard. Essai d'évaluation des critères susceptibles d'en préciser l'hérédité pré-métamorphique (Alpes valaisannes, Suisse).- Thèse de doctorat, Université de Lausanne.
- Torquato, J.R. (1977): Geotectonic correlation between SE Brazil and SW Africa.- *An. Acad. Bras. Cienc.*, **48** (Supl.), 353-363.
- Touret, J.L.R.; Dietvorst, P. (1983): Fluid inclusions in high-grade anatetic metamorphites.- *J. Geol. Soc. London*, **140**, 635-649.

- Tracy, R.J.; Miller, S.J.; Solberg, T.N.; Waldron, K.A. (1992): Documentation of diffusional modification of garnet growth zoning in amphibolite facies garnets.- Geol. Soc. Am. Abstracts with Programs, **24**, A219.
- Tröger, W.E. (1982): Optische Bestimmung der gesteinsbildenden Minerale, Teil 2.- E. Schweizerbartsche Verlagsbuchhandlung , Stuttgart, 822 S.
- Trompette, R. (1994): Geology of western Gondwana (2000 - 500 Ma), Pan-African-Brasiliano aggregation of South America and Africa.- Balkema, Rotterdam, 350 p.
- Unrug, R. (ed.) (1996): The Geodynamic Map of Gondwana Supercontinent Assembly; Scale 1 : 10 million.- Publ. by Bureau de Recherches Géologiques et Minières, France.
- Unrug, R. (1997): Rodinia to Gondwana: The Geodynamic Map of Gondwana Supercontinent Assembly.- GSA Today, **7**, 1, 1-5.
- Veh, M.W. von (1993): The stratigraphy and structural evolution of the late Proterozoic Gariep belt in the Sendelingsdrif-Annisfontein area, northwestern Cape Province.- Bull. Precamb. Res. Unit, Univ. Cape Town, **38**, 174p.
- Vielzeuf, D.; Holloway, J.R. (1988): Experimental determination of the fluid-absent melting relations in the pelitic system.- Contrib. Mineral. Petrol., **98**, 257-276.
- Vielzeuf, D.; Montel, J.M. (1994): Partial melting of metagreywackes. Part 1. Fluid-absent experiments and phase realtionships.- Contrib. Mineral. Petrol., **117**, 375-393.
- Visser, D.; Kloprogge, J.T.; Maijer, C. (1994): An infrared spectroscopic (IR) and light element (Li, Be, Na) study of cordierites from the Bamble Sector, South Norway.- Lithos, **32**, 95-107.
- Vry, J.K.; Brown, P.E.; Valley, J.W. (1990) Cordierite volatile content and the role of CO₂ in high-grade metamorphism.- Am. Mineral., **75**, 71-88.
- Waldbaum, D.R.; Thompson, J.B. (1969): Mixing properties of sanidine crystalline solutions; IV. Phase diagrams from equations of state.- Am. Mineral., **54**, 1274-1298.
- Will, T.M. (1997): Kursunterlagen zur Lehrveranstaltung Phasendiagramme.- Mineralogisches Institut, Universität Würzburg.
- Will, T.M. (1998a): Phase diagrams and their application to determine pressure-temperature paths of metamorphic rocks.-N. Jb. Mineral. Abh., **174**, 2, 103-130.
- Will, T.M. (1998b): Phase equilibria in metamorphic rocks: thermodynamic background and petrological applications.- Lecture notes in earth sciences, **71**, Springer-Verlag, Berlin-Heidelberg, 315 p.
- Will, T.M.; Okrusch, M.; Schmädicke, E.; Chen, G. (1998): Phase relations in greenschist-blueschist-amphibolite-eclogite facies: Calculated mineral equilibria in the system Na₂O-CaO-FeO-MgO-Al₂O₃-SiO₂-H₂O, with applications to the PT evolution of metamorphic rocks from Samos, Greece.- Contrib. Mineral. Petrol., **132**, 85-102.
- Wimmenauer, W. (1984): Das prävariskische Kristallin im Schwarzwald.- Fortschr. Mineral., Beih., **62**, 69-86.
- Wolfsdorff, P.; Schreyer, W. (1992): Synthesis of sodian cordierites in the system Na₂O-MgO-Al₂O₃-SiO₂.- N. Jb. Mineral., Mh., **2**, 80-96.
- Xu, G.; Will, T.M.; Powell, R. (1994): A calculated petrogenetic grid for the system K₂O-FeO-MgO-Al₂O₃-SiO₂-H₂O, with particular reference to contact-metamorphosed pelites.- J. Metamorph. Geol., **12**, 99-119.
- Zeh, A. (1995): Die Druck-Temperatur-Deformations-Entwicklung des Ruhlaer Kristallins (Mitteldeutsche Kristallinzone).- Dissertation, Universität Würzburg, Würzburg, 254 S.