

## 6 LITERATURVERZEICHNIS

- AGRE P. (1997) *Biol. Cell.* (89): 255-57
- AGRE P. (1998) Aquaporin null phenotypes: The importance of classical physiology. *Proc. Natl. Acad. Sci. USA* (95): 9061-9063
- AGRE P., LEE M. D., DEVIDAS S., GUGGINO W. B., SASAKI S., *et al.* (1997) *Science* (275): 1490-92
- AGRE P., SABOORI A. M., ASIMOS A., SMITH B. L. (1987) Purification and partial characterization of the M<sub>r</sub> 30,000 integral membrane protein associated with the erythrocyte Rh(D) antigen. *J. Biol. Chem.* (262): 17497-503
- BALLING A. AND ZIMMERMANN U. (1990) Comparative measurements of the xylem pressure of *Nicotiana* plants by means of the pressure bomb and pressure probe. *Planta* (182): 325-338
- BARONE L. M., SHIH C. AND WASSERMAN B. P. (1997) Mercury-induced Conformational Changes and Identifikation of Conserved Surface Loops in Plasma Membrane Aquaporins from Higher Plants: Topology of PMIP31 from *Beta vulgaris* L. *J. Biological Chemistry* 272(49): 30672-30677
- BIELA A., GROTE K., OTTO B., HOTH S., HEDRICH R. AND KALDENHOFF R. (1999) The *Nicotiana tabacum* plasma membrane aquaporin NtAQP1 is mercury-insensitive and permeable for glycerol. *The Plant Journal* 18(5): 565-570
- BORGNIA M., NIELSEN S., ENGEL A. AND AGRE P. (1999) Cellular and Molecular Biology of the Aquaporin Water Channels. *Annu. Rev. Biochem.* (68): 425-458
- CARVAJAL M., COOKE D. T., CLARKSON D. T. (1996) Responses of wheat plants to nutrient deprivation may involve the regulation of water-channel function. *Planta* (199): 372-381
- CHENG A., VAN HOEK A. N., YEAGER M., VERKMAN A. S. AND MITRA A. K. (1997) Three-dimensional organization of a human water channel. *Nature* (387): 627-630
- CONKLING M. A., CHENG C. L., YAMAMOTO Y. T. AND GOODMAN H. M. (1990) Isolation of transcriptionally regulated root-specific genes from tobacco. *Plant Physiology* (93): 1203-1211
- COOPER G. J. AND BORON W. F. (1998) Effect of PCMBs on CO<sub>2</sub> permeability of *Xenopus* oocytes expressing aquaporin 1 or its C189S mutant. *Am. J. Physiol.* 275 (*Cell Physiol.* 44): C1481-C1486
- CUBITT A. B., HEIM R., ADAMS S. R., BOYD A. E., GROSS L. A. AND TSIEN R. Y. (1995) Understanding, improving and using green fluorescent proteins. *TIPS* (20) November 1995: 448-455
- DAINTY J. (1963) Water relation of plant cells. *Adv. Bot. Res.* (1): 279-326
- DANIELS M. J., CHAUMONT F., MIRKOV T. E., CHRISPEELS M. J. (1996) Characterization of a new vacuolar membrane aquaporin sensitive to mercury at a unique site. *Plant Cell* (8): 587-99
- DANIELS M. J., MIRKOV T. E., CHRISPEELS M. J. (1994) The plasma membrane of *Arabidopsis thaliana* contains a mercury-insensitive aquaporin that is a homolog of the tonoplast water channel protein TIP. *Plant Physiol.* (106): 1325-33

- DENKER B. M., SMITH B. L., KUHAJDA F. P., AGRE P. (1988) *J. Biol. Chem.* (263): 15634-42
- ECHIVARRÍA M., WINDHAGER E. E., FRINDT G. (1996) Selectivity of the renal collecting duct water channel aquaporin-3. *J Biol Chem* Oct 11; (271:41): 25079-82
- FEINBERG P. A., VOGELSTEIN B. (1983) A technique for radiolabelling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* (132): 6-13
- FEINBERG P. A., VOGELSTEIN B. (1984) Addendum: A technique for radiolabelling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* (137): 266-267
- FETTIPLACE R. AND HAYDON D. A. (1980) Water permeability of lipid membranes. *Physiological Reviews* (60): 510-550
- FINKEL A. S., GAGE P. W. (1985) Conventional voltage clamping with two intracellular microelectrodes. In: Voltage and Patch Clamping with Microelectrodes (Smith T. G., Lecar H., Redman S. J., Gage P. W., Eds., William & Wilkins Co., Baltimore, Maryland: 47-94
- FLEURAT-LESSARD P., FRANGNE N., MAESHIMA M., RATAJCZAK, BONNEMAIN J.-L., MARTINOIA E. (1997) Increased Expression of Vakuolar Aquaporin and H<sup>+</sup>-ATPase Related to motor Cell Funktion in *Mimosa pudica* L. *Plant Physiol.* (114): 827-834
- GORIN, M. B., YANCEY, S. B., CLINE, J., REVEL, J.-P. AND HORWITZ, J. (1984) The major intrinsic protein (MIP) of the bovine lens fiber membrane: characterization and structure based on cDNA cloning, *Cell* (39): 49-59.
- GORTON H. L. (1987) Water Relations in Pulvini from *Samanea Saman*. *Plant Physiol.* (83): 945-950
- GORTON H. L. (1990) In: Satter R. L., Gorton H. L., Vogelmann T. C. (eds) The pulvinus: motor organ for leaf movement. *American Society of Plant Physiologists, Bethesda*, Vol. 3: 223-237
- GROTE K., VON TRZEBIATOVSKI P. AND KALDENHOFF R. (1998) RNA levels of plasma membrane aquaporins in *Arabidopsis thaliana*. *Protoplasma* (204): 139-144
- GUERRERO F. D., JONES J. T., MULLET J. E. (1990) Turgor-responsive gene transcription and RNA levels increase rapidly when pea shoots are wilted. Sequence and expression of three inducible genes. *Plant Mol. Biol.* (15): 11-26
- GUTKNECHT J. (1967) Membranes of *Valonia ventricosa*: apparent absence of water-filled pores. *Science* (158): 787-788
- HABERLANDT G. (1884) Physiologische Pflanzenanatomie. 1. Auflage, Leipzig: Engelmann Verlag
- HENZLER T., STEUDLE E. (1995) Reversible closing of water channels in *Chara* internodes provides evidence for a composite transport model of the plasma membrane. *J. Exp. Bot.* (46): 199-209
- HERTEL A., STEUDLE E. (1997) The function of water channels in *Chara*: The temperature dependence of water and solute flows provides evidence for composite membrane transport and for a slippage of small organic solutes across water channels. *Planta* (202): 324-335

- HEYMANN J. B., AGRE P., ENGEL A. (1998) Progress on the Structure and Function of Aquaporin 1, *Journal of Structural Biology* 121(2): 191-206
- HÖFTE H., HUBBARD L., REIZER J., LUDEVID D., HERMAN E. M. AND CHRISPEELS M. J. (1992) Vegetative and seed-specific forms of tonoplast intrinsic protein in the vacuolar membrane of *Arabidopsis thaliana*. *Plant Physiology* (99): 561-570
- ISHIBASHI K., KUWAHARA M., KAGEYAMA Y., TOHSAKA A., MARUMO F., SASAKI S. (1997a) Cloning and functional expression of a second new aquaporin abundantly expressed in testis. *Biochem Biophys Res Commun* Aug 28; (237:3): 714-718
- ISHIBASHI K., KUWAHARA M., KAGEYAMA Y., TOHSAKA A., MARUMO F., SASAKI S. (1997b) Cloning and functional expression of a new aquaporin abundantly expressed in the testis permeable to water, glycerol and urea. *Journal Biol. Chem.* Aug 15; (272:33): 20782-20786
- JAP B. K., LI H. (1995) Structure of the osmo-regulated H<sub>2</sub>O-channel, AQP-CHIP, in projection at 3.5 Å resolution. *J. Mol. Biol.* (251): 413-420
- JOHANSSON I., KARLSON M., SHUKLA V. K., CHRISPEELS M. J., LARSON C., KJELLBOM P. (1998) Water Transport Activity of the Plasma Membrane Aquaporin PM28A Is Regulated by Phosphorylation. *The Plant Cell* (10): 451-459
- JOHNSON K. D., HERMAN E. M., CHRISPEELS M. J. (1989) An abundant, highly conserved tonoplast protein in seeds. *Plant Physiol.* (91): 1006-13
- JOHNSON K. D., CHRISPEELS M. J. (1992) Tonoplast-bound protein kinase phosphorylates tonoplast intrinsic protein. *Plant Physiol.* (100): 1787-95
- JOHNSON K. D., HÖFTE H., CHRISPEELS M. J. (1990) An intrinsic tonoplast protein of storage vacuoles in seeds is structurally related to a bacterial solute transporter (GlpF). *Plant Cell* (2): 525-532
- JUNG J. S., PRESTON G. M., SMITH B. L., GUGGINO W. B., AGRE P. (1994) Molecular structure of the water channel through aquaporin CHIP: the hourglass model. *J. Biol. Chem.* (269): 14648-54
- KALDENHOFF R., GROTE K., ZHU J., ZIMMERMANN U. (1998) Significance of plasmalemma aquaporins for watertransport in *Arabidopsis thaliana*. *The Plant Journal* 14 (1): 121-128
- KALDENHOFF R., KÖLLING A., MEYERS A., KARMAN U., RUPPEL G., RICHTER G. (1995) The blue light-responsive AthH2 gene of *Arabidopsis thaliana* is primarily expressed in expanding as well as in differentiating cells and encodes a putative channel protein. *Plant Journal* (7): 87-95
- KALDENHOFF R., KÖLLING A., RICHTER G. (1993) A novel blue light- and abscisic acid-inducible gene of *Arabidopsis thaliana* encoding an intrinsic membrane protein. *Plant Mol. Biol.* (23): 1187-98
- KAMMERLOHER W., FISCHER U., PIECHOTTKA G. P., SCHÄFFNER A. R. (1994) Water channels in the plant plasma membrane cloned by immunoselection from a mammalian expression system. *Plant J.* (6): 187-99

- KIYOSAWA K. AND OGATA K. (1987) Influence of external osmotic pressure on water permeability and electrical conductance of *Chara* cell membrane. *Plant Cell Physiol.* (28): 1013-1022
- KOEFOED-JOHNSEN V., USSING H. H. (1953) The contributions of diffusion and flow to the passage of D<sub>2</sub>O through living membranes. Effect of neurohypophyseal hormone on isolated anuran skin. *Acta Physiol. Scand.* (28): 60-76
- KRAJINSKI F., BIELA A., SCHUBERT D., GIANINAZZI-PEARSON V., KALDENHOFF R., FRANKEN P. (2000) Arbuscular mycorrhiza development regulates the mRNA abundance of MtAQP1 encoding a mercury insensitive aquaporin of *Medicago truncatula*. *Planta* (in Druck)
- KURIYAMA H., KAWAMOTO S., ISHIDA N., OHNO I., MITA S., MATSUZAWA Y., MATSUBARA K., OKUBO K. (1997) Molecular cloning and expression of a novel human aquaporin from adipose tissue with glycerol permeability. *Biochem. Biophys. Res. Commun.* Dec 8; (241:1): 53-58
- KUWAHARA M., GU Y., ISHIBASHI K., MARUMO F., SASAKI S. (1997) Mercury-sensitive residues and pore site in AQP3 water channel. *Biochemistry* 1997 Nov 18; 36(46): 13973-8
- KYTE J. AND DOOLITTLE R. (1982) A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* (157): 105-132
- LEVITT D. G. (1974) A new theory of transport for cell membrane pores. I. General theory and application to red cell. *Biochim. Biophys. Acta* (373): 115-131
- LIMAN E. R., TYTGAT J., HESS P. (1992) Subunit stoichiometry of a mammalian K<sup>+</sup> channel determined by construction of multimeric cDNAs. *Neuron* (9): 861-871
- LOO D. D. F., ZEUTHEN T., CHANDY G., WRIGHT E. M. (1996) Cotransport of water by Na<sup>+</sup>/glucose cotransporter. *Proceedings of the National Academy of Sciences USA* (93): 13367-13370
- LUDEVID D., HÖFTE H., HIMMELBLAU E., CHRISPPEELS M. J. (1992) The expression pattern of the tonoplast intrinsic protein TIP in *Arabidopsis thaliana* is correlated with cell enlargement. *Plant Physiology* (100): 1633-1639
- MACEY R. I. AND FARMER R. E. I. (1970) Inhibition of water and solute permeability in human red cells, *Biochim. Biophys. Acta* (211): 104-106
- MACEY R. I. (1984) Transport of water and urea in red blood cells. *Am. J. Physiol.* (246): C195-203
- MAGGIO A., JOLY R. J. (1995) Effects of mercury chloride on the hydraulic conductivity of tomato root systems. Evidence for a channel-mediated water pathway. *Plant Physiology* (109): 331-335
- MAUREL C. (1997) Aquaporins and water permeability of plant membranes. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* (48): 399-429
- MAUREL C., CHRISPPEELS M. J., LURIN C., TACNET F., GEELEN D., RIPOCHE P., GUERN J. (1997) Function and regulation of seed aquaporins. *J. Exp. Bot.* (48): 421-430
- MAUREL C., KADO R. T., GUERN J., CHRISPPEELS M. J. (1995) Phosphorylation regulates the water channel activity of the seed-specific aquaporin -TIP. *EMBO Journal* (14): 3028-35

- MAUREL C., REIZER J., SCHROEDER J. I. AND CHRISPEELS M. J. (1993) The vacuolar membrane protein -TIP creates water specific channels in *Xenopus* oocytes. *EMBO Journal* (12): 2241-2247
- MAUREL C., REIZER J., SCHROEDER J. I., CHRISPEELS M. J., SAIER M. H. JR. (1994) Functional characterization of the *Escherichia coli* glycerol facilitator, GlpF, in *Xenopus* oocytes. *J. Biol. Chem.* 269(16): 11869-11872
- MITRA A. K., VAN HOEK A. N., WIENER M. C., VERKMAN A. S., YEAGER M. (1995) The CHIP28 water channel visualized in ice by electron cristallography. *Nat. Struct. Biol.* (2): 726-29
- NACCACHE P., SHA`AFI R. I. (1974) Effect of pCMBS on water transfer across biological membranes. *Journal of Cellular Physiology* (83): 449-456
- NAKHOUL N. L., DAVIS A. D., ROMERO M. F. AND BORON W. F. (1998) Effect of expressing the water channel aquaporin-1 on the CO<sub>2</sub> permeability of *Xenopus* oocytes. *Am. J. Physiol.* 274 (*Cell Physiol.* 43): C543-C548
- OTTO B., KALDENHOFF R. (2000) Cell specific expression of the mercury-insensitive plasma membrane aquaporin NtAQP1 from *Nicotiana tabacum*. *Planta* (in Druck)
- OTTO B., GRIMM B., OTTERSBAACH P., KLOPPSTECH K. (1988) Circadian control of the accumulation of mRNA for light- and heat-inducible chloroplast proteins in pea. *Plant Physiol.*(88): 21-25
- PFEFFER W., Osmotische Untersuchungen, 2. Aufl. Leipzig 1921
- PHILLIPS A. L., HUTTLY A. K. (1994) Cloning of two gibberellin-regulated cDNAs from *Arabidopsis thaliana* by subtractive hybridisation: expression of the tonoplast water channel -TIP is increased by GA<sub>3</sub>. *Plant Mol. Biol.* (24): 603-615
- PRESTON G. M. AND AGRE P. (1991) Isolation of the cDNA for erythrocyte integral membrane protein of 28 kilodaltons: Member of an ancient channel family. *Proc. Natl. Acad. Sci USA* (88): 11110-14
- PRESTON G. M., CARROL T. P., GUGGINO W. B. AND AGRE P. (1992) Appearance of water channels in *Xenopus* oocytes expressing red cell CHIP28 protein. *Science* (256): 385-87
- PRESTON G. M., JUNG G. S., GUGGINO W. B., AGRE P. (1993) The mercury-sensitive residue at cysteine 189 in the CHIP28 water channel. *J. Biol. Chem.* (268): 17-20
- PRESTON G. M., JUNG G. S., GUGGINO W. B., AGRE P. (1994) Membrane topology of aquaporin CHIP. Analysis of functional epitope-scanning mutants by vectorial proteolysis. *J. Biol. Chem.* (269): 1668-73
- RAY P. M. (1960) On the theory of osmotic water movement. *Plant Physiol.* (35): 783-795
- REIZER J., REIZER A., SAIER M. H. J. (1993) The MIP family of integral proteins: sequence comparisons, evolutionary relationships, reconstructed pathway of evolution and proposed functional differentiation of the two repeated halves of the proteins. *Crit. Rev. Biochem. Mol. Biol.* (28): 235-57

- RIVERS R. L., DEAN R. M., CHANDY G., HALL J. E., ROBERTS D. M., ZEIDEL M. L. (1997) Functional analysis of nodulin 26, an aquaporin in soybean root nodule symbiosomes. *J. Biol. Chem.* (272:26): 16256-61
- ROBINSON D. G., SIEBER H., KAMMERLOHER W., SCHÄFFNER A. R. (1996) PIP1 aquaporins are concentrated in plasmalemmasomes of *Arabidopsis thaliana* mesophyll. *Plant Physiol* (111): 645-649
- SAMBROOK J., MANIATIS T., FRITSCH E. F. (1989) Molecular cloning. A Laboratory Manual. Second Edition, Cold Spring Harbor Laboratory New York
- SATTER R. L., GEBALLE G. T., APPLEWHITE P. B., GALSTON A. W. (1974) Potassium flux and leaf movement in *Samanea Saman*. I. Rhythmic movement. *J. Gen. Physiol.* (64): 413-430
- SCHEURING S., RINGLER P., BORGNIA M., STAHLBERG H., MÜLLER D. J., AGRE P. AND ENGEL A. (1999) High resolution AFM topographs of the *Escherichia coli* water channel aquaporin Z. *The EMBO Journal* Vol. (18) No. 18: 4981-4987
- SCHÜTZ K., TYERMAN S. D. (1997) Water channels in *Chara corallina*. *Journal of experimental Botany* (48): 1511-1518
- SHI L.-B., VERKMAN A. S. (1994) Functional independence of monomeric CHIP28 water channels revealed by expression of wild-type mutant heterodimers. *J. Biol. Chem.* (269): 10417-22
- SIDEL V. W. AND SALOMON A. K. (1957) Entrance of water into human red cells under an osmotic pressure gradient, *J. Gen. Physiol.* (41): 243-257
- SOKOLOVSKY V., KALDENHOFF R., RICCI M., RUSSO V. E. A. (1990) Fast and reliable extraction of RNA from *Neurospora crassa*. *Fungal Genetics Newsletter* (37): 40-43
- STÜRZL M., ROTH W. K. (1990) "Run off" synthesis and application of defined single-stranded DNA-hybridization probes. *Anal. Biochem.* (185): 164-169
- TAZAWA M., OHKUMA E., SHIBASAKA M., NAKASHIMA S. (1997) Mercurial-sensitive water transport in barley roots. *Journal of Plant Research* (110): 435-442
- TYERMAN S. D., BOHNERT H. J., MAUREL C., STEUDLE E. AND SMITH J. A. C. (1999) Plant aquaporins: their molecular biology, biophysics and significance for plant water relations. *Journal of Experimental Botany* (50): 1055-1071
- VAN OS C. H., DEEN P. M. T., DEMPSTER J. A. (1994) Aquaporins: water selective channels in biological membranes. Molecular structure and tissue distribution. *Biochim. Biophys. Acta* (1197): 291-309
- VERKMAN A. S., VAN HOECK A. N., MA T. FRIGERI A., SKACH W. R. (1996) Water transport across mammalian cell membranes. *Am. J. Physiol.* (270): C12-30
- WALZ T, SMITH BL, AGRE P, ENGEL A (1994) The three-dimensional structure of human erythrocyte aquaporin CHIP. *EMBO J* Jul 13 (13) 2985-93

- WALZ T., HIRAI T., MURATA K., HEYMANN J. B., MITSUOKA K., FUJIYOSHI Y., SMITH B. L., AGRE P. AND ENGEL A. (1997) The 6Å three-dimensional structure of aquaporin-1. *Nature* (387): 624-627
- WALZ T., TYPKE D., SMITH B. L., AGRE P., ENGEL A. (1995) Projection map of aquaporin-1 determined by electron cristallography. *Nat. Struct. Biol.* (2): 730-32
- WAYNE R., TAZAWA M. (1990) Nature of the water channels in the internodal cells of *Nitellopsis*. *J. Membr. Biol.* (116): 31-39
- WEI C., TYREE M. T., STEUDLE E. (1999) Direct measurement of xylem pressure in leaves of intact maize plants. A test of the cohesion-tension theory taking hydraulic architecture into consideration. *Plant Physiol.* Dec;121(4):1191-206
- WINTOUR E. M. (1997) Water channels and urea transporters. *Clin. Exp. Pharmacol. Physiol.* Jan 24(1): 1-9
- YAMADA S., KATSUHARA M., KELLY W. B., MICHALOWSKI C. B., BOHNERT H. J. (1995) A family of transcripts encoding water channel proteins: tissue-specific expression in the common ice plant. *Plant Cell* (7): 1129-1142
- YAMADA S., KOMORI T., MYERS P. N., KUWATA S., KUBO T., IMASEKI H. (1997) Expression of Plasma Membrane Water Channel Genes under Water Stress in *Nicotiana excelsior*. *Plant Cell Physiol.* 38 (11): 1226-1231
- YAMADA S., NELSON D. E., LEY E., MARQUEZ S., BOHNERT H. J. (1997b) The expression of an aquaporin promoter from *Mesembryanthemum crystallinum* in tobacco. *Plant Cell Physiology* (38): 1326-1332
- YAMAGUCHI-SHINOZAKI K., KOIZUMI M., URAO S., SHINOZAKI K. (1992) Molecular cloning and characterization of 9 cDNAs for genes that are responsive to desiccation in *Arabidopsis thaliana*. Sequence analysis of one cDNA clone that encodes a putative transmembrane channel protein. *Plant Cell Physiol.* (33): 217-224
- YAMAMOTO Y. T., TAYLOR C. G., ACEDO G. N., CHENG C. L. AND CONKLING M. A. (1991) Characterization of *cis*-acting sequences regulating root-specific gene expression in tobacco. *The Plant Cell* (3): 371-382
- YASUI M., HAZAMA A., KWON T.-H., NIELSEN S., GUGGINO WM. B., AGRE P. (1999) Rapid gating and anion permeability of an intracellular aquaporin. *Nature* Vol. 402(11): 184-187
- YOOL A. J., STAMER W. D., REGAN J. W. (1996) Forskolin stimulation of water and cation permeability in aquaporin 1 water channels. *Science* (273): 1216-1218
- ZEUTHEN T., MEINILD A.-K., KLAERKE D. A., LOO D. D. F., WRIGHT E. M., BELHAGE B. AND LITMAN T. (1997) Water transport by the Na<sup>+</sup>/glucose cotransporter under isotonic conditions, *Biology of the Cell* (89): 307-312

- ZEUTHEN T. AND KLAERKE D. A. (1999) Transport of Water and Glycerol in Aquaporin 3 Is Gated by H<sup>+</sup>. *Journal of Biol. Chem.* Vol. 274(31): 21631-21636
- ZEUTHEN T., *Molecular Mechanism of Water Transport*; Springer Verlag 1996
- ZHANG R. B., LOGEE K. A., VERKMAN A. S. (1990) Expression of mRNA coding for kidney and red cell water channels in *Xenopus* oocytes. *Journal of Cell Biol.* (265): 15375-78
- ZHANG R. B. AND VERKMAN A. S. (1991) Water and urea permeability properties of *Xenopus* oocytes: Expression of mRNA from toad urinary bladder. *Am. J. Physiol.* (260): C26-C34
- ZHANG W. H., TYERMAN S. D. (1991) Effect of low O<sub>2</sub> concentration and azide on hydraulic conductivity and osmotic volume of cortical cells of wheat roots. *Australian Journal of Plant Physiology* (18): 603-613