

## 6 Literaturverzeichnis

- Ackman, R. G., Hansen, R. P. (1967). The occurrence of diastereomers of phytanic and pristanic acids and their determination by gas-liquid chromatography. *Lipids* **2**, 357 - 362.
- Adams, S. S., Bresloff, P. & Mason, C. G. (1976). Pharmacological differences between the optical isomers of ibuprofen: evidence for metabolic inversion of the (R)-isomer. *J Pharm Pharmacol* **28**(3), 256-7.
- Albertini, M., Rehling, P., Erdmann, R., Girzalsky, W., Kiel, J. A., Veenhuis, M. & Kunau, W. H. (1997). Pex14p, a peroxisomal membrane protein binding both receptors of the two PTS-dependent import pathways. *Cell* **89**(1), 83-92.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990). Basic local alignment search tool. *J Mol Biol* **215**(3), 403-10.
- Asselineau, J. (1966). *The Bacterial Lipids*, pp. 372. Hermann-Editeurs des Sciences et des Arts. Paris.
- Avigan, J., Steinberg, D., Gutman, A., Mize, C. E. & Milne, G. W. (1966). Alpha-decarboxylation, an important pathway for degradation of phytanic acid in animals. *Biochem Biophys Res Commun* **24**(6), 838-44.
- Baes, M., Gressens, P., Baumgart, E., Carmeliet, P., Casteels, M., Fransen, M., Evrard, P., Fahimi, D., Declercq, P. E., Collen, D., van Veldhoven, P. P. & Mannaerts, G. P. (1997). A mouse model for Zellweger syndrome. *Nat Genet* **17**(1), 49-57.
- Bandlow, W., Strobel, G. & Schricker, R. (1998). Influence of N-terminal sequence variation on the sorting of major adenylate kinase to the mitochondrial intermembrane space in yeast. *Biochem J* **329**(Pt 2), 359-67.
- Banerjee, A., Burg, J., Conzelmann, E., Carroll, M. & Sandhoff, K. (1984). Enzyme-linked immunosorbent assay for the ganglioside GM2-activator protein. Screening of normal human tissues and body fluids, of tissues of GM2 gangliosidosis, and for its subcellular localization. *Hoppe Seylers Z Physiol Chem* **365**(3), 347-56.
- Batta, A. K., Salen, G., Shefer, S., Dayal, B. & Tint, G. S. (1983). Configuration at C-25 in 3 alpha, 7 alpha, 12 alpha-trihydroxy-5 beta-cholestane-26-oic acid isolated from human bile. *J Lipid Res* **24**(1), 94-6.
- Beard, M. E., Baker, R., Conomos, P., Pugatch, D. & Holtzman, E. (1985). Oxidation of oxalate and polyamines by rat peroxisomes. *J Histochem Cytochem* **33**(5), 460-4.
- Bohme, H. J., Kopperschlager, G., Schulz, J. & Hofmann, E. (1972). Affinity chromatography of phosphofructokinase using Cibacron blue F3G-A. *J Chromatogr* **69**(1), 209-14.
- Bout, A., Teunissen, Y., Hashimoto, T., Benne, R. & Tager, J. M. (1988). Nucleotide sequence of human peroxisomal 3-oxoacyl-CoA thiolase. *Nucleic Acids Res* **16**(21), 10369.
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**, 248-54.
- Braverman, N., Steel, G., Obie, C., Moser, A., Moser, H., Gould, S. J. & Valle, D. (1997). Human PEX7 encodes the peroxisomal PTS2 receptor and is responsible for rhizomelic chondrodysplasia punctata. *Nat Genet* **15**(4), 369-76.
- Brocard, C., Lametschwandtner, G., Koudelka, R. & Hartig, A. (1997). Pex14p is a member of the protein linkage map of Pex5p. *Embo J* **16**(18), 5491-500.
- Casteels, M., Croes, K., Van Veldhoven, P. P. & Mannaerts, G. P. (1997). Peroxisomal localization of alpha-oxidation in human liver. *J Inherit Metab Dis* **20**(5), 665-73.

- Catty, D., Ed. (1988). *Antibodies - A Practical Approach*. 1 edit. Vol. 1. *Antibodies*. Edited by Rickwood, D., Hames, B. D. 2 vols. Oxford: IRL Press.
- Chen, C. S., Shieh, W. R., Lu, P. H., Harriman, S. & Chen, C. Y. (1991). Metabolic stereoisomeric inversion of ibuprofen in mammals. *Biochim Biophys Acta* **1078**(3), 411-7.
- Chou, Z.-F., Chen, F. and Wilusz, J. (1994). Sequence and position requirements for uridylate-rich downstream elements of polyadenylation signals. *Nucl. Acids Res.* **22**, 2525-2531.
- Christensen, E., Van Eldere, J., Brandt, N. J., Schutgens, R. B., Wanders, R. J., Eyissen, H. J. (1990). A new peroxisomal disorder: di- and trihydroxycholestanaemia due to a presumed trihydroxycholestanyl-CoA oxidase deficiency. *J Inherit Metab Dis* **13** (3) 363-6.
- Clark, J. M. (1988). Novel non-templated nucleotide addition reactions catalyzed by prokaryotic and eucaryotic DNA polymerases. *Nucleic Acids Res* **16**(20), 9677-86.
- Contreras, M., Sengupta, T. K., Sheikh, F., Aubourg, P. & Singh, I. (1996). Topology of ATP-binding domain of adrenoleukodystrophy gene product in peroxisomes. *Arch Biochem Biophys* **334**(2), 369-79.
- Crookes, W. J. & Olsen, L. J. (1998). The effects of chaperones and the influence of protein assembly on peroxisomal protein import. *J Biol Chem* **273**(27), 17236-42.
- Danpure, C. J. (1998). The molecular basis of alanine: glyoxylate aminotransferase mistargeting: the most common single cause of primary hyperoxaluria type 1. *J Nephrol* **11** Suppl 1, 8-12.
- Danpure, C. J., Guttridge, K. M., Fryer, P., Jennings, P. R., Allsop, J. & Purdue, P. E. (1990). Subcellular distribution of hepatic alanine:glyoxylate aminotransferase in various mammalian species. *J Cell Sci* **97**(Pt 4), 669-78.
- Danpure, C. J. & Jennings, P. R. (1986). Peroxisomal alanine:glyoxylate aminotransferase deficiency in primary hyperoxaluria type I. *FEBS Lett* **201**(1), 20-4.
- De Duve, C. & Baudhuin, P. (1966). Peroxisomes (microbodies and related particles). *Physiol Rev* **46**(2), 323-57.
- de Hoop, M. J. & Ab, G. (1992). Import of proteins into peroxisomes and other microbodies. *Biochem J* **286**(Pt 3), 657-69.
- Dean, P. D. G., Johnson, W.S., Middle, F.A., Ed. (1985). *Affinity Chromatography - A Practical Approach*. Oxford: IRL Press.
- DeLabre, M. L., Nett, J. H. & Trumper, B. L. (1999). The cleaved presequence is not required for import of subunit 6 of the cytochrome bc<sub>1</sub> complex into yeast mitochondria or assembly into the complex. *FEBS Lett* **449**(2-3), 201-5.
- Devchand, P. R., Keller, H., Peters, J. M., Vazquez, M., Gonzalez, F. J. & Wahli, W. (1996). The PPARalpha-leukotriene B4 pathway to inflammation control. *Nature* **384**(6604), 39-43.
- Distel, B., Erdmann, R., Gould, S. J., Blobel, G., Crane, D. I., Cregg, J. M., Dodt, G., Fujiki, Y., Goodman, J. M., Just, W. W., Kiel, J. A., Kunau, W. H., Lazarow, P. B., Mannaerts, G. P., Moser, H. W., Osumi, T., Rachubinski, R. A., Roscher, A., Subramani, S., Tabak, H. F., Tsukamoto, T., Valle, D., van der Klei, I., van Veldhoven, P. P. & Veenhuis, M. (1996). A unified nomenclature for peroxisome biogenesis factors. *J Cell Biol* **135**(1), 1-3.
- Dreyer, C., Krey, G., Keller, H., Givel, F., Helftenbein, G. & Wahli, W. (1992). Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors. *Cell* **68**(5), 879-87.
- Eitzen, G. A., Szilard, R. K. & Rachubinski, R. A. (1997). Enlarged peroxisomes are present in oleic acid-grown *Yarrowia lipolytica* overexpressing the PEX16 gene encoding an intraperoxisomal peripheral membrane peroxin. *J Cell Biol* **137**(6), 1265-78.

- Eitzen, G. A., Titorenko, V. I., Smith, J. J., Veenhuis, M., Szilard, R. K. & Rachubinski, R. A. (1996). The Yarrowia lipolytica gene PAY5 encodes a peroxisomal integral membrane protein homologous to the mammalian peroxisome assembly factor PAF-1. *J Biol Chem* **271**(34), 20300-6.
- Elgersma, Y., Kwast, L., Klein, A., Voorn Brouwer, T., van den Berg, M., Metzig, B., America, T., Tabak, H. F. & Distel, B. (1996). The SH3 domain of the *Saccharomyces cerevisiae* peroxisomal membrane protein Pex13p functions as a docking site for Pex5p, a mobile receptor for the import PTS1-containing proteins. *J Cell Biol* **135**(1), 97-109.
- Elgersma, Y. & Tabak, H. F. (1996). Proteins involved in peroxisome biogenesis and functioning. *Biochim Biophys Acta* **1286**(3), 269-83.
- Elgersma, Y., van Roermund, C. W., Wanders, R. J. & Tabak, H. F. (1995). Peroxisomal and mitochondrial carnitine acetyltransferases of *Saccharomyces cerevisiae* are encoded by a single gene. *Embo J* **14**(14), 3472-9.
- Engel, C. & Wierenga, R. (1996). The diverse world of coenzyme A binding proteins. *Curr Opin Struct Biol* **6**(6), 790-797.
- Erdmann, R. & Blobel, G. (1995). Giant peroxisomes in oleic acid-induced *Saccharomyces cerevisiae* lacking the peroxisomal membrane protein Pmp27p. *J Cell Biol* **128**(4), 509-23.
- Erdmann, R. & Blobel, G. (1996). Identification of Pex13p a peroxisomal membrane receptor for the PTS1 recognition factor. *J Cell Biol* **135**(1), 111-21.
- Erdmann, R., Wiebel, F. F., Flessau, A., Rytka, J., Beyer, A., Frohlich, K. U. & Kunau, W. H. (1991). PAS1, a yeast gene required for peroxisome biogenesis, encodes a member of a novel family of putative ATPases. *Cell* **64**(3), 499-510.
- Fingerhut, R., Schmitz, W. & Conzelmann, E. (1993). Accumulation of phytanic acid alpha-oxidation intermediates in Zellweger fibroblasts. *J Inherit Metab Dis* **16**(3), 591-4.
- Forman, B. M., Chen, J. & Evans, R. M. (1996). The peroxisome proliferator-activated receptors: ligands and activators. *Ann N Y Acad Sci* **804**, 266-75.
- Fransen, M., Brees, C., Baumgart, E., Vanhooren, J. C., Baes, M., Mannaerts, G. P. & Van Veldhoven, P. P. (1995). Identification and characterization of the putative human peroxisomal C-terminal targeting signal import receptor. *J Biol Chem* **270**(13), 7731-6.
- Fujiki, Y. (1997). Molecular defects in genetic diseases of peroxisomes. *Biochim Biophys Acta* **1361**(3), 235-50.
- Fujiki, Y., Fowler, S., Shio, H., Hubbard, A. L. & Lazarow, P. B. (1982). Polypeptide and phospholipid composition of the membrane of rat liver peroxisomes: comparison with endoplasmic reticulum and mitochondrial membranes. *J Cell Biol* **93**(1), 103-10.
- Fujiki, Y., Rachubinski, R. A., Mortensen, R. M. & Lazarow, P. B. (1985). Synthesis of 3-ketoacyl-CoA thiolase of rat liver peroxisomes on free polyribosomes as a larger precursor. Induction of thiolase mRNA activity by clofibrate. *Biochem J* **226**(3), 697-704.
- Garcia, P. D. & Walter, P. (1988). Full-length prepro-alpha-factor can be translocated across the mammalian microsomal membrane only if translation has not terminated. *J Cell Biol* **106**(4), 1043-8.
- Gärtner, J., Moser, H. & Valle, D. (1992). Mutations in the 70K peroxisomal membrane protein gene in Zellweger syndrome. *Nat Genet* **1**(1), 16-23.
- Gaume, B., Klaus, C., Ungermann, C., Guiard, B., Neupert, W. & Brunner, M. (1998). Unfolding of preproteins upon import into mitochondria. *Embo J* **17**(22), 6497-507.

- Gautheret, D., Poirot, O., Lopez, F., Audic, S. and Claverie, J.-M. (1998). Alternate polyadenylation in human mRNAs: A large scale analysis by EST clustering. *Genome Res.* (8), 524-530.
- Goebl, M. & Yanagida, M. (1991). The TPR snap helix: a novel protein repeat motif from mitosis to transcription. *Trends Biochem Sci* **16**(5), 173-7.
- Goldman, D. S. (1954). Fatty acid oxidizing system of animal tissues. (VII).  $\beta$ -Ketoacetyl coenzyme A cleavage enzyme. *J. Biol. Chem.* **208**, 345-357.
- Gould, S. J., Kalish, J. E., Morrell, J. C., Bjorkman, J., Urquhart, A. J. & Crane, D. I. (1996). Pex13p is an SH3 protein of the peroxisome membrane and a docking factor for the predominantly cytoplasmic PTs1 receptor. *J Cell Biol* **135**(1), 85-95.
- Gould, S. J., Keller, G. A., Hosken, N., Wilkinson, J. & Subramani, S. (1989). A conserved tripeptide sorts proteins to peroxisomes. *J Cell Biol* **108**(5), 1657-64.
- Green, D. E., Mii, S., Maler, H.R., Boch, R.M. (1954). Fatty acid oxidizing system of animal tissues. (III). Butyryl coenzyme dehydrogenase. *J. Biol. Chem.* **206**, 1-12.
- Gustafsson, J. (1980). Biosynthesis of cholic acid in rat liver: formation of cholic acid from 3 alpha, 7 alpha, 12 alpha-trihydroxy- and 3 alpha, 7 alpha, 12 alpha, 24-tetrahydroxy-5 beta-cholestanoic acids. *Lipids* **15**(2), 113-21.
- Hanahan, D. (1983). Studies on transformation of Escherichia coli with plasmids. *J Mol Biol* **166**(4), 557-80.
- Holtta, E. (1977). Oxidation of spermidine and spermine in rat liver: purification and properties of polyamine oxidase. *Biochemistry* **16**(1), 91-100.
- Horng, J. T., Behari, R., Burke, L. E. & Baker, A. (1995). Investigation of the energy requirement and targeting signal for the import of glycolate oxidase into glyoxysomes. *Eur J Biochem* **230**(1), 157-63.
- Hübl, P., Bretschneider, R. (1964). Titanyl sulfate method for the detection of catalase in blood, serum and urine. *Hoppe-Seyler's Z. Physiol. Chem.* **335**, 146 - 155.
- Huhse, B., Rehling, P., Albertini, M., Blank, L., Meller, K. & Kunau, W. H. (1998). Pex17p of *Saccharomyces cerevisiae* is a novel peroxin and component of the peroxisomal protein translocation machinery. *J Cell Biol* **140**(1), 49-60.
- Imanaka, T., Small, G. M. & Lazarow, P. B. (1987). Translocation of acyl-CoA oxidase into peroxisomes requires ATP hydrolysis but not a membrane potential. *J Cell Biol* **105**(6 Pt 2), 2915-22.
- Issemann, I. & Green, S. (1990). Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* **347**(6294), 645-50.
- Jacob, J. (1976). *Chemistry and Biochemistry of Natural Waxes*, pp. 93-146. Kolattukudy, P.E., Elsevier. Amsterdam
- Johnson, G. (1983). Gel Sieving Electrophoresis: A description of procedures and analysis of errors. *Methods in Biochemical Analysis* (Glick, D., ed.), Vol. 29, pp. 25-58. J. Wiley & Sons, New York.
- Kaiser, D. G., Vangiessen, G. J., Reischer, R. J. & Wechter, W. J. (1976). Isomeric inversion of ibuprofen (R)-enantiomer in humans. *J Pharm Sci* **65**(2), 269-73.
- Kamijo, K., Taketani, S., Yokota, S., Osumi, T. & Hashimoto, T. (1990). The 70-kDa peroxisomal membrane protein is a member of the Mdr (P-glycoprotein)-related ATP-binding protein superfamily. *J Biol Chem* **265**(8), 4534-40.

- Kase, B. F. (1989). Role of liver peroxisomes in bile acid formation: inborn error of C27-steroid side chain cleavage in peroxisome deficiency (Zellweger syndrome). *Scand J Clin Lab Invest* **49**(1), 1-10.
- Kennedy, E. P. u. L., A.L. (1949). Oxidation of fatty acids and tricarboxylic acid cycle intermediates by isolated liver mitochondria. *J. Biol. Chem.* **179**, 957-972.
- Kliewer, S. A., Forman, B. M., Blumberg, B., Ong, E. S., Borgmeyer, U., Mangelsdorf, D. J., Umesono, K. & Evans, R. M. (1994). Differential expression and activation of a family of murine peroxisome proliferator-activated receptors. *Proc Natl Acad Sci U S A* **91**(15), 7355-9.
- Kliewer, S. A., Lenhard, J. M., Willson, T. M., Patel, I., Morris, D. C. & Lehmann, J. M. (1995). A prostaglandin J2 metabolite binds peroxisome proliferator-activated receptor gamma and promotes adipocyte differentiation. *Cell* **83**(5), 813-9.
- Knihinicki, R. D., Williams, K. M. & Day, R. O. (1989). Chiral inversion of 2-arylpropionic acid non-steroidal anti-inflammatory drugs--1. In vitro studies of ibuprofen and flurbiprofen. *Biochem Pharmacol* **38**(24), 4389-95.
- Knoop, F. (1904). *Der Abbau aromatischer Fettsäuren im Tierkörper*, Ernst Kuttruff, Freiburg.
- Knox, C., Sass, E., Neupert, W. & Pines, O. (1998). Import into mitochondria, folding and retrograde movement of fumarase in yeast. *J Biol Chem* **273**(40), 25587-93.
- Koike, K., Ohta, S., Urata, Y., Kagawa, Y. & Koike, M. (1988). Cloning and sequencing of cDNAs encoding alpha and beta subunits of human pyruvate dehydrogenase. *Proc Natl Acad Sci U S A* **85**(1), 41-5.
- Kolvraa, S. & Gregersen, N. (1986). In vitro studies on the oxidation of medium-chain dicarboxylic acids in rat liver. *Biochim Biophys Acta* **876**(3), 515-25.
- Kornberg, A. u. P., W.E., Jr. (1953). Enzymatic synthesis of the coenzyme A derivatives of long-chain fatty acids. *J. Biol. Chem.* **204**, 329-343.
- Kozak, M. (1986). Point mutations define a sequence flanking the AUG initiator codon that modulates translation by eukaryotic ribosomes. *Cell* **44**, 283-292.
- Kragler, F., Langeder, A., Raupachova, J., Binder, M. & Hartig, A. (1993). Two independent peroxisomal targeting signals in catalase A of *Saccharomyces cerevisiae*. *J Cell Biol* **120**(3), 665-73.
- Kunau, W. H., Beyer, A., Franken, T., Gotte, K., Marzioch, M., Saidowsky, J., Skaletz Rorowski, A. & Wiebel, F. F. (1993). Two complementary approaches to study peroxisome biogenesis in *Saccharomyces cerevisiae*: forward and reversed genetics. *Biochemie* **75**(3-4), 209-24.
- Kuroda, R., Ikenoue, T., Honsho, M., Tsujimoto, S., Mitoma, J. Y. & Ito, A. (1998). Charged amino acids at the carboxyl-terminal portions determine the intracellular locations of two isoforms of cytochrome b5. *J Biol Chem* **273**(47), 31097-102.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**(259), 680-5.
- Lametschwandtner, G., Brocard, C., Fransen, M., Van Veldhoven, P., Berger, J. & Hartig, A. (1998). The difference in recognition of terminal tripeptides as peroxisomal targeting signal 1 between yeast and human is due to different affinities of their receptor Pex5p to the cognate signal and to residues adjacent to it. *J Biol Chem* **273**(50), 33635-43.
- Lazarow, P. B. & Fujiki, Y. (1985). Biogenesis of peroxisomes. *Annu Rev Cell Biol* **1**, 489-530.
- Lee, C., Levin, A. & Branton, D. (1987). Copper staining: a five-minute protein stain for sodium dodecyl sulfate-polyacrylamide gels. *Anal Biochem* **166**(2), 308-12.

- Lipmann, F. (1945). Acetylation of sulfanilamide by liver homogenates and extracts. *J. Biol. Chem.* **160**, 173-190.
- Lough, A. K. (1973). The chemistry and biochemistry of phytanic, pristanic and related acids. *Prog Chem Fats Other Lipids* **14**(1), 1-48.
- Mahler, H. R. (1953). Role of coenzyme A in fatty acid metabolism. *Fed. Proc.* **12**, 694-702.
- Mannaerts, G. P., Van Veldhoven, P., Van Broekhoven, A., Vandebroek, G. & Debeer, L. J. (1982). Evidence that peroxisomal acyl-CoA synthetase is located at the cytoplasmic side of the peroxisomal membrane. *Biochem J* **204**(1), 17-23.
- Marshall, P. A., Dyer, J. M., Quick, M. E. & Goodman, J. M. (1996). Redox-sensitive homodimerization of Pex11p: a proposed mechanism to regulate peroxisomal division. *J Cell Biol* **135**(1), 123-37.
- Marzioch, M., Erdmann, R., Veenhuis, M. & Kunau, W. H. (1994). PAS7 encodes a novel yeast member of the WD-40 protein family essential for import of 3-oxoacyl-CoA thiolase, a PTS2-containing protein, into peroxisomes. *Embo J* **13**(20), 4908-18.
- Mazumder, R., Sasakawa, T., Kajiro, Y., Ochoa, S. (1962). Metabolism of propionic acid in animal tissues. (IX). Methylmalonyl coenzyme A racemase. *J. Biol. Chem.* **237**, 3065-3068.
- McCollum, D., Monosov, E. & Subramani, S. (1993). The pas8 mutant of *Pichia pastoris* exhibits the peroxisomal protein import deficiencies of Zellweger syndrome cells—the PAS8 protein binds to the COOH-terminal tripeptide peroxisomal targeting signal, and is a member of the TPR protein family [published erratum appears in *J Cell Biol* 1993 Sep;122(5):following 1143]. *J Cell Biol* **121**(4), 761-74.
- McNew, J. A. & Goodman, J. M. (1996). The targeting and assembly of peroxisomal proteins: some old rules do not apply. *Trends Biochem Sci* **21**(2), 54-8.
- Mihalik, S. J., Moser, H. W., Watkins, P. A., Danks, D. M., Poulos, A. & Rhead, W. J. (1989). Peroxisomal L-pipeolic acid oxidation is deficient in liver from Zellweger syndrome patients. *Pediatr Res* **25**(5), 548-52.
- Miura, S., Kasuya Arai, I., Mori, H., Miyazawa, S., Osumi, T., Hashimoto, T. & Fujiki, Y. (1992). Carboxyl-terminal consensus Ser-Lys-Leu-related tripeptide of peroxisomal proteins functions in vitro as a minimal peroxisome-targeting signal. *J Biol Chem* **267**(20), 14405-11.
- Miura, S., Mori, M., Takiguchi, M., Tatibana, M., Furuta, S., Miyazawa, S. & Hashimoto, T. (1984). Biosynthesis and intracellular transport of enzymes of peroxisomal beta-oxidation. *J Biol Chem* **259**(10), 6397-402.
- Miyazawa, S., Osumi, T., Hashimoto, T., Ohno, K., Miura, S. & Fujiki, Y. (1989). Peroxisome targeting signal of rat liver acyl-coenzyme A oxidase resides at the carboxy terminus. *Mol Cell Biol* **9**(1), 83-91.
- Moser, H. W. (1988). The peroxisome: nervous system role of a previously underrated organelle. The 1987 Robert Wartenberg lecture. *Neurology* **38**(10), 1617-27.
- Moser, H. W. (1989). Peroxisomal diseases. *Adv Pediatr* **36**, 1-38.
- Mosser, J., Douar, A. M., Sarde, C. O., Kioschis, P., Feil, R., Moser, H., Poustka, A. M., Mandel, J. L. & Aubourg, P. (1993). Putative X-linked adrenoleukodystrophy gene shares unexpected homology with ABC transporters. *Nature* **361**(6414), 726-30.
- Nakatani, T., Kawasaki, Y., Minatogawa, Y., Okuno, E. & Kido, R. (1985). Peroxisome localized human hepatic alanine-glyoxylate aminotransferase and its application to clinical diagnosis. *Clin Biochem* **18**(5), 311-6.
- Nobumoto, M., Yamada, M., Song, S., Inouye, S. & Nakazawa, A. (1998). Mechanism of mitochondrial import of adenylate kinase isozymes. *J Biochem Tokyo* **123**(1), 128-35.

- Noguchi, T., Minatogawa, Y., Takada, Y., Okuno, E. & Kido, R. (1978). Subcellular distribution of pyruvate (glyoxylate) aminotransferases in rat liver. *Biochem J* **170**(1), 173-5.
- Noguchi, T. & Takada, Y. (1979). Peroxisomal localization of alanine: glyoxylate aminotransferase in human liver. *Arch Biochem Biophys* **196**(2), 645-7.
- Nuttley, W. M., Brade, A. M., Eitzen, G. A., Veenhuis, M., Aitchison, J. D., Szilard, R. K., Glover, J. R. & Rachubinski, R. A. (1994). PAY4, a gene required for peroxisome assembly in the yeast *Yarrowia lipolytica*, encodes a novel member of a family of putative ATPases. *J Biol Chem* **269**(1), 556-66.
- O'Farrell, P. H. (1975). High resolution two-dimensional electrophoresis of proteins. *J Biol Chem* **250**(10), 4007-21.
- Oda, T., Funai, T. & Ichiyama, A. (1990). Generation from a single gene of two mRNAs that encode the mitochondrial and peroxisomal serine:pyruvate aminotransferase of rat liver. *J Biol Chem* **265**(13), 7513-9.
- Okuno, E., Minatogawa, Y., Nakanishi, J., Nakamura, M., Kamoda, N., Makino, M. & Kido, R. (1979). The subcellular distribution of alanine-glyoxylate aminotransferase and serine-pyruvate aminotransferase in dog liver. *Biochem J* **182**(3), 877-9.
- Olsen, L. J. (1998). The surprising complexity of peroxisome biogenesis. *Plant Mol Biol* **38**(1-2), 163-89.
- Ooi, C. E. & Weiss, J. (1992). Bidirectional movement of a nascent polypeptide across microsomal membranes reveals requirements for vectorial translocation of proteins. *Cell* **71**(1), 87-96.
- Osumi, T. & Fujiki, Y. (1990). Topogenesis of peroxisomal proteins. *Bioessays* **12**(5), 217-22.
- Patton, S. & Benson, A. A. (1966). Phytol metabolism in the bovine. *Biochim Biophys Acta* **125**(1), 22-32.
- Pedersen, J. I., Kase, B.F., Prydz, K., Björkhem, I. (1987). *Peroxisomes in Biology and Medicine* (Fahimi, H. D., Sies, H., Ed.), Springer-Verlag, Berlin.
- Pedersen, J. I. & Gustafsson, J. (1980). Conversion of 3 alpha, 7 alpha, 12 alpha-trihydroxy-5 beta-cholestanoic acid into cholic acid by rat liver peroxisomes. *FEBS Lett* **121**(2), 345-8.
- Pedersen, J. I., Veggan, T. & Bjorkhem, I. (1996). Substrate stereospecificity in oxidation of (25S)-3 alpha, 7 alpha, 12 alpha-trihydroxy-5 beta-cholestanyl-CoA by peroxisomal trihydroxy-5 beta-cholestanyl-CoA oxidase. *Biochem Biophys Res Commun* **224**(1), 37-42.
- Poulos, A., Sharp, P., Fellenberg, A. J. & Johnson, D. W. (1988). Accumulation of pristanic acid (2, 6, 10, 14 tetramethylpentadecanoic acid) in the plasma of patients with generalised peroxisomal dysfunction. *Eur J Pediatr* **147**(2), 143-7.
- Poulos, A., Sharp, P. & Whiting, M. (1984). Infantile Refsum's disease (phytanic acid storage disease): a variant of Zellweger's syndrome? *Clin Genet* **26**(6), 579-86.
- Prydz, K., Kase, B. F., Bjorkhem, I. & Pedersen, J. I. (1986). Formation of chenodeoxycholic acid from 3 alpha, 7 alpha-dihydroxy-5 beta-cholestanoic acid by rat liver peroxisomes. *J Lipid Res* **27**(6), 622-8.
- Purdue, P. E. & Lazarow, P. B. (1996). Targeting of human catalase to peroxisomes is dependent upon a novel COOH-terminal peroxisomal targeting sequence. *J Cell Biol* **134**(4), 849-62.
- Purdue, P. E., Lumb, M. J. & Danpure, C. J. (1992). Molecular evolution of alanine/glyoxylate aminotransferase 1 intracellular targeting. Analysis of the marmoset and rabbit genes. *Eur J Biochem* **207**(2), 757-66.

- Reichel, C., Brugger, R., Bang, H., Geisslinger, G. & Brune, K. (1997). Molecular cloning and expression of a 2-arylpropionyl-coenzyme A epimerase: a key enzyme in the inversion metabolism of ibuprofen. *Mol Pharmacol* **51**(4), 576-82.
- Rhead, W. J., Hall, C. L. & Tanaka, K. (1981). Novel tritium release assays for isovaleryl-CoA and butyryl-CoA dehydrogenases. *J Biol Chem* **256**(4), 1616-24.
- Rhodin, J. (1954). Correlation of ultrastructural organization and function in normal and experimentally changed proximal tubule cells of the mouse kidney. Akitbolaget Godvil. Karolinska Institut.
- Roels, F., Espeel, M., Pauwels, M., De Craemer, D., Egberts, H. J. & van der Spek, P. (1991). Different types of peroxisomes in human duodenal epithelium. *Gut* **32**(8), 858-65.
- Sakai, Y., Marshall, P. A., Saiganji, A., Takabe, K., Saiki, H., Kato, N. & Goodman, J. M. (1995). The *Candida boidinii* peroxisomal membrane protein Pmp30 has a role in peroxisomal proliferation and is functionally homologous to Pmp27 from *Saccharomyces cerevisiae*. *J Bacteriol* **177**(23), 6773-81.
- Sambrook, J., Fritsch, E.F., Maniatis, T., Ed. (1989). *Molecular Cloning. A Laboratory Manual*. 2 edit. 3 vols: Cold Spring Harbor Laboratory Press.
- Sandhoff, K., Conzelmann, E. & Nehrkorn, H. (1977). Specificity of human liver hexosaminidases A and B against glycosphingolipids GM2 and GA2. Purification of the enzymes by affinity chromatography employing specific elution. *Hoppe Seylers Z Physiol Chem* **358**(7), 779-87.
- Schmitz, W., Albers, C., Fingerhut, R. & Conzelmann, E. (1995). Purification and characterization of an alpha-methylacyl-CoA racemase from human liver. *Eur J Biochem* **231**(3), 815-22.
- Schmitz, W. & Conzelmann, E. (1997). Stereochemistry of peroxisomal and mitochondrial beta-oxidation of alpha-methylacyl-CoAs. *Eur J Biochem* **244**(2), 434-40.
- Schmitz, W., Fingerhut, R. & Conzelmann, E. (1994). Purification and properties of an alpha-methylacyl-CoA racemase from rat liver. *Eur J Biochem* **222**(2), 313-23.
- Schmitz, W., Helander, H. M., Hiltunen, J. K. & Conzelmann, E. (1997). Molecular cloning of cDNA species for rat and mouse liver alpha-methylacyl-CoA racemases. *Biochem J* **326**(Pt 3), 883-9.
- Schneider, H. C., Berthold, J., Bauer, M. F., Dietmeier, K., Guiard, B., Brunner, M. & Neupert, W. (1994). Mitochondrial Hsp70/MIM44 complex facilitates protein import. *Nature* **371**(6500), 768-74.
- Schneider, W. C. u. P., V.R. (1949). Intracellular distribution of enzymes. (IV). Distribution of oxalacetic oxidase activity in liver and kidney fractions. *J. Biol. Chem.* **177**, 893-903.
- Schutgens, R. B., Romeyn, G. J., Wanders, R. J., van den Bosch, H., Schrakamp, G. & Heymans, H. S. (1984). Deficiency of acyl-CoA: dihydroxyacetone phosphate acyltransferase in patients with Zellweger (cerebro-hepato-renal) syndrome. *Biochem Biophys Res Commun* **120**(1), 179-84.
- Sheets, M. D., Ogg, S.C and Wickens, M.P. (1990). Point mutations in AAUAAA and the poly (A) addition site: effects on the accuracy and efficiency of cleavage and polyadenylation *in vitro*. *Nucl. Acids Res.* **18**(19), 5799-5805.
- Shefer, S., Cheng, F. W., Batta, A. K., Dayal, B., Tint, G. S., Salen, G. & Mosbach, E. H. (1978). Stereospecific side chain hydroxylations in the biosynthesis of chenodeoxycholic acid. *J Biol Chem* **253**(18), 6386-92.
- Shieh, W. R. & Chen, C. S. (1993). Purification and characterization of novel "2-arylpropionyl-CoA epimerases" from rat liver cytosol and mitochondria. *J Biol Chem* **268**(5), 3487-93.

- Shimozawa, N., Tsukamoto, T., Suzuki, Y., Orii, T., Shirayoshi, Y., Mori, T. & Fujiki, Y. (1992). A human gene responsible for Zellweger syndrome that affects peroxisome assembly. *Science* **255**(5048), 1132-4.
- Short, J. M., Fernandez, J. M., Sorge, J. A. & Huse, W. D. (1988). Lambda ZAP: a bacteriophage lambda expression vector with in vivo excision properties. *Nucleic Acids Res* **16**(15), 7583-600.
- Siebert, P. D., Chenchik, A., Kellogg, D. E., Lukyanov, K. A. & Lukyanov, S. A. (1995). An improved PCR method for walking in uncloned genomic DNA. *Nucleic Acids Res* **23**(6), 1087-8.
- Singh, H., Usher, S., Johnson, D. & Poulos, A. (1990). A comparative study of straight chain and branched chain fatty acid oxidation in skin fibroblasts from patients with peroxisomal disorders. *J Lipid Res* **31**(2), 217-25.
- Small, G. M., Szabo, L. J. & Lazarow, P. B. (1988). Acyl-CoA oxidase contains two targeting sequences each of which can mediate protein import into peroxisomes. *Embo J* **7**(4), 1167-73.
- Soltys, B. J. & Gupta, R. S. (1999). Mitochondrial-matrix proteins at unexpected locations: are they exported? *Trends Biochem Sci* **24**(5), 174-7.
- South, S. T. & Gould, S. J. (1999). Peroxisome synthesis in the absence of preexisting peroxisomes. *J Cell Biol* **144**(2), 255-66.
- Spong, A. P. & Subramani, S. (1993). Cloning and characterization of PAS5: a gene required for peroxisome biogenesis in the methylotrophic yeast *Pichia pastoris*. *J Cell Biol* **123**(3), 535-48.
- Stabler, S. P., Marcell, P. D. & Allen, R. H. (1985). Isolation and characterization of DL-methylmalonyl-coenzyme A racemase from rat liver. *Arch Biochem Biophys* **241**(1), 252-64.
- Steinberg, D., Herndon, J. H., Jr., Uhlendorf, B. W., Mize, C. E., Avigan, J. & Milne, G. W. (1967). Refsum's disease: nature of the enzyme defect. *Science* **156** (783), 1740-2
- Steinberg, D. (1995). *Refsum disease*. 7 edit. The Metabolic and Molecular Bases of Inherited Disease (Scriver, C. R., Beaudet, A. L., Sly, W.S. and Valle, D., Ed.), 2. 3 vols, McGraw-Hill, New York.
- Subramani, S. (1993). Protein import into peroxisomes and biogenesis of the organelle. *Annu Rev Cell Biol* **9**, 445-78.
- Subramani, S. (1997). PEX genes on the rise. *Nat Genet* **15**(4), 331-3.
- Suzuki, T., Yoshida, T. & Tuboi, S. (1992). Evidence that rat liver mitochondrial and cytosolic fumarases are synthesized from one species of mRNA by alternative translational initiation at two in-phase AUG codons. *Eur J Biochem* **207**(2), 767-72.
- Suzuki, Y., Jiang, L. L., Souri, M., Miyazawa, S., Fukuda, S., Zhang, Z., Une, M., Shimozawa, N., Kondo, N., Orii, T. & Hashimoto, T. (1997). D-3-hydroxyacyl-CoA dehydratase/D-3-hydroxyacyl-CoA dehydrogenase bifunctional protein deficiency: a newly identified peroxisomal disorder. *Am J Hum Genet* **61**(5), 1153-62.
- Swell, L., Gustafsson, J., Danielsson, H., Schwartz, C. C. & Vlahcevic, Z. R. (1981). Biosynthesis of bile acids in man. An in vivo evaluation of the conversion of R and S 3 alpha, 7 alpha, 12 alpha-trihydroxy-5 beta-cholestanoic and 3 alpha, 7 alpha, 12 alpha, 12 alpha-24 xi-tetrahydroxy-5 beta-cholestanoic acids to cholic acid. *J Biol Chem* **256**(2), 912-6.

- Swinkels, B. W., Gould, S. J. & Subramani, S. (1992). Targeting efficiencies of various permutations of the consensus C-terminal tripeptide peroxisomal targeting signal. *FEBS Lett* **305**(2), 133-6.
- Takada, Y., Kaneko, N., Esumi, H., Purdue, P. E. & Danpure, C. J. (1990). Human peroxisomal L-alanine: glyoxylate aminotransferase. Evolutionary loss of a mitochondrial targeting signal by point mutation of the initiation codon. *Biochem J* **268**(2), 517-20.
- Takada, Y. & Noguchi, T. (1982). Subcellular distribution, and physical and immunological properties of hepatic alanine: glyoxylate aminotransferase isoenzymes in different mammalian species. *Comp Biochem Physiol B* **72**(4), 597-604.
- Tanford, C., Pain, R. H. & Otchin, N. S. (1967). Equilibrium and kinetics of the unfolding of lysozyme (muramidase) by guanidine hydrochloride. *J Mol Biol* **15**(2), 489-504.
- Tsukamoto, T., Shimozawa, N. & Fujiki, Y. (1994). Peroxisome assembly factor 1: nonsense mutation in a peroxisome-deficient Chinese hamster ovary cell mutant and deletion analysis. *Mol Cell Biol* **14**(8), 5458-65.
- Ungermann, C., Neupert, W. & Cyr, D. M. (1994). The role of Hsp70 in conferring unidirectionality on protein translocation into mitochondria. *Science* **266**(5188), 1250-3.
- Vamecq, J. & Draye, J. P. (1987). Interactions between the omega- and beta-oxidations of fatty acids. *J Biochem Tokyo* **102**(1), 225-34.
- van Grunsven, E. G., van Berkel, E., Ijlst, L., Vreken, P., de Klerk, J. B., Adamski, J., Lemonde, H., Clayton, P. T., Cuevas, D. A. & Wanders, R. J. (1998). Peroxisomal D-hydroxyacyl-CoA dehydrogenase deficiency: resolution of the enzyme defect and its molecular basis in bifunctional protein deficiency. *Proc Natl Acad Sci U S A* **95** (5), 2128-33.
- Van Veldhoven, P. P., Croes, K., Asselberghs, S., Herdewijn, P. & Mannaerts, G. P. (1996). Peroxisomal beta-oxidation of 2-methyl-branched acyl-CoA esters: stereospecific recognition of the 2S-methyl compounds by trihydroxycoprostanoyl-CoA oxidase and pristanoyl-CoA oxidase. *FEBS Lett* **388**(1), 80-4.
- Verhoeven, N. M., Schor, D. S., ten Brink, H. J., Wanders, R. J., Jakobs, C. (1997a) Resolution of the phytanic acid alpha-oxidation pathway: identification of pristanal as product of the decarboxylation of 2-hydroxyphytanoyl-CoA. *Biochem Biophys Res Commun* **237**(1), 33-6
- Verhoeven, N. M., Wanders, R. J., Schor, D. S., Jansen, G. A., Jakobs, C. (1997b) Phytanic acid alpha-oxidation: decarboxylation of 2-hydroxyphytanoyl-CoA to pristanic acid in human liver, *J Lipid Res* **38**(10), 2062-70
- Verhoeven, N. M., Roe, D. S., Kok, R. M., Wanders, R. J., Jakobs, C. & Roe, C. R. (1998a). Phytanic acid and pristanic acid are oxidized by sequential peroxisomal and mitochondrial reactions in cultured fibroblasts. *J Lipid Res* **39**(1), 66-74.
- Verhoeven, N. M., Wanders, R. J., Poll The, B. T., Saudubray, J. M. & Jakobs, C. (1998b). The metabolism of phytanic acid and pristanic acid in man: a review. *J Inher Metab Dis* **21**(7), 697-728.
- Wakil, S. J., Green, D.E., Mii, S., Maler, H.R. (1954). Fatty acid oxidizing system of animal tissues. (VI).  $\beta$ -Hydroxyacyl coenzyme A dehydrogenase. *J. Biol. Chem.* **207**, 631-638.
- Walton, P. A., Wendland, M., Subramani, S., Rachubinski, R. A. & Welch, W. J. (1994). Involvement of 70-kD heat-shock proteins in peroxisomal import. *J Cell Biol* **125**(5), 1037-46.

- Wanders, R. J., van Roermund, C. W., Schutgens, R. B., Barth, P. G., Heymans, H. S., van den Bosch, H. & Tager, J. M. (1990). The inborn errors of peroxisomal beta-oxidation: a review. *J Inherit Metab Dis* **13**(1), 4-36.
- Wanders, R. J., Casteels, M., Mannaerts, G. P., van Roermund, C. W., Schutgens, R. B., Kozich, V., Zeman, J., Hynek, J. (1991). Accumulation and impaired in vivo metabolism of di- and trihydroxycholestanoic acid in two patients. *Clin Chim Acta* **202**(3), 123-32.
- Waterham, H. R., de Vries, Y., Russel, K. A., Xie, W., Veenhuis, M. & Cregg, J. M. (1996). The *Pichia pastoris* PER6 gene product is a peroxisomal integral membrane protein essential for peroxisome biogenesis and has sequence similarity to the Zellweger syndrome protein PAF-1. *Mol Cell Biol* **16**(5), 2527-36.
- Watkins, P. A., Chen, W. W., Harris, C. J., Hoefler, G., Hoefler, S., Blake, D. C., Jr., Balfe, A., Kelley, R. I., Moser, A. B., Beard, M. E. & et al. (1989). Peroxisomal bifunctional enzyme deficiency. *J Clin Invest* **83**(3), 771-7.
- Wendland, M. & Subramani, S. (1993). Cytosol-dependent peroxisomal protein import in a permeabilized cell system. *J Cell Biol* **120**(3), 675-85.
- Wray, W., Boulikas, T., Wray, V. P. & Hancock, R. (1981). Silver staining of proteins in polyacrylamide gels. *Anal Biochem* **118**(1), 197-203.
- Yahraus, T., Braverman, N., Dodt, G., Kalish, J. E., Morrell, J. C., Moser, H. W., Valle, D. & Gould, S. J. (1996). The peroxisome biogenesis disorder group 4 gene, PXAAA1, encodes a cytoplasmic ATPase required for stability of the PTS1 receptor. *Embo J* **15**(12), 2914-23.
- Yang, M., Omura, S., Bonifacino, J. S. & Weissman, A. M. (1998). Novel aspects of degradation of T cell receptor subunits from the endoplasmic reticulum (ER) in T cells: importance of oligosaccharide processing, ubiquitination, and proteasome-dependent removal from ER membranes. *J Exp Med* **187**(6), 835-46.
- Yokota, S., Oda, T. & Ichiyama, A. (1987). Immunocytochemical localization of serine: pyruvate aminotransferase in peroxisomes of the human liver parenchymal cells. *Histochemistry* **87**(6), 601-6.
- Zhang, J. W. & Lazarow, P. B. (1995). PEB1 (PAS7) in *Saccharomyces cerevisiae* encodes a hydrophilic, intra-peroxisomal protein that is a member of the WD repeat family and is essential for the import of thiolase into peroxisomes. *J Cell Biol* **129**(1), 65-80.