

## Response to azacitidine is independent of p53 expression in higher-risk myelodysplastic syndromes and secondary acute myeloid leukemia

The demethylating agent azacitidine (AZA) is currently the standard of care for patients with higher-risk myelodysplastic syndromes (MDS) not eligible for allogeneic stem cell transplant (HSCT). Although approximately 50% of patients show a response to AZA it is not currently possible to accurately predict which patients will respond. In addition, higher-risk MDS frequently progress to secondary acute myeloid leukemia (sAML) within months, even in the presence of continuous therapy with AZA. Somatic gene mutations affecting prognosis have recently been identified in MDS.<sup>1,4</sup> Among these, mutations in *TP53*, *EZH2*, *ETV6*, *RUNX1* and *ASXL1* are associated with poor overall survival.<sup>1,4</sup> *TP53* mutations are mainly seen in high-risk MDS with abnormal chromosome 5 or complex karyotype.<sup>5,6</sup> In particular, *TP53* mutations have been shown to confer resistance to lenalidomide in MDS with del(5q).<sup>7</sup> In a recent publication by Saft *et al.*, p53 expression as determined by immunohistochemistry (IHC) was shown to predict for lower cytogenetic response rate, higher risk for transformation to sAML and shorter overall survival (OS) in lenalidomide-treated MDS patients with del(5q).<sup>8</sup> However, whether p53 expression also influences response to AZA is unclear. To answer this question, we analyzed the prevalence of p53 expression in a cohort of 100 patients with higher-risk MDS (IPSS intermediate-2 or high), sAML or MDS/MPN treated with at least one complete cycle of AZA (75 mg/m<sup>2</sup>/day for 7 days) and correlated this to outcome.

Bone marrow (BM) trephine biopsies were obtained from all patients before the start of AZA. Sections (2 µm) on SuperFrost microscope slides were de-paraffinized and pre-treated at 95°C for 7 min in citrate buffer (pH 6). For quantification and assessment of distribution of progenitor cells, samples were stained for CD34 (Cellmarque Rocklin, CA, USA). The DO-7 antibody (DakoCytomation, Denmark), which labels both wild-type and mutant-type p53 proteins, was used to detect p53 protein expression.<sup>6</sup> The entire trephine section was assessed for p53 protein nuclear staining in hematopoietic progenitor cells, as previously described.<sup>6,9</sup> In order to minimize the possibility of false positive results, p53 protein expression was considered positive only if strong nuclear staining (score 3+) was present in at least 5% of hematopoietic cells in the entire BM.<sup>6,9</sup> To ensure correct staining, a positive control (urothelial carcinoma) was included on each slide. Molecular *TP53* mutation analysis was performed by deep sequencing, as described previously.<sup>2</sup>

Patients' demographic and clinical characteristics are detailed in Table 1. Thirty-five patients were positive for p53 expression, 69% of whom had higher-risk MDS (Table 1). In sAML, patients lacking p53 expression were significantly more frequent (46%) than p53-positive patients (26%) ( $P=0.046$ ). Cytogenetics were scored according to IPSS-R.<sup>10</sup> P53 positive patients were predominantly found in the very poor risk group (60% p53-positive vs. 14% p53-negative;  $P<0.001$ ) (Table 1) and 74% had involvement of chromosome 5 aberrations. Five patients showed an isolated del(5q) aberration and TP53 mutations were found in 2 of these (40%). The majority of patients with monosomal karyotypes were also p53-positive (78%;  $P<0.001$ ).<sup>11</sup> The chance of exhibiting strong p53 expression was significantly increased in patients with very poor risk cytogenetics (Odds Ratio (OR): 9.333; 95%CI: 3.517; 24.771;  $P<0.001$ ) and with chromosome 5 aberrations (OR: 7.364; 95%CI:

2.886; 18.791;  $P<0.001$ ) as assessed by univariate binary logistic regression. In multivariable logistic regression analysis, a significantly increased chance of p53 expression was seen only for patients with very poor risk cytogenetics (OR: 1.488; 95%CI: 1.115; 1.986;  $P=0.007$ ).

The overall response rate (ORR) to AZA was assessed according to IWG 2006 criteria.<sup>12</sup> Complete remission in the bone marrow (BM-CR) without improvement of peripheral blood values and stable disease without hematologic

**Table 1.** Patients' demographic and clinical characteristics.

N	p53 negative patients (n = 65)	p53 positive patients (n = 35)	P
Age, years	71 (55-87)	71 (47-86)	
Median (range)			
MDS (WHO 2008)	29 (45%)	24 (69%)	
RCMD	0 (0%)	3 (9%)	
RAEB-1	4 (6%)	4 (11%)	
RAEB-2	25 (39%)	17 (49%)	
MDS/MPN (WHO 2008)	6 (9%)	2 (6%)	
sAML (WHO 2008)	30 (46%)	9 (26%)	0.046†
IPSS			
Intermediate-2	18 (28%)	13 (37%)	
High	14 (22%)	11 (31%)	
Not assessable	33 (51%)	11 (31%)	
Cytogenetics (IPSS-R)			
Very good	1 (2%)	0 (0%)	<0.001 <sup>††</sup>
Good	29 (45%)	8 (23%)	
Intermediate	14 (22%)	1 (3%)	
Poor	12 (19%)	5 (14%)	
Very poor	9 (14%)	21 (60%)	<0.001*
Very poor cytogenetics with abnormal Chr. 5	6/23 (26%)	17/23 (74%)	<0.001*
with monosomal KT	4/18 (22%)	14/18 (78%)	<0.001*
Allogeneic HSCT	7 (11%)	5 (15%)	

For comparison between patients with and without TP53 mutations a Fisher's exact (\*),  $\chi^2$  (†) or Mann-Whitney-U test (‡) was used. † $\chi^2$  test for the difference in very good to intermediate cytogenetics.

**Table 2.** Response to azacitidine.

N	p53 negative patients	p53 positive patients	P
5-Aza cycles			
Median (range)	4 (1-29)	5 (1-22)	0.926‡
All	n=65	n=35	
ORR (CR, PR, SD with HI)	16 (25%)	16 (46%)	0.033*
SD without HI	23/32 (72%)	9/32 (28%)	0.020*
MDS	n=29	n=24	
ORR (CR, PR, SD with HI)	4 (14%)	11 (46%)	0.008*
sAML	n=30	n=9	
ORR (CR, PR, SD with HI)	11 (37%)	3 (33%)	
Very poor cytogenetics			
ORR (CR, PR, SD with HI)	3/9 (33%)	7/21 (33%)	
+ abnormal Chr. 5	2/6 (33%)	7/17 (41%)	
+ monosomal KT	1/4 (25%)	5/14 (36%)	

For comparison between patients with and without TP53 mutations a Fisher's exact (\*),  $\chi^2$  (+) or Mann-Whitney-U test (‡) was used. ORR: overall response rate; CR: complete remission; PR: partial remission; SD: stable disease; HI: hematologic improvement.

**Table 3.** TP53 mutations causing false negative p53 IHC.

p53 IHC	TP53 mutation	Type of TP53 mutation
1 negative	p.Cys242Alafs*5	frameshift
2 negative	[p.Asn239*(;);p.Pro316Trpfs*22]	nonsense
3 negative	p.Gly245Ser	non-functional
4 negative	[p.Gly245Alafs*2(;)p.Gly105Valfs*23]	non-functional

IHC: immunohistochemistry.

improvement (HI) were not included in the calculation of ORR. The ORR for the total cohort was 46% for p53-positive patients and 25% for p53-negative patients ( $P=0.033$ ) (Table 2). The number of AZA cycles administered was similar in both cohorts. Of note, MDS patients with strong p53 expression showed a significantly higher ORR than p53 negative patients (46% vs. 14%;  $P=0.008$ ). Interestingly, there was no significant difference in ORR to AZA in sAML patients according to p53 status (33% ORR for p53-positive and 37% ORR for p53-negative patients). In terms of cytogenetic risk, the subgroup of patients with very poor risk cytogenetics and strong p53 expression responded as well as patients without p53 expression to treatment with AZA (33% vs. 33%, respectively). There appeared to be a better ORR in patients with very poor risk cytogenetics, chromosome 5 aberration and p53 expression than in similar patients without p53 expression (ORR 41% vs. 33%, respectively). A potentially greater ORR also appeared to be seen in patients with very poor risk cytogenetics, monosomal karyotype and p53 expression than in similar patients without p53 expression (ORR 36% vs. 25%, respectively). However, both subgroups were too small to reach statistical significance. Interestingly, the considerable subgroup of patients achieving stable disease without HI ( $n=32$ ) consisted mainly of p53-negative patients (72% vs. 28%;  $P=0.020$ ). Given the retrospective nature of the analysis, and the fact that patients were not part of a clinical trial, there was no regular response assessment during follow up.

Overall survival was measured beginning from start of AZA. Patients who underwent allogeneic HSCT were censored at the day of transplantation. The median OS of our total cohort was 380 days (95%CI: 217; 543), with 246 days (95%CI: 62; 430) for p53-positive patients and 410 days (95%CI: 272; 548) for p53-negative patients ( $P=NS$ ), confirming the very poor survival for p53-positive patients despite response to AZA.<sup>1,5,7</sup> A limitation of our study is that we were not able to evaluate response duration. Of note, compared to the randomized trial of higher-risk MDS patients treated with AZA, the ORR in our cohort was lower.<sup>15</sup> This is most likely due to the fact that our patients were older and included a higher percentage of complex karyotypes (33% poor or very poor karyotype in our p53-negative group, 74% poor or very poor karyotype in our p53-positive group compared to 28% poor risk karyotype in the AZA-001 trial). Although patients in our cohort with p53 positivity responded to AZA, response duration may be shorter than for p53-negative patients; one probable reason for the observed poor overall survival.

To validate the quality of p53 IHC to detect TP53 mutations, we performed TP53 sequencing of 37 randomly selected BM samples. TP53 sequencing revealed mutations in 14 of 37 samples and p53 IHC was positive in 10 of 37 samples. Substantial agreement (Cohens Kappa=0.645;  $P<0.001$ ) was seen between both methods. Analysis of the four TP53 mutations not detected by p53 IHC showed two

non-functional mutations: one frameshift and one nonsense mutation (Table 3). Frameshift and nonsense mutations represent approximately 6-10% and 3-8% of TP53 mutations, respectively.<sup>6,14</sup> Regarding the value of TP53 sequencing in relation to IHC for p53 expression, it is worth noting that sequencing data do not always directly reflect p53 function since not all TP53 mutations lead to loss of p53 function. In contrast, strong p53 expression is an indicator of loss of p53 function due to impairment of the auto-regulatory feedback loop inducing p53 degradation. Finally, IHC for p53 expression has recently been extensively validated for MDS and can thus be considered a reliable method to assess for dysfunctional TP53.<sup>5,7,8</sup>

In summary, we show in a large cohort of 100 patients with higher-risk MDS and sAML treated with AZA that p53 expression as a surrogate for the presence of TP53 mutations does not negatively impact on treatment response. Accordingly, the combination of AZA and lenalidomide may confer a benefit in higher-risk MDS patients with del(5q) harboring TP53 mutations and should be assessed in future trials.<sup>15</sup>

Catharina Müller-Thomas,<sup>1</sup> Martina Rudelius,<sup>2</sup> Ina-Christine Rondak,<sup>3</sup> Torsten Haferlach,<sup>4</sup> Julie Schanz,<sup>5</sup> Christina Huberle,<sup>1</sup> Burkhard Schmidt,<sup>6</sup> Rainer Blaser,<sup>3</sup> Marcus Kremer,<sup>7</sup> Christian Peschel,<sup>1,8</sup> Ulrich Germing,<sup>9</sup> Uwe Platzbecker,<sup>8,10</sup> and Katharina Götze<sup>1,8</sup>

<sup>1</sup>III. Department of Medicine, Technical University of Munich; <sup>2</sup>Institute of Pathology, Julius-Maximilians-University Würzburg; <sup>3</sup>Institute for Medical Statistics and Epidemiology, Technical University of Munich; <sup>4</sup>MLL, Munich Leukemia Laboratory; <sup>5</sup>Department of Hematology and Oncology, Georg-August-University Göttingen; <sup>6</sup>Hematology and Oncology Pasing, Munich; <sup>7</sup>Institute of Pathology Harlaching, Munich; <sup>8</sup>German Cancer Consortium (DKTK) and German Cancer Research Center (DKFZ), Heidelberg; <sup>9</sup>Department of Hematology and Oncology, Heinrich-Heine-University Düsseldorf; and <sup>10</sup>I. Department of Medicine, Gustav-Carus-University Dresden, Dresden, Germany

Correspondence: k.goetze@lrz.tum.de  
doi:10.3324/haematol.2014.104760

Key words: MDS, TP53 mutations, p53 expression, azacitidine.

Information on authorship, contributions, and financial & other disclosures was provided by the authors and is available with the online version of this article at [www.haematologica.org](http://www.haematologica.org).

## References

- Bejar R, Stevenson K, Abdel-Wahab O, Galili N, Nilsson B, Garcia-Manero G, et al. Clinical Effect of Point Mutations in Myelodysplastic Syndromes. *N Engl J Med*. 2011;364(26):2496-506.
- Haferlach T, Nagata Y, Grossmann V, Okuno Y, Bacher U, Nagae G, et al. Landscape of genetic lesions in 944 patients with myelodysplastic syndromes. *Leukemia*. 2014;28(2):241-7.
- Papaemmanuil E, Gerstung M, Malcovati L, Tauro S, Gündem G, Van Loo P, et al. Clinical and biological implications of driver mutations in myelodysplastic syndromes. *Blood*. 2013;122(22):3616-27.
- Thol F, Friesen I, Damm F, Yun H, Weissinger EM, Krauter J, et al. Prognostic Significance of ASXL1 Mutations in Patients With Myelodysplastic Syndromes. *J Clin Oncol*. 2011;29(18):2499-506.
- Kita-Sasai Y, Horiike S, Misawa S, Kaneko H, Kobayashi M, Nakao M, et al. International prognostic scoring system and TP53 mutations are independent prognostic indicators for patients with myelodysplastic syndrome. *Br J Haematol*. 2001;115(2):309-12.
- Kulasekararaj AG, Smith AE, Mian SA, Mohamedali AM, Krishnamurthy P, Lea NC, et al. TP53 mutations in myelodysplastic syndrome are strongly correlated with aberrations of chromosome 5, and correlate with adverse prognosis. *Br J Haematol*. 2013;160(5):660-72.
- Jädersten M, Saft L, Smith A, Kulasekararaj A, Pomplun S, Göhring

- G, et al. TP53 Mutations in Low-Risk Myelodysplastic Syndromes With del(5q) Predict Disease Progression. *J Clin Oncol*. 2011;29(15):1971-9.
8. Saft L, Karimi M, Ghaderi M, Matolcsy A, Mufti GJ, Kulasekararaj A, et al. P53 protein expression independently predicts outcome in patients with lower-risk myelodysplastic syndromes with del(5q). *Haematologica*. 2014;99(6):1041-9.
  9. Iwasaki T, Murakami M, Sugisaki C, Sobue S, Ohashi H, Asano H, et al. Characterization of myelodysplastic syndrome and aplastic anemia by immunostaining of p53 and hemoglobin F and karyotype analysis: Differential diagnosis between refractory anemia and aplastic anemia. *Pathol Int*. 2008;58(6):353-60.
  10. Schanz J, Tüchler H, Solé F, Mallo M, Luno E, Cervera J, et al. New Comprehensive Cytogenetic Scoring System for Primary Myelodysplastic Syndromes (MDS) and Oligoblastic Acute Myeloid Leukemia After MDS Derived From an International Database Merge. *J Clin Oncol*. 2012;30(8):820-9.
  11. Breems DA, Van Putten WLJ, De Greef GE, Van Zelder-Bhola SL, Gerssen-Schoorl KBJ, Mellink CHM, et al. Monosomal Karyotype in Acute Myeloid Leukemia: A Better Indicator of Poor Prognosis Than a Complex Karyotype. *J Clin Oncol*. 2008;26(29):4791-7.
  12. Cheson BD, Greenberg PL, Bennett JM, Lowenberg B, Wijermans PW, Nimer SD, et al. Clinical application and proposal for modification of the International Working Group (IWG) response criteria in myelodysplasia. *Blood*. 2006;108(2):419-25.
  13. Fenaux P, Mufti GJ, Hellström-Lindberg E, Santini V, Finelli C, Giagounidis A, et al. Efficacy of azacitidine compared with that of conventional care regimens in the treatment of higher-risk myelodysplastic syndromes: A randomized, open-label phase III study. *Lancet Oncol*. 2009;10:223-32.
  14. Petitjean A, Mathe E, Kato S, Ishioka C, Tavtigian SV, Hainaut P, et al. Impact of mutant p53 functional properties on TP53 mutation patterns and tumor phenotype: lessons from recent developments in the IARC TP53 database. *Hum Mutat*. 2007;28(6):622-9.
  15. Platzbecker U, Braulke F, Kündgen A, Götze K, Bug G, Schonefeldt C, et al. Sequential combination of azacitidine and lenalidomide in del(5q) higher-risk myelodysplastic syndromes or acute myeloid leukemia: a phase I study. *Leukemia*. 2013;27(6):1403-7.