

5. LITERATURVERZEICHNIS

- Achong, B. G., Mansell, P. W. A., Epstein, M. A., und Clifford, P. (1971). An unusual virus in cultures from a human nasopharyngeal carcinoma. *J. Natl. Cancer Inst.* 46, 299-307.
- Ali, M., Taylor, G. P., Pitman, R. J., Parker, D., Rethwilm, A., Cheingsong Popov, R., Weber, J. N., Bieniasz, P. D., Bradley, J., und McClure, M. O. (1996). No evidence of antibody to human foamy virus in widespread human populations. *AIDS Res Hum Retroviruses* 12, 1473-83.
- Baldwin, D. N., und Linial, M. L. (1998). The roles of Pol and Env in the assembly pathway of human foamy virus. *J Virol* 72, 3658-65.
- Baltimore, D. (1970). RNA-dependent DNA polymerase in virions of RNA tumor viruses. *Nature* 226, 1209-1211.
- Baunach, G., Maurer, B., Hahn, H., Kranz, M., und Rethwilm, A. (1993). Functional analysis of human foamy virus accessory reading frames. *J Virol* 67, 5411-8.
- Berkowitz, R., Fisher, J., und Goff, S. P. (1996). RNA packaging. *Curr. Top. Microbiol. Immunol.* 214, 177-218.
- Bieniasz, P. D., Rethwilm, A., Pitman, R., Daniel, M. D., Chrystie, I., und McClure, M. O. (1995). A comparative study of higher primate foamy viruses, including a new virus from a gorilla. *Virology* 207, 217-28.
- Birnboim, H. C., und Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acid Research* 7, 1513.
- Blair, W. S., Bogerd, H., und Cullen, B. R. (1994). Genetic analysis indicates that the human foamy virus Bel-1 protein contains a transcription activation domain of the acidic class. *J Virol* 68, 3803-8.
- Bock, M., Heinkelein, M., Lindemann, D., und Rethwilm, A. (1998). Cells expressing the human foamy virus (HFV) accessory Bet protein are resistant to productive HFV superinfection. *Virology* 250, 194-204.
- Bonifacino, J. S., Cosson, P., Shah, N., und Klausner, R. D. (1991). Role of potentially charged transmembrane residues in targeting proteins for retention and degradation within the endoplasmic reticulum. *Embo J* 10, 2783-93.
- Bonneville, J.-M., und Hohn, T. (1993). A reverse transcriptase for cauliflower mosaic virus - state of the art, 1992. In *Reverse Transcriptase*, A. M. Skalka and S. P. Goff, eds. (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press), pp. 357-390.
- Boulanger, P., und Jones, I. (1996). Use of heterologous expression systems to study retroviral morphogenesis. *Curr Top Microbiol Immunol* 214, 237-60.
- Brody, B. A., Rhee, S. S., und Hunter, E. (1994). Postassembly cleavage of a retroviral glycoprotein cytoplasmic domain removes a necessary incorporation signal and activates fusion activity. *J Virol* 68, 4620-7.
- Broussard, S. R., Comuzzie, A. G., Leighton, K. L., Leland, M. M., Whitehead, E. M., und Allan, J. S. (1997). Characterization of new simian foamy viruses from African nonhuman primates. *Virology* 237, 349-59.
- Brown, P. O. (1997). Integration. In *Retroviruses*, J. M. Coffin, S. H. Hughes und H. E. Varmus, eds. (Plainview, NY 11803: Cold Spring Harbor Laboratory Press), pp. 161-204.
- Chan, D. C., Fass, D., Berger, J. M., und Kim, P. S. (1997). Core structure of gp41 from the HIV envelope glycoprotein. *Cell* 89, 263-73.
- Chan, D. C., und Kim, P. S. (1998). HIV entry and its inhibition. *Cell* 93, 681-4.
- Chiswell, D. J., und Pringle, C. R. (1977). Infectious DNA from cells infected with feline syncytium-forming virus (Spumavirinae). *J Gen Virol* 36, 551-5.

- Cocquerel, L., Duvet, S., Meunier, J. C., Pillez, A., Cacan, R., Wychowski, C., und Dubuisson, J. (1999). The transmembrane domain of hepatitis C virus glycoprotein E1 is a signal for static retention in the endoplasmic reticulum. *J Virol* 73, 2641-9.
- Coffin, J. M. (1992a). Genetic diversity and evolution of retroviruses. In *Current Topics in Microbiology and Immunology*, pp. 143-164.
- Coffin, J. M. (1996). Retroviridae and their replication. In *Virology*, B. N. e. a. Fields, ed. (New York: Raven Press), pp. 1767-1848.
- Coffin, J. M. (1992b). Structure and classification of retroviruses. In *The Retroviridae*, J. A. Levy, ed. (New York: Plenum Press), pp. 19-49.
- Cole, S. R., Ashman, L. K., und Ev, P. L. (1987). Biotinylation: an alternative to radioiodination for the identification of cell surface antigens in immunoprecipitates. *Molecular Immunology* 24, 699-705.
- Dong, J., Roth, M. G., und Hunter, E. (1992). A chimeric avian retrovirus containing the influenza virus hemagglutinin gene has an expanded host range. *Journal of Virology* 66, 7374-7382.
- Du Bridge, R. B., Tang, P., Hsia, H. C., Leong, P. M., Miller, J. H., und Calos, M. P. (1987). Analysis of mutation in human cells by using an Epstein-Barr virus shuttle system. *Mol. Cell. Biol.* 7, 379-87.
- Dubay, J. W., Roberts, S. J., Hahn, B. J., und Hunter, E. (1992). Truncation of human immunodeficiency virus type 1 transmembrane glycoprotein cytoplasmic domain blocks virus infectivity. *Journal of Virology* 66, 6616-6625.
- Duvet, S., Cocquerel, L., Pillez, A., Cacan, R., Verbert, A., Moradpour, D., Wychowski, C., und Dubuisson, J. (1998). Hepatitis C virus glycoprotein complex localization in the endoplasmic reticulum involves a determinant for retention and not retrieval. *J Biol Chem* 273, 32088-95.
- Enders, J., und Peebles, T. (1954). Propagation in tissue culture of cytopathogenic agents from patients with measles. *Proc. Soc. Biol. Med.* 86, 277-287.
- Enssle, J., Fischer, N., Moebes, A., Mauer, B., Smola, U., und Rethwilm, A. (1997). Carboxy-terminal cleavage of the human foamy virus Gag precursor molecule is an essential step in the viral life cycle. *J-Virol* 71, 7312-7.
- Enssle, J., Jordan, I., Mauer, B., und Rethwilm, A. (1996). Foamy virus reverse transcriptase is expressed independently from the Gag protein. *Proc Natl Acad Sci U S A* 93, 4137-41.
- Enssle, J., Moebes, A., Heinkelein, M., Panhuysen, M., Mauer, B., Schweizer, M., Neumann-Haefelin, D., und Rethwilm, A. (1999). An active foamy virus integrase is required for virus replication. *J Gen Virol* 80, 1445-52.
- Fischer, N., Heinkelein, M., Lindemann, D., Enssle, J., Baum, C., Werder, E., Zentgraf, H., Muller, J. G., und Rethwilm, A. (1998). Foamy virus particle formation. *J Virol* 72, 1610-5.
- Flügel, R. M., Rethwilm, A., Maurer, B., und Darai, G. (1987). Nucleotide sequence analysis of the env gene and its flanking regions of the human spumaretrovirus reveals two novel genes. *EMBO J.* 6, 2077-2084.
- Freed, E. O., und Martin, M. A. (1996). Domains of the human immunodeficiency virus type 1 matrix and gp41 cytoplasmic tail required for envelope incorporation into virions. *J Virol* 70, 341-51.
- Freed, E. O., und Martin, M. A. (1995). Virion incorporation of envelope glycoproteins with long but not short cytoplasmic tails is blocked by specific, single amino acid substitutions in the human immunodeficiency virus type 1 matrix. *J Virol* 69, 1984-9.
- Friedmann, T., und Yee, J. K. (1995). Pseudotyped retroviral vectors for studies of human gene therapy. *Nat Med* 1, 275-7.
- Ganem, D. (1996). Hepadnaviridae and their replication. In *Virology*, B. N. Fields, D. M. Knipe und P. M. Howley, eds. (New York, NY: Lippincott-Raven), pp. 2703-2737.
- Goepfert, P., Shaw, K., Wang, G., Bansal, A., Edwards, B., und Mulligan, M. (1999). An endoplasmic reticulum retrieval signal partitions human foamy virus maturation to intracytoplasmic membranes. *J. Virol.* 73, 7210-7217.

- Goepfert, P. A., Shaw, K. L., Ritter, G. D., Jr., und Mulligan, M. J. (1997). A sorting motif localizes the foamy virus glycoprotein to the endoplasmic reticulum. *J Virol* *71*, 778-84.
- Goepfert, P. A., Wang, G., und Mulligan, M. J. (1995). Identification of an ER retrieval signal in a retroviral glycoprotein [letter]. *Cell* *82*, 543-4.
- Hahn, H., Baunach, G., Brautigam, S., Mergia, A., Neumann Haefelin, D., Daniel, M. D., McClure, M. O., und Rethwilm, A. (1994). Reactivity of primate sera to foamy virus Gag and Bet proteins. *J Gen Virol* *75*, 2635-44.
- Hallenberger, S., Bosch, V., Angliker, H., Shaw, E., Klenk, H.-D., und Garten, W. (1992). Inhibition of furin-mediated cleavage activation of HIV-1 glycoprotein gp160. *Nature* *360*, 358-361.
- Hatton, T., Zhou, S., und Standring, D. N. (1992). RNA- and DNA-binding activities in hepatitis B virus capsid protein: a model for their roles in virus replication. *Journal of Virology* *66*, 5232-5241.
- He, F., Blair, W. S., Fukushima, J., und Cullen, B. R. (1996). The human foamy virus Bel-1 transcription factor is a sequence-specific DNA binding protein. *J Virol* *70*, 3902-8.
- Heinkelein, M., Pietschmann, T., Jarmy, G., Dressler, M., Thurow, J., Imrich, H., Lindemann, D., Bock, M., Moebes, A., Roy, J., Herchenröder, O., und Rethwilm, A. (2000). Highly efficient intra-cellular retrotransposition of an exogenous primate retrovirus. Zur Publikation eingereicht.
- Helseth, E., Olshevsky, U., Gabuzda, D., Ardman, B., Haseltine, W., und Sodroski, J. (1990). Changes in the transmembrane region of the human immunodeficiency virus type 1 gp41 envelope glycoprotein affect membrane fusion. *J Virol* *64*, 6314-8.
- Herchenröder, O., Renne, R., Loncar, D., Cobb, E. K., Murthy, K. K., Schneider, J., Mergia, A., und Luciw, P. A. (1994). Isolation, cloning, and sequencing of simian foamy viruses from chimpanzees (SFVcpz): high homology to human foamy virus (HFV). *Virology* *201*, 187-99.
- Herchenröder, O., Turek, R., Neumann Haefelin, D., Rethwilm, A., und Schneider, J. (1995). Infectious proviral clones of chimpanzee foamy virus (SFVcpz) generated by long PCR reveal close functional relatedness to human foamy virus. *Virology* *214*, 685-9.
- Hosaka, M., Nagahama, M., Kim, W. S., Watanabe, T., Hatsuzawa, K., Ikemizu, J., Murakami, K., und Nakayama, K. (1991). Arg-X-Lys/Arg-Arg motif as a signal for precursor cleavage catalyzed by furin within the constitutive secretory pathway. *J. Biol. Chem.* *266*, 12127-12130.
- Hunter, E. (1994). Macromolecular interactions in the assembly of HIV and other retroviruses. *Seminars in Virology* *5*, 71-83.
- Hunter, E. (1997). Viral Entry and Receptors. In *Retroviruses*, J. M. Coffin, S. H. Hughes und V. H.E., eds. (Plainview, NY 11803: Cold Spring Harbor Laboratory Press).
- Johnson, R. H., Oginnusi, A. A., und Ladds, P. W. (1983). Isolations and serology of bovine spumavirus. *Aust Vet J* *60*, 147.
- Johnston, P. B., Dubay, J. W., und Hunter, E. (1993). Truncations of the simian immunodeficiency virus transmembrane protein confer expanded virus host range by removing a block to virus entry into cells. *J Virol* *67*, 3077-86.
- Jordan, I., Enssle, J., Guttler, E., Mauer, B., und Rethwilm, A. (1996). Expression of human foamy virus reverse transcriptase involves a spliced pol mRNA. *Virology* *224*, 314-9.
- Kemble, G. W., Danieli, T., und White, J. M. (1994). Lipid-anchored influenza hemagglutinin promotes hemifusion, not complete fusion. *Cell* *76*, 383-91.
- Konvalinka, J., Löchelt, M., Zentgraf, H., Flügel, R. M., und Krausslich, H. G. (1995). Active foamy virus proteinase is essential for virus infectivity but not for formation of a Pol polyprotein. *J Virol* *69*, 7264-8.
- Lee, P. P., und Linial, M. L. (1994). Efficient particle formation can occur if the matrix domain of human immunodeficiency virus type 1 Gag is substituted by a myristylation signal. *Journal of Virology* *68*, 6644-6654.

- Lehrmann, H., Lindemann, D., Bräutigam, S., und Rethwilm, A. Expression of foamy virus envelope proteins by recombinant baculoviruses and generation of protein-specific antisera. (submitted for publication).
- Letourneur, F., und Cosson, P. (1998). Targeting to the Endoplasmic Reticulum in Yeast Cells by Determinants Present in Transmembrane Domains. *J Biol Chem* 273, 33273-33278.
- Lindemann, D., Bock, M., Schweizer, M., und Rethwilm, A. (1997). Efficient pseudotyping of murine leukemia virus particles with chimeric human foamy virus envelope proteins. *J. Virol.* 71, 4815-20.
- Lindemann, D., und Rethwilm, A. (1998). Characterization of a human foamy virus 170-kilodalton Env-Bet fusion protein generated by alternative splicing. *J Virol* 72, 4088-94.
- Linial, M. L. (1999). Foamy viruses are unconventional retroviruses. *J Virol* 73, 1747-55.
- Löchelt, M., Muranyi, W., und Flügel, R. M. (1993). Human foamy virus genome possesses an internal, Bel-1-dependent and functional promoter. *Proc Natl Acad Sci U S A* 90, 7317-21.
- Löchelt, M., Zentgraf, H., und Flügel, R. M. (1991). Construction of an infectious DNA clone of the full-length human spumaretrovirus genome and mutagenesis of the bel 1 gene. *Virology* 184, 43-54.
- Loeb, D. D., und Ganem, D. (1993). Reverse transcription pathway of the hepatitis B virus. In Reverse transcriptase, A. M. Skalka und S. P. Goff, eds. (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press), pp. 329-355.
- Mammano, F., Kondo, E., Sodroski, J., Bukovsky, A., und Gottlinger, H. G. (1995). Rescue of human immunodeficiency virus type 1 matrix protein mutants by envelope glycoproteins with short cytoplasmic domains. *J Virol* 69, 3824-30.
- McClure, M. O., Marsh, M., und Weiss, R. A. (1988). Human immunodeficiency virus infection of CD4-bearing cells occurs by a pH-independent mechanism. *Embo J* 7, 513-8.
- McClure, M. O., Sommerfelt, M. A., Marsh, M., und Weiss, R. A. (1990). The pH independence of mammalian retrovirus infection. *J Gen Virol* 71, 767-73.
- Melikyan, G. B., White, J. M., und Cohen, F. S. (1995). GPI-anchored influenza hemagglutinin induces hemifusion to both red blood cell and planar bilayer membranes. *J Cell Biol* 131, 679-91.
- Mergia, A., und Luciw, P. A. (1991). Replication and regulation of primate foamy viruses. *Virology* 184, 475-82.
- Moebes, A., Enssle, J., Bieniasz, P. D., Heinkelein, M., Lindemann, D., Bock, M., McClure, M. O., und Rethwilm, A. (1997). Human foamy virus reverse transcription that occurs late in the viral replication cycle. *J-Virol* 71, 7305-11.
- Mulligan, M. J., Yamshchikov, G. V., Ritter, G. D., Jr., Gao, F., Jin, M. J., Nail, C. D., Spies, C. P., Hahn, B. H., und Compans, R. W. (1992). Cytoplasmic domain truncation enhances fusion activity by the exterior glycoprotein complex of human immunodeficiency virus type 2 in selected cell types. *J Virol* 66, 3971-5.
- Muranyi, W., und Flügel, R. M. (1991). Analysis of splicing patterns of human spumaretrovirus by polymerase chain reaction reveals complex RNA structures. *J Virol* 65, 727-35.
- Nassal, M. (1992). The arginine-rich domain of the hepatitis B virus core protein is required for pregenome encapsidation and productive viral positive-strand DNA synthesis but not for virus assembly. *Journal of Virology* 66, 4107-4116.
- Nassal, M., und Schaller, H. (1993). Hepatitis B virus replication. *Trends Microbiol.* 1, 221-228.
- Netzer, K. O., Rethwilm, A., Maurer, B., und ter Meulen, V. (1990). Identification of the major immunogenic structural proteins of human foamy virus. *J Gen Virol* 71, 1237-41.
- Neves, M., Peries, J., und Saib, A. (1998). Study of human foamy virus proviral integration in chronically infected murine cells. *Res-Virol* 149, 393-401.
- Odell, D., Wanas, E., Yan, J. und Ghosh, H. P. (1997). Influence of membrane anchoring and cytoplasmic domains on the fusogenic activity of vesicular stomatitis virus glycoprotein G. *J Virol* 71, 7996-8000.

- Owens, R. J., Burke, C., und Rose, J. K. (1994). Mutations in the membrane-spanning domain of the human immunodeficiency virus envelope glycoprotein that affect fusion activity. *J Virol* 68, 570-4.
- Owens, R. J., Dubay, J. W., Hunter, E., und Compans, R. W. (1991). Human immunodeficiency virus envelope protein determines the site of virus release in polarized epithelial cells. *Proc Natl Acad Sci U S A* 88, 3987-3991.
- Pear, W. S., Nolan, G. P., Scott, M. L., und Baltimore, D. (1993). Production of high-titer helper-free retroviruses by transient transfection. *Proc Natl Acad Sci U S A* 90, 8392-6.
- Perez, L. G., Davis, G. L., und Hunter, E. (1987). Mutants of the Rous sarcoma virus envelope glycoprotein that lack the transmembrane anchor and cytoplasmic domains: Analysis of intracellular transport and assembly into virions. *Journal of Virology* 61, 2981-2988.
- Pfreppe, K. I., Löchelt, M., Schnolzer, M., und Flügel, R. M. (1997). Expression and molecular characterization of an enzymatically active recombinant human spumaretrovirus protease. *Biochem-Biophys-Res-Commun* 237, 548-53.
- Pietschmann, T., Heinkelein, M., Heldmann, M., Zentgraf, H., Rethwilm, A., und Lindemann, D. (1999). Foamy virus capsids require the cognate envelope protein for particle export. *J Virol* 73, 2613-21.
- Ragheb, J. A., und Anderson, W. F. (1994a). Uncoupled expression of moloney murine leukemia virus envelope polypeptides SU and TM: a functional analysis of the role of TM domains in viral entry. *J. Virol.* 68, 3207-3219.
- Ragheb, J. A., und Anderson, W. F. (1994b). pH-independent murine leukemia virus ecotropic envelope-mediated cell fusion: implications for the role of the R peptide and p12E TM in viral entry. *J. Virol.* 68, 3220-3231.
- Redmond, S., Peters, G., und Dickson, C. (1984). Mouse mammary tumor virus can mediate cell fusion at reduced pH. *Virology* 133, 393-402.
- Rein, A., Mirro, J., Gordon Haynes, J., Ernst, S. M., und Nagashima, K. (1994). Function of the cytoplasmic domain of a retroviral transmembrane protein: p15E-p2E cleavage activates the membrane fusion capability of the murine leukemia virus env protein. *J. Virol.* 68, 1773-1781.
- Rethwilm, A. (1996). Unexpected replication pathways of foamy viruses. *J Acquir Immune Defic Syndr Hum Retrovirol* 13 Suppl 1, S248-53.
- Rethwilm, A., Baunach, G., Netzer, K. O., Maurer, B., Borisch, B., und ter Meulen, V. (1990). Infectious DNA of the human spumaretrovirus. *Nucleic Acids Res* 18, 733-8.
- Rhee, S. S., und Hunter, E. (1990). A single amino acid substitution within the matrix protein of a type D retrovirus converts its morphogenesis to that of a type C retrovirus. *Cell* 63, 77-86.
- Ritter, G. D., Jr., Mulligan, M. J., Lydy, S. L., und Compans, R. W. (1993). Cell fusion activity of the simian immunodeficiency virus envelope protein is modulated by the intracytoplasmic domain. *Virology* 197, 255-64.
- Rizvi, T. A., und Panganiban, A. T. (1992). Simian immunodeficiency virus vectors: replication and pseudotyping. *J Med Primatol* 21, 69-73.
- Rolls, M. M., Webster, P., Balba, N. H., und Rose, J. K. (1994). Novel infectious particles generated by expression of the vesicular stomatitis virus glycoprotein from a self-replicating RNA. *Cell* 79, 497-506.
- Rose, J. K., und Bergmann, J. E. (1983). Altered cytoplasmic domains affect intracellular transport of the vesicular stomatitis virus glycoprotein. *Cell* 34, 513-524.
- Salzwedel, K., Johnston, P. B., Roberts, S. J., Dubay, J. W., und Hunter, E. (1993). Expression and characterization of glycopospholipid-anchored human immunodeficiency virus type 1 envelope glycoproteins. *J Virol* 67, 5279-88.
- Sambrook, J., Fritsch, E. F., und Maniatis, T. (1989). *Molecular Cloning, A Laboratory Manual*, 2 Edition: Cold Spring Harbor Laboratory Press).
- Schliephake, A. W., und Rethwilm, A. (1994). Nuclear localization of foamy virus Gag precursor protein. *J Virol* 68, 4946-54.

- Schmidt, M., und Rethwilm, A. (1995). Replicating foamy virus-based vectors directing high level expression of foreign genes. *Virology* 210, 167-78.
- Schweizer, M., Turek, R., Hahn, H., Schliephake, A., Netzer, K. O., Eder, G., Reinhardt, M., Rethwilm, A., und Neumann Haefelin, D. (1995). Markers of foamy virus infections in monkeys, apes, and accidentally infected humans: appropriate testing fails to confirm suspected foamy virus prevalence in humans. *AIDS Res Hum Retroviruses* 11, 161-70.
- Soneoka, Y., Cannon, P. M., Ramsdale, E. E., Griffiths, J. C., Romano, G., Kingsman, S. M., und Kingsman, A. J. (1995). A transient three-plasmid expression system for the production of high titer retroviral vectors. *Nucleic Acids Res* 23, 628-33.
- Swanstrom, R., und Wills, J. W. (1997). Synthesis, Assembly, and Processing of Viral Proteins. In *Retroviruses*, J. M. Coffin, S. H. Hughes und V. H.E., eds. (Plainview, NY 11803: Cold Spring Harbor Laboratory Press), pp. 263-334.
- Temin, H. M. (1964). Nature of the provirus of Rous sarcoma. *National Cancer Institute Monographs* 17, 557-570.
- Temin, H. M., und Mizutani, S. (1970). RNA-directed DNA polymerase in virions of Rous sarcoma virus. *Nature* 226, 1211-1213.
- Tuttleman, J. S., Pourcel, C., und Summers, J. (1986). Formation of the pool of covalently closed circular viral DNA in hepadnavirus-infected cells. *Cell* 47, 451-460.
- Udenfriend, S., und Kodukula, K. (1995). How glycosylphosphatidylinositol-anchored membrane proteins are made. *Annu Rev Biochem* 64, 563-91.
- Venkatesh, L. K., Yang, C., Theodorakis, P. A., und Chinnadurai, G. (1993). Functional dissection of the human spumaretrovirus transactivator identifies distinct classes of dominant-negative mutants. *J Virol* 67, 161-9.
- Vogt, P. K. (1997). Historical Introduction to the General Properties of Retroviruses. In *Retroviruses*, J. M. Coffin, H. S. Hughes und H. E. Varmus, eds. (Plainview, NY 11803: Cold Spring Harbor Laboratory Press), pp. 1-26.
- Wang, C. T., Zhang, Y., McDermott, J., und Barklis, E. (1993). Conditional infectivity of human immunodeficiency virus matrix domain deletion mutant. *Journal of Virology* 67, 7067-7076.
- Wang, G., und Mulligan, M. J. (1999). Comparative sequence analysis and predictions for the envelope glycoproteins of foamy viruses. *J Gen Virol* 80, 245-54.
- Weissenhorn, W., Dessen, A., Calder, L. J., Harrison, S. C., Skehel, J. J., und Wiley, D. C. (1999). Structural basis for membrane fusion by enveloped viruses. *Mol Membr Biol* 16, 3-9.
- Weissenhorn, W., Dessen, A., Harrison, S. C., Skehel, J. J., und Wiley, D. C. (1997). Atomic structure of the ectodomain from HIV-1 gp41 [see comments]. *Nature* 387, 426-30.
- Weldon, R. A., und Hunter, E. (1997). Molecular Requirements for Retrovirus Assembly. In *Structural Biology of Viruses*, W. Chiu, R. M. Burnett und R. L. Garcea, eds. (New York, NY 10016: Oxford University Press), pp. 381-410.
- White, J. M. (1992). Membrane fusion. *Science* 258, 917-924.
- White, J. M. (1990). Viral and cellulare membrane fusion proteins. *Annual Review of Physiology* 52, 675-697.
- Yang, P., Zemba, M., Aboud, M., Flügel, R. M., und Löchelt, M. (1997). Deletion analysis of both the long terminal repeat and the internal promoters of the human foamy virus. *Virus Genes* 15, 17-23.
- Yee, J. K., Friedmann, T., und Burns, J. C. (1994). Generation of high-titer pseudotyped retroviral vectors with very broad host range. *Methods Cell Biol* 43, 99-112.
- Young, J. A. T., Bates, P., Willert, K., und Varmus, H. E. (1990). Efficient incorporation of human CD4 protein into avian leukosis virus particles. *Science* 250, 1421-1423.

- Yu, S. F., Stone, F. J., und Linial, M. L. (1993). Analysis of the role of the *bel* and *bet* open reading frames of human foamy virus by using a new quantitative assay. *J Virol* 67, 6618-24
- Yu, S. F., Baldwin, D. N., Gwynn, S. R., Yendapalli, S., und Linial, M. L. (1996a). Human foamy virus replication: a pathway distinct from that of retroviruses and hepadnaviruses. *Science* 271, 1579-82.
- Yu, S. F., Edelman, K., Strong, R. K., Moebes, A., Rethwilm, A., und Linial, M. L. (1996b). The carboxyl terminus of the human foamy virus Gag protein contains separable nucleic acid binding and nuclear transport domains. *J Virol* 70, 8255-62.
- Yu, S. F., Sullivan, M. D., und Linial, M. L. (1999). Evidence that the human foamy virus genome is DNA. *J-Virol* 73, 1565-72.
- Yu, X., Yuan, X., Matsuda, Z., Lee, T.-H., und Essex, M. (1992). The matrix protein of human immunodeficiency virus type 1 is required for incorporation of viral envelope protein into mature virions. *Journal of Virology* 66, 4966-4971.
- Yu, X. X., Yuan, X., McLane, M. F., Lee, T. H., und Essex, M. (1993). Mutations in the cytoplasmic domain of human immunodeficiency virus type 1 transmembrane protein impair the incorporation of Env proteins into mature virions. *Journal of Virology* 67, 213-221.