



Evolution of dispersal polymorphism and local adaptation of dispersal distance in spatially structured landscapes

Dries Bonte, Thomas Hovestadt and Hans-Joachim Poethke

D. Bonte (dries.bonte@ugent.be), Dept of Biology, Ghent Univ., K.L. Ledeganckstraat 35, BE–9000 Ghent, Belgium. – T. Hovestadt and H.-J. Poethke, Würzburg Univ., Field Station Fabriksschleichbach, Glashuettenstrasse 5, DE–96181 Raubenebrach, Germany. TH also at: Field Station Muséum Natl d'Histoire Naturelle, CNRS UMR 7179, 1 Avenue du petit château, FR–91800 Brunoy, France.

Many organisms show polymorphism in dispersal distance strategies. This variation is particularly ecological relevant if it encompasses a functional separation of short- (SDD) and long-distance dispersal (LDD). It remains, however, an open question whether both parts of the dispersal kernel are similarly affected by landscape related selection pressures.

We implemented an individual-based model to analyze the evolution of dispersal traits in fractal landscapes that vary in the proportion of habitat and its spatial configuration. Individuals are parthenogenetic with dispersal distance determined by two alleles on each individual's genome: one allele coding for the probability of global dispersal and one allele coding for the variance σ of a Gaussian local dispersal with mean value zero.

Simulations show that mean distances of local dispersal and the probability of global dispersal, increase with increasing habitat availability, but that changes in the habitat's spatial autocorrelation impose opposing selective pressure: local dispersal distances decrease and global dispersal probabilities increase with decreasing spatial autocorrelation of the available habitat. Local adaptation of local dispersal distance emerges in landscapes with less than 70% of clumped habitat.

These results demonstrate that long and short distance dispersal evolve separately according to different properties of the landscape. The landscape structure may consequently largely affect the evolution of dispersal distance strategies and the level of dispersal polymorphism.

Dispersal has important ecological and evolutionary consequences for organisms living in spatially structured populations (Kokko and López-Sepulcre 2006). It plays therefore a crucial role in the context of habitat fragmentation. Dispersal has repeatedly been shown to be a plastic process conditional on the individual's social and ecological environment (Clobert et al. 2009). Further, the presence of heritable variation and strong selection pressures related to e.g. landscape composition may induce fast evolution in dispersal traits (reviewed by Bowler and Benton 2005, Kokko and López-Sepulcre 2006, Ronce 2007). The dispersal strategy should be considered as an integrated process, encompassing dispersal propensity as well as transfer and settlement, the latter two being reflected in an individual's effective displacement distance.

Three kinds of mechanisms have been shown to favor the evolution of increased dispersal rates: kin competition (Hamilton and May 1977, Frank 1986, Taylor 1988, Ronce et al. 2000, Poethke et al. 2007), inbreeding avoidance (Waser et al. 1986, Gandon 1999) and environmental stochasticity in space and time (Van Valen 1971, Olivieri et al. 1995, Friedenberg 2003, Poethke et al. 2003). Evidently, these benefits are balanced by intrinsic or landscape-related dispersal costs (McPeck and Holt 1992, Travis and Dytham 1999, Heino and Hanski 2001, Poethke and Hovestadt 2002,

Poethke et al. 2003). Although the evolution of dispersal distance has received far less attention, the available studies (Rousset and Gandon 2002 for kin competition, Hovestadt et al. 2001 for relationship with dispersal mortality, Murrell et al. 2002 for correlations with subpopulation dynamics), point out that mechanisms that act on the evolution of emigration rates may equally affect the evolution of dispersal distance. It is therefore reasonable to assume that the spatial extent of these selective forces will affect the evolution of increased dispersal distances as well. Because dispersal propensity and dispersal distance determine gene-flow, they may strongly interfere with, and often constrain, processes of local adaptation and trait polymorphism in natural landscapes (Lenormand 2002, Billiard and Lenormand 2005, Garant et al. 2007). Thus, dispersal may be subject to local adaptation (Hovestadt et al. 2001, Gros et al. 2006) and may, at the same time, influence spatial patterns in local adaptation and its evolution.

Many organisms show polymorphism in dispersal distance strategies. Animals may disperse by different mechanisms in the larval life phase compared to the adult stage (Ghent 1999, Bonte et al. 2007). Many anemochorous (i.e. wind dispersed) plants show distinct seed dimorphism with heavy short-distance dispersing seeds and light seeds with pappus-structures for long-distance aerial dispersal (Imbert 2001, Cheptou et al. 2008), may 'exploit' different dispersal

vectors like birds with small and large home-ranges, or plants may either distribute clonally by stolones or by seeds (Gardner and Mangel 1999).

The dispersal polymorphism is of particular ecological relevance if it encompasses a functional separation of short- and long-distance dispersal (further on abbreviated as respectively SDD and LDD). Presumably, such different dispersal modes are characterised by different cost–benefit ratios due to evolutionary and environmental constraints (Bonte et al. 2006). Therefore different parts of the dispersal kernel may evolve independently (Ronce 2007, Bonte et al. 2008). For instance, long distance dispersal (i.e. dispersal beyond the bounds of the local habitat cluster) can be considered to be most advantageous in landscapes with high habitat availability or under conditions with strong enemy interactions (Hovestadt et al. 2001, Muller-Landau et al. 2003).

At large spatial scales, we can expect that polymorphisms in dispersal strategies are maintained by among-landscape variation in environmental stochasticity, habitat quality and dispersal mortality. Even at smaller spatial scales, i.e. within metapopulations, polymorphism in dispersal may evolve as a consequence of local adaptation towards local habitat configuration. This was empirically demonstrated by Hanski et al. (2004) and Bonte et al. (2006). These fast evolutionary responses towards local landscape structure are important for species conservation because they may either rescue or hasten species extinction through genetic and demographic feedbacks. This has for instance been shown by Colas et al. (1997) for an extremely rare, endemic cliff-dwelling plant species. Within single patches of habitat, recent modelling work (Gros et al. 2006) suggests that local adaptation in dispersal strategies may only evolve beyond a minimum patch size at the edges of patches.

While insights into dispersal evolution are predominantly delivered by modeling and microcosm experiments in simple landscapes, field studies highlight the importance of landscape structure for dispersal evolution (Hill et al. 1999, Hanski et al. 2004, Bonte et al. 2006, 2007, Merckx and Van Dyck 2006, Schtrickzelle et al. 2006, Cheptou et al. 2008). According to this empirical work, low dispersal distances in habitat embedded in hostile matrix (isolated habitat, edges) and high dispersal distances in continuous habitat are hypothesised to be an adaptation to the local landscape structure. Our understanding of dispersal evolution and the environmental conditions for local dispersal adaptation within realistic landscapes needs consequently to be bridged by theoretical studies. Therefore, we developed a spatially explicit, individual based model for dispersal in fractal landscapes differing in their degree of habitat availability and habitat clumping (With and King 1997). These are useful to assess the evolution of dispersal distance in a conceptual, though realistic way (With 1997). We explicitly address the question how changes of both, habitat availability and its spatial autocorrelation or contagion affect the evolution of dispersal distance polymorphism, by inferring dispersal kernels that are a function of two different dispersal functions. In contrast to earlier theoretical work (Hovestadt et al. 2001, Murell et al. 2002, Rousset and Gandon 2002), we here explored under which landscape configurations adaptive polymorphisms in dispersal distance strategy emerge.

Material and methods

Neutral landscapes

We used the program QRULE (Gardner 1999, Gardner and Urban 2007) to generate spatially structured landscapes. The generated landscape models are binary (i.e. only with a binary distinction between suitable habitat and unsuitable matrix) and measure 256×256 grid cells. For details on the landscape generator algorithm we refer to Gardner (1999). In short, we used fractal landscapes that are generated using the mid-point displacement algorithm in which both the fraction (P) and the degree of spatial autocorrelation (H) can be controlled. H represents the spatial autocorrelation or clumping of habitat, with low values indicating dispersed distribution of habitat cells; high values represent a strongly aggregated habitat distribution (With 1997). It should be noted that such fractal landscapes are always clumped and never overdispersed as compared to a pure Poisson process. Nevertheless we will use the term ‘clumped’ for highly autocorrelated (large values of H) and ‘dispersed’ for less autocorrelated (low H) landscapes. Figure 1 provides examples of four landscapes differing in clumping (H) and fraction of suitable habitat (P). Fractal landscapes provide a tool to tease apart the effects of habitat fragmentation (with high values of H indicating strong connectivity) from habitat loss (with high values of P indicating high amounts of preserved habitat). To avoid edge effects in the simulations, landscapes were wrapped at the edges into a torus so that edges matched, and can consequently be considered as being infinite but periodic landscapes.

We investigated the evolution of dispersal distance in fractal landscapes that systematically varied in P between 0.1 and 0.9 and H between 0 and 1 (Table 1). Intervals were 0.1 in both cases, subsequently resulting in 99 landscapes with different combinations of H and P. We chose to generate five replicates of 99 interdependent landscape (the landscapes are generated from the same baseline fractal landscape; Gardner 1999). Consequently, a total of 495 simulations (described hereunder) were run.

The model

Individuals

We modelled the evolution of dispersal distance in parthenogenetic organisms with discrete generations. Individuals either disperse globally with a certain probability (i.e. the individual is displaced toward a random cell in the landscape), or they draw a distance from a Gaussian probability distribution. We chose to implement this dispersal function because (1) it allows the generation of fat-tailed dispersal kernels by increased global dispersal and (2) because of its realism in many terrestrial arthropods with polymorphic dispersal strategies (Zera and Denno 1997). Dispersal costs emerge in first instance due to arrival in unsuitable matrix cells, but we alternatively tested a scenario with additional distance dependent dispersal mortality μ_{disp} . The latter is positively linearly correlated with dispersal distance according to the rule $\mu_{\text{disp}} = \text{dist} / \text{dist}_{\text{max}}$. In our landscape of 256×256 grids, dist_{max} equals 362.




Figure 1. Examples of fractal landscapes used in simulations. Upper panels contain $P = 0.2$ habitat (black), lower panels $P = 0.8$. Left panels show a more dispersed distribution ($H = 0.2$), right ones are more clumped ($H = 0.8$).

Local population dynamics

Each habitat grid cell was characterised by a carrying capacity K beyond which reproduction failed (Table 1). Reproduction takes place after dispersal, i.e. after the building-up of local population sizes (N). The realised number of offspring λ was determined as a linear function of the local density $\lambda = KN$ with complete loss of individuals through failed reproduction when $N \geq K$ in local grid cells, but with the possibility for recolonization from reproducing individual in the landscape. Individuals die immediately when they disperse into the hostile matrix. These local population dynamics resemble typical arthropod herbivores with distinct genetically determined dispersal polymorphism and scramble competition (Zera and Denno 1997).

Alternatively, we tested the model for logistic growth population dynamics (i.e. contest competition), hypothesised to plant systems. Therefore, we modified local population dynamics according to Hassell and Comins (1976). Local

population dynamics are governed by density-dependent reproduction of individuals. Each generation, a female gives birth to Λ offspring, where $\Lambda \sim \text{Pois } \lambda$. In our simulations, λ was set to 2, typical for arthropod demography (Bellows 1981). Offspring develop into mature individuals with a density-dependent survival probability s due to contest competition:

$$s = \frac{1}{(1 + aN)} \quad (1) \quad \text{with } a = \frac{\lambda - 1}{K}$$

Here N represents the local population size and K carrying capacity.

Evolution of dispersal

Dispersal distance is determined by two alleles on each individual's genome: one allele (allele G_p) coding for the probability

Table 1. Summary of the model parameters.

Parameter	Description	Ranges tested (sensitivity analysis)
K	Carrying capacity local populations	5 (7, 10, 30*)
λ	mean offspring number (logistic growth submodel)	2 (3, 4, 5)
μ	dispersal mortality (distance dependent mortality submodel)	dist/dist _{max}
P	Proportion of available habitat in the landscape	0.1, 0.2, ..., 0.9
H	Spatial autocorrelation of the available habitat	0, 0.1, 0.2, ...1

*Analysis done in grid landscape of 128×128 cells for computational reasons.

of global dispersal p_{global} . Alternatively, that is with probability $1 - p_{\text{global}}$ individuals will disperse locally according to a Gaussian dispersal kernel (mean value of 0) with the genetically determined variance (σ). The Gaussian kernel is two-dimensional with randomly selected directionality (isotropic). Thus the second allele (G_p) codes for the variance (σ) of the Gaussian distribution. Individual allele frequencies were initialised for G_p from a uniform distribution [0..1], and for G_g from a uniform distribution [1..128]. Offspring inherit both alleles from their parent, but alleles may mutate with a probability of 0.001. Mutations are uniformly sampled within the interval 0.9x to 1.1x. As a consequence, all individuals are theoretically dispersing, but philopatry (dispersal with distance zero) occurs when both G_p and G_g approach zero.

Data analysis

Stable distributions of trait values were reached within 1000 time-steps in all scenarios. Mean trait values were calculated for all surviving individuals in one landscape. Beyond the global evolutionary results we were also interested to investigate under which conditions local adaptation of dispersal traits can occur. To characterize local conditions, we estimated the proportion of habitat cells (p) in the vicinity of 1000 randomly selected habitat cells in a similar way as the developed univariate O-ring statistics by Wiegand et al. (1999). The number of suitable habitat cells in 100 increasing radii of 1 grid cell was counted for each of these randomly selected cells. In a fractal landscape the availability of habitat is expected to more or less gradually decline with distance for any cell of suitable habitat, ultimately equilibrating at the proportion of available habitat in the entire landscape (P). We use the slope α of the fitted negative exponential function $p' = e^{-(\alpha \times \text{radius})}$ to describe the change in habitat availability with distance for the individual cells, with high values representing steeply declining availability (as would be the case for isolated patches in the landscape, or patches at edges of clumped habitat cells). Intermediate values of α indicate gradually declining habitat availability, while values of α close to 0 would indicate a homogeneous habitat distribution within the landscape sector analyzed. In a region with rapidly declining habitat availability (large α) it should be beneficial to limit local dispersal to the near vicinity, i.e. selection should favour low levels of the allele G_g . We thus use the strength of the Spearman rank correlation (R_s) between the cell's value for α (a habitat attribute) and the values of the two dispersal alleles (G_p respectively G_g) of the individual residing in this cell to estimate the degree of local adaptation (or selection driven polymorphism). We assume the presence of local adaptation only when $R_s^2 > 10\%$.

Mean individual trait values and Spearman correlations between local trait values and local landscape structure were subsequently averaged over the five independent replicas for each landscape type (i.e. each landscape determined by one of the 99 H, p-values). These averaged trait values are presented in this contribution.

Results

Evolutionary stable Gaussian dispersal distances (Fig. 2A, 2C, 2E) evolve towards higher values with an increase in the

proportion of available habitat (P) as well as with an increase in spatial autocorrelation (H). The proportion of individuals performing global dispersal also increases with increasing P but declines with increasing H (Fig. 2B, 2D, 2F), i.e. selection favours the investment into local dispersal as autocorrelation becomes larger (at a given overall habitat availability). We thus observed the prevalence of more fat-tailed dispersal kernels due to increased global dispersal in landscapes characterized by more evenly distributed suitable habitat. The obtained outcome is consistent over the three different simulation experiments (Fig. 2) and does, with exception of a slight decrease in global dispersal, not depend on the applied submodels for local population dynamics or the implementation of distance dependent dispersal mortality. Neither did we detect deviations in scenarios with different λ or K (results not shown).

No local adaptation within landscapes was recorded for global dispersal trait values (all $R^2 < 0.03$). However, local adaptation in the Gaussian component of dispersal emerge when the availability of habitat P falls below 0.7 and when at the same time clumping is high (H within range 0.4-1 (Fig. 3, only data presented for simple scramble competition). These ranges of landscape structure correspond with the evolution of local dispersal distances (trait G_g) in the range of 1.5-3 units (Fig. 2A).

Discussion

The availability of habitat and its spatial distribution has a prominent effect on the evolution of dispersal distances. We show that an increasing proportion of suitable habitat in a landscape selects – unsurprisingly – for increased dispersal distance, as reflected by the increased investments into global dispersal and the parallel increase in the local dispersal distances. Yet a change in autocorrelation inflicts opposing selective pressure on the local and global component of dispersal. This gives rise to particularly fat-tailed dispersal kernels under conditions of spatially uncorrelated habitat availability (large fraction of global dispersal but short mean dispersal distance for local dispersal). Polymorphism in dispersal tactics only emerged in landscapes with low habitat availability and high spatial autocorrelation and emerged only with respect to the trait coding for the Gaussian dispersal tactics. This local adaptation emerged in landscapes that select against global dispersal and for intermediate Gaussian dispersal rates, i.e. in landscapes with less than 70% but strongly clumped habitat.

The retrieved patterns are obviously relevant for organisms with mixed dispersal strategies encompassing (1) more or less controlled short distance dispersal (SDD) and (2) uncontrolled passive long distance dispersal (LDD). These conditions hold for seed dimorphic plants with investments for anemochorous dispersal, insects with facultative wing formation like aphids and some butterflies, arthropods that use silk as a dispersal mode and probably many organisms that combine passive, uncontrolled dispersal phases with more active dispersal modes. Neither the implementation of local logistic growth with contest competition nor the distance dependent dispersal mortality affected this pattern. Because we implemented asexual reproduction and low local carrying capacities, strong




Figure 2. Mean values for dispersal traits according to the proportion of available habitat (P) and its degree of autocorrelation (H). (A), (C), (E): mean value of the allele for the Gaussian distance, (B), (D), (F): frequency of individuals performing global dispersal. We present data for respectively the baseline smodel (scramble competition; A, B), logistic growth model (C, D) and baseline model with distance dependent dispersal mortality (E,F).




Figure 3. A. Strength of local adaptation (R^2 , see material and methods for details) of Gaussian dispersal distance strategies according to the proportion of available habitat (P) and its degree of autocorrelation (H).

kin competition can here be attributed to overrule the expected decreasing dispersal under logistic growth or additional dispersal mortality (Hamilton and May 1977, Ronce et al. 2000, Poethke et al. 2007).

The few empirical studies that detected a geographic variation in dispersal tactics confirm our theoretical predictions that (passive) LDD would be advantageous in landscapes with high degrees of habitat availability (Cody and Overton 1996, Imbert 2001, Bonte et al. 2006, 2007, Cheptou et al. 2008), but no evidence for shifts towards more controlled movement methods in highly connective landscapes were found. To our knowledge, no empirical studies have explicitly tested the relationship between spatial configuration and the evolution of dispersal polymorphism within heterogeneous landscapes. However, if we consider detectable movements of butterflies as short distance dispersal that contrast with probable, but rarely quantified long distance dispersal (Van Dyck and Baguette 2004), patterns found in a meadow-inhabiting species (Baguette and Schtickzelle 2006) fit our predictions.

Because our model deals with the evolution of ecological phenomena typical for passively dispersing organisms, we hypothesised the polymorphism to be adaptive within a landscape if distances decrease with increased habitat isolation and availability (Bonte et al. 2007, Cheptou et al. 2008). Evidently, different responses can be expected for actively moving organisms with certain habitat detection abilities (Lima and Zollner 1996). In first instance, our simulations point out the absence of local adaptation in LDD dispersal strategies. This is not unexpected given the overall low dispersal rates, rendering the trait more subject to drift than to selection, and the simultaneous gene flow over very large distances associated with this mode of dispersal. In contrast, local adaptation in Gaussian dispersal distance was detected in landscapes with high degrees of autocorrelation and habitat availability up to 70%. These comprise landscapes that on average select for intermediate dispersal distances and accord with earlier theoretical work on the relationship between gene flow and local adaptation

(Lenormand 2002, Billiard and Lenormand 2005, Garant et al. 2007). Because dispersal distance is expected to be tightly related to gene-flow, extremely low dispersal distances will lead to the absence of gene flow and consequently low genetic variation on which natural selection can act. In contrast, high levels of gene flow through large dispersal distances will be responsible for gene swamping. High dispersal distances will consequently erode any emergence of local adaptation in dispersal distance.

By modelling combined dispersal kernels, we show that different parts of the kernel (i.e. SDD and LDD) are affected by different landscape-related selection pressures. Both SDD and LDD consequently evolve independently, despite the simplicity of the applied simulations. Moreover, it can be expected that the consideration of (1) different associated dispersal costs and (2) additional environmental attributes like different rates and spatial extents of disturbance (Muller-Landau et al. 2003) will presumably amplify these independent evolutionary mechanisms.

Given the relevance of LDD for biological invasions (Muller-Landau et al. 2003) and range expansion under climate change (Phillips et al. 2008), we can expect serious geographic variation in range expansion rates for species with distinct dispersal polymorphisms. This geographic variation is expected to be related to the landscape configuration of source populations with larger range expansion fronts for those populations, inhabiting landscapes with high levels of naturally or already long-term fragmented habitat. Evolutionary responses of dispersal kernels at range expansion fronts are documented (Travis and Dytham 2002, Phillips et al. 2008). Our simulations additionally point out that the underlying landscape structure should be equally considered when predicting invasive range expansion of species under global change.

Acknowledgements - DB received a mobility grant from the Fund for Scientific Research – Flanders (FWO) for a long-term stay at Würzburg Univ. DB was funded by FWO research grant (G.0057.09). HJP and TH are partially supported by a grant from the “Deutsche Forschungsgemeinschaft” (DFG PO244/4).

References

- Baguette, M. and Schtickzelle, N. 2006. Negative relationship between dispersal distance and demography in butterfly metapopulations. – *Ecology* 87: 648–654.
- Bellows, T. S. 1981. The descriptive properties of some models for density dependence. – *J. Anim. Ecol.* 50: 139–156.
- Billiard, S. and Lenormand, T. 2005. Evolution of migration under kin selection and local adaptation. – *Evolution* 59: 13–23.
- Bonte, D. et al. 2006. Geographic variation in wolfspider dispersal behaviour is related to landscape structure. – *Anim. Behav.* 72: 655–662.
- Bonte, D. et al. 2007. Aerial dispersal plasticity under different wind velocities in a salt marsh wolf spider. – *Behav. Ecol.* 18: 438–443.
- Bonte, D. et al. 2008. Thermal conditions during juvenile development affect adult dispersal in a spider. – *Proc. Natl Acad. Sci. USA* 105: 17000–17005.
- Bowler, D. E. and Benton, T. G. 2005. Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. – *Biol. Rev.* 80: 205–225.

- Cheptou, P. O. et al. 2008. Rapid evolution of seed dispersal in an urban environment in the weed *Crepis sancta*. – Proc. Natl Acad. Sci. USA 105: 3796–3799.
- Clobert, J. et al. 2009. Informed dispersal, heterogeneity in animal dispersal syndromes and the dynamics of spatially structured populations. – Ecol. Lett. 12: 197–209.
- Cody, M. L. and Overton, J. M. 1996. Short-term evolution of reduced dispersal in island plant populations. – J. Ecol. 84: 53–61.
- Colas, B. et al. 1997. *Centaurea corymbosa*, a cliff-dwelling species tottering on the brink of extinction: a demographic and genetic study. – Proc. Natl Acad. Sci. USA 94: 3471–3476.
- Frank, S. A. 1986. Dispersal polymorphism in subdivided populations. – J. Theor. Biol. 122: 303–309.
- Friedenberg, N. A. 2003. Experimental evolution of dispersal in spatiotemporally variable microcosms. – Ecol. Lett. 6: 953–959.
- Gandon, S. 1999. Kin competition, the cost of inbreeding and the evolution of dispersal. – J. Theor. Biol. 200: 345–364.
- Garant, D. et al. 2007. The multifarious effects of dispersal and gene flow on contemporary adaptation. – Funct. Ecol. 21: 434–443.
- Gardner, R. H. 1999. RULE: map generation and spatial analysis program. – In: Klopatek, J. M. and Gardner, R. H. (eds), Landscape ecological analysis: issues and applications. Springer, pp. 280–303.
- Gardner, R. H. and Urban, D. L. 2007. Neutral models for testing landscape hypotheses. – Landscape Ecol. 22: 15–29.
- Gardner, S. N. and Mangel, M. 1999. Modeling investments in seeds, clonal offspring, and translocation in a clonal plant. – Ecology 80: 1202–1220.
- Ghent, A. W. 1999. Studies of ballooning and resulting patterns of locally contagious distribution of the bagworm *Thyridopteryx ephemeriformis* (Haworth) (Lepidoptera: Psychidae). – Am. Midl. Nat. 142: 291–313.
- Gros, A. et al. 2006. Evolution of spatial pattern in dispersal strategies. – Oikos 114: 544–552.
- Hamilton, W. D. and May, R. M. 1977. Dispersal in stable habitats. – Nature 269: 578–581.
- Hanski, I. et al. 2004. Variation in migration propensity among individuals maintained by landscape structure. – Ecol. Lett. 7: 958–966.
- Hassell, M. P. and Comins, H. N. 1976. Discrete time models for two-species competition. – Theor. Popul. Biol. 9: 202–221.
- Heino, M. and Hanski, I. 2001. Evolution of migration rate in a spatially realistic metapopulation model. – Am. Nat. 157: 495–511.
- Hill, J. K. et al. 1999. Flight morphology in fragmented populations of a rare British butterfly, *Hesperia comma*. – Biol. Conserv. 87: 277–283.
- Hovestadt, T. et al. 2001. Evolution of reduced dispersal mortality and ‘fat-tailed’ dispersal kernels in autocorrelated landscapes. – Proc. R. Soc. Lond. B 268: 385–391.
- Imbert, E. 2001. Capitulum characteristics in a seed heteromorphic plant, *Crepis sancta* (Asteraceae): variance partitioning and inference for the evolution of dispersal rate. – Heredity 86: 78–86.
- Kokko, H. and Lopez-Sepulcre, A. 2006. From individual dispersal to species ranges: perspectives for a changing world. – Science 313: 789–791.
- Lenormand, T. 2002. Gene flow and the limits to natural selection. – Trends Ecol. Evol. 17: 183–189.
- Lima S. L. and Zollner, P. A. 1996. Towards a behavioral ecology of ecological landscapes. – Trends Ecol. Evol. 11: 131–135.
- McPeck, M. A. and Holt, R. D. 1992. The evolution of dispersal in spatially and temporally varying environments. – Am. Nat. 140: 1010–1027.
- Merckx, T. and van Dyck, H. 2007. Habitat fragmentation affects habitat-finding ability of the speckled wood butterfly, *Pararge aegeria* L. – Anim. Behav. 74: 1029–1037.
- Muller-Landau, H. C. et al. 2003. Theoretical perspectives on evolution of long-distance dispersal and the example of specialized pests. – Ecology 84: 1957–1967.
- Murrell, D. J. et al. 2002. The evolution of dispersal distance in spatially-structured populations. – Oikos 97: 229–236.
- Olivieri, I. et al. 1995. Metapopulation genetics and the evolution of dispersal. – Am. Nat. 146: 202–228.
- Phillips, B. L. et al. 2008. Reid’s paradox revisited: the evolution of dispersal kernels during range expansion. – Am. Nat. 172: 34–48.
- Poethke, H. J. and Hovestadt, T. 2002. Evolution of density- and patch-size-dependent dispersal rates. – Proc. R. Soc. Lond. B 269: 637–645.
- Poethke, H. J. et al. 2003. Local extinction and the evolution of dispersal rates: causes and correlations. – Am. Nat. 161: 631–640.
- Poethke, H. J. et al. 2007. The relative contributions of individual- and kin-selection in the evolution of density-dependent dispersal rates. – Evol. Ecol. Res. 9: 41–50.
- Ronce, O. 2007. How does it feel to be like a Rolling Stone? Ten questions about dispersal evolution. – Annu. Rev. Ecol. Evol. Syst. 38: 231–253.
- Ronce, O. et al. 2000. Kin selection and natal dispersal in an age structured population. – Theor. Popul. Biol. 58: 143–159.
- Rousset, F. and Gandon, S. 2002. Evolution of the distribution of dispersal distance under distance-dependent cost of dispersal. – J. Evol. Biol. 15: 515–523.
- Schickzelle, N. et al. 2006. Dispersal depression with habitat fragmentation in the bog fritillary butterfly. – Ecology 87: 1057–1065.
- Taylor, P. D. 1988. An inclusive fitness model for dispersal of offspring. – J. Theor. Biol. 130: 363–378.
- Travis, J. M. J. and Dytham, C. 1999. Habitat persistence, habitat availability and the evolution of dispersal. – Proc. R. Soc. Lond. B 266: 723–728.
- Travis, J. M. J. and Dytham, C. 2002. Dispersal evolution during invasions. – Evol. Ecol. Res. 4: 1119–1129.
- van Dyck, H. and Baguette, M. 2005. Dispersal behaviour in fragmented landscapes: routine or special movements? – Basic Appl. Ecol. 6: 535–545.
- van Valen L. 1971. Group selection and the evolution of dispersal. – Evolution 25: 591–598.
- Waser, P. W. et al. 1986. When should animals tolerate inbreeding? – Am. Nat. 128: 529–537.
- Wiegand, T. et al. 1999. Finding the missing link between landscape structure and population dynamics: a spatially explicit perspective. – Am. Nat. 154: 605–627.
- With, K. A. 1997. The application of neutral landscape models in conservation biology. – Conserv. Biol. 11: 1069–1080.
- With, K. A. and King, A. W. 1997. The use and misuse of neutral landscape models in ecology. – Oikos 79:219–229.
- Zera, A. J. and Denno, R. F. 1997. Physiology and ecology of dispersal polymorphism in insects. – Annu. Rev. Entomol. 42: 207–230.